

MAGNESIUM SALTS VOL I #62

D18

Magnesium Salts-Volume I

#62

9/21/73

VOLUME 1

**GRAS MONOGRAPH SERIES**

**MAGNESIUM SALTS**

prepared for  
**THE FOOD AND DRUG ADMINISTRATION**  
**DEPARTMENT OF HEALTH, EDUCATION**  
**AND WELFARE**

**SEPTEMBER 21, 1973**

prepared by  
**Tracor Jitco, Inc.**

## Table of Contents

### Summary

### CHEMICAL INFORMATION

MAGNESIUM CARBONATE		Page
I.	Nomenclature	1
II.	Empirical Formula	1
III.	Structural Formula	1
IV.	Molecular Weight	1
V.	Specifications	1
VI.	Description	4
MAGNESIUM CHLORIDE		
I.	Nomenclature	5
II.	Empirical Formula	5
III.	Structural Formula	5
IV.	Molecular Weight	5
V.	Specifications	5
VI.	Description	7
MAGNESIUM HYDROXIDE		
I.	Nomenclature	8
II.	Empirical Formula	8
III.	Structural Formula	8
IV.	Molecular Weight	8
V.	Specifications	8
VI.	Description	11
MAGNESIUM OXIDE		
I.	Nomenclature	12
II.	Empirical Formula	12
III.	Structural Formula	12
IV.	Molecular Weight	12
V.	Specifications	12
VI.	Description	15

## Table of Contents (Cont'd)

MAGNESIUM PHOSPHATE, DIABASIC		Page
I.	Nomenclature	16
II.	Empirical Formula	16
III.	Structural Formula	16
IV.	Molecular Weight	16
V.	Specifications	16
VI.	Description	19
MAGNESIUM PHOSPHATE, TRIBASIC		
I.	Nomenclature	20
II.	Empirical Formula	20
III.	Structural Formula	20
IV.	Molecular Weight	20
V.	Specifications	20
VI.	Description	23
MAGNESIUM STEARATE		
I.	Nomenclature	24
II.	Empirical Formula	24
III.	Structural Formula	24
IV.	Molecular Weight	24
V.	Specifications	24
VI.	Description	28
MAGNESIUM SULFATE		
I.	Nomenclature	29
II.	Empirical Formula	29
III.	Structural Formula	29
IV.	Molecular Weight	29
V.	Specifications	29
VI.	Description	
VII.	Analytical Methods	32
VIII.	Occurrence	34
<u>BIOLOGICAL DATA</u>		
I.	Acute Toxicity	39
	Aquatic Animals	39
	Mice	39
	Rats	39
	Rabbits	42
II.	Short-Term Toxicity	44
	Rats	44
	Humans	45
III.	Long-Term Toxicity	56
IV.	Special Studies	57

## Table of Contents (Cont'd)

	Page
Frogs	57
Mice	58
Guinea Pigs	64
Rabbits	66
Dogs	68
Humans	69
<u>BIOCHEMICAL ASPECTS</u>	
I. Breakdown	72
II. Absorption and Distribution	72
Rats	72
Rabbits	74
Dogs	75
Humans	79
III. Metabolism and Excretion	84
Rats	84
Dogs	87
Pigs	89
Humans	90
IV. Enzymes and Other Biochemical Parameters	96
<u>In Vitro</u>	96
Rats	100
Guinea Pigs	100
Dogs	100
V. Drug Interaction	104
Rats	104
Rabbits	106
Dogs	110
Humans	111
VI. Consumer Exposure	113

## SUMMARY

### Description and Specifications

1. Magnesium carbonate: Magnesium carbonate is a light, white, friable mass or a bulky, white powder. It occurs naturally as the minerals Landsfordite and Magnesite. The formula for magnesium carbonate is approximately  $(\text{MgCO}_3)_4 \cdot \text{Mg}(\text{OH})_2 \cdot 5\text{H}_2\text{O}$ . It is an odorless substance which is practically insoluble in water (1 part in 3300 parts of  $\text{CO}_2$ -free water), is insoluble in alcohol, and is dissolved by dilute acids with effervescence. An assay of food grade magnesium carbonate should yield the equivalent of not less than 40.0% and not more than 43.5% of  $\text{MgO}$ .
2. Magnesium chloride: Magnesium chloride, anhydrous and hexahydrate, occurs as colorless, odorless, deliquescent flakes or crystals which are very soluble in water and freely soluble in alcohol. The molecular formulae are  $\text{MgCl}_2$  (MW=95.23) and  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$  (MW = 203.30) for the anhydrous and hexahydrate forms, respectively. The anhydrous form melts at  $712^\circ$ , with rapid heating. Food grade magnesium chloride should be not less than 99% and not more than the equivalent of 105% of  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ .
3. Magnesium hydroxide: Magnesium hydroxide with the molecular formula  $\text{Mg}(\text{OH})_2$  and molecular weight 58.34, is a white, bulky powder which occurs naturally as the

mineral Brucite. It is soluble in dilute acids and is practically insoluble in water. An assay of food grade magnesium hydroxide should yield not less than 95%  $\text{Mg}(\text{OH})_2$  after drying. Loss on ignition should be between 30% and 33%.

4. Magnesium oxide: Magnesium oxide, with the molecular formula  $\text{MgO}$  and molecular weight 40.32, occurs as a very bulky, white powder (light), 5 grams of which occupy 40-50 ml, or as a relatively dense, white powder (heavy), 5 grams of which occupy 10-20 ml. Periclase is a naturally occurring mineral of  $\text{MgO}$ . The melting point of magnesium oxide is  $2800^\circ$ . It is practically insoluble in water and is insoluble in alcohols. An assay of food grade magnesium oxide should yield not less than 96%  $\text{MgO}$  after ignition, with not less than 1.5%  $\text{CaO}$ . Loss on ignition should be not more than 10%.

5. Magnesium phosphate, dibasic: Dibasic magnesium phosphate is a white odorless crystalline powder with the molecular formula  $\text{MgHPO}_4 \cdot 3\text{H}_2\text{O}$  and molecular weight 174.36. It occurs naturally as the minerals Newbergite and Phosphor-roesslerite. It has a density of 2.13 and is slightly soluble in water, soluble in dilute acids and is insoluble in alcohol. An assay of food grade dibasic magnesium phosphate should yield not less than 96% of  $\text{Mg}_2\text{P}_2\text{O}_7$  calculated on the ignited basis. Loss on ignition should be between 29% and 36%.

6. Magnesium phosphate, tribasic: Tribasic magnesium phosphate, which may contain 4, 5 or 8 molecules of water of hydration, is a white, odorless, tasteless crystalline powder. The pentahydrate occurs most frequently, and the molecular formula for the anhydrous form is  $\text{Mg}_3(\text{PO}_4)_2$  with molecular weight, 262.86. It is insoluble in water but readily soluble in dilute mineral acids. An assay of food grade tribasic magnesium phosphate should yield not less than 98% and not more than the equivalent of 101.5% of  $\text{Mg}_3(\text{PO}_4)_2$ , calculated on the ignited basis. Loss on heating should be between 15% and 23% for the tetrahydrate; 20% and 27% for the pentahydrate; and 30% and 37% for the octahydrate.

7. Magnesium stearate: Magnesium stearate with molecular formula  $\text{Mg}(\text{C}_{18}\text{H}_{35}\text{O}_2)_2$  contains 96% stearic acid with some palmitic acid in the article of commerce. It occurs as a fine, white, bulky powder, having a faint characteristic odor. It is insoluble in water, alcohol and ether and is decomposed by dilute acids. An assay of food grade magnesium stearate should yield not less than 6.8% and not more than 8% of the equivalent of 8% of  $\text{MgO}$ .

8. Magnesium sulfate: Magnesium sulfate,  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , has a molecular weight of 246.47. It occurs as small, colorless crystals, usually needle-like, with a cooling, saline, bitter taste. It occurs naturally as the mineral



Kieserite. It is freely soluble in water giving neutral solutions, is slowly soluble in glycerin, and is sparingly soluble in alcohol. An assay of food grade magnesium sulfate should yield not less than 99.5%  $\text{MgSO}_4$  after ignition.

#### Acute Toxicity

Dowden and Bennet (198) found the median tolerance limit in mg/liter ( $\text{TL}_m$ ) for water fleas (Daphnia magna) to be: 3,391 at 25 hours; 3,699 at 50 hours; and 3,484 at 100 hours. All in standard reference water support medium.

The same authors found the  $\text{TL}_m$  for 3 species of aquatic animals to be: (a) water fleas (Daphnia magna); 963 at 24 hours, 929 at 48 hours, 861 at 72 hours, and 788 at 96 hours, all in glass wool filtered University Lake water support medium. In standard reference water support medium the  $\text{TL}_m$  was 3,803 at 96 hours. (b) Fish (Lepomis macrochirus); 19,000 at 24 hours; and (c) snail eggs (Lymnaea sp.); 10,530 at 24 hours, 6,525 at 48 hours, 6,300 at 72 hours and 6,250 at 96 hours in glass wool filtered University Lake water support medium.

Barbour and Winter (037) found the  $\text{LD}_{50}$  for the subcutaneous injection of magnesium chloride in white mice to be 1050 mg/kg BW.

Selisko and Ackerman (681) determined that the  $\text{LD}_{100}$  for intraperitoneal administration to Wistar albino rats was 100 mg Mg.

The same authors (689) found that the LD<sub>50</sub>'s for intraperitoneal administration to Wistar albino rats at various concentrations of magnesium chloride were: 100 mg Mg at 6.5 mg/ml; 90 mg Mg at 13.0 mg/ml; and 66 mg Mg at 17.8 mg/ml.

Selisko and Ackerman (682) found that the LD<sub>50</sub>'s for intraperitoneal administration to Wistar albino rats at various concentrations of magnesium sulfate were: 150 mg Mg at 6.5 mg/ml; 130 mg Mg at 13.0 mg/ml; and 140 mg Mg at 17.8 mg/ml.

Barbour and Taylor (639) found the LD<sub>50</sub> for subcutaneous administration of magnesium chloride to rabbits to be 725 mg/kg BW.

#### Short-Term

#### Toxicity

Moinuddin and Lee (515) fed magnesium sulfate in amounts from 0.88 mmole/kg feed to 138 mmole/kg feed to weanling rats for 4 weeks. They observed chronic diarrhea and depressed intake of food. At the highest amount (138 mmole/kg feed), there was a large increase in cecal weight and length. The authors pointed out that this change had also been reported when MgCl<sub>2</sub> or MgCO<sub>3</sub> were fed in large quantity to rats.

Many magnesium sulfate poisonings were reported in the literature over the past 40 years. Some of the significant aspects of nine such cases are: (1) Thatcher (766) reported the case of a 26-year-old male who died

an hour after accidentally ingesting a large amount of magnesium sulfate solution. The quantity and concentration were not given. The author cautioned that considering other reports of death and toxicity from magnesium sulfate administration, there are individuals for whom even an average dose may be toxic. (2) As reported by Roller (636), an injection of 2 ml of 20% magnesium sulfate caused tetany with respiratory distress in a 21-year-old woman. (3) Byron (113) discussed the cases of 5 young children who died after a strong magnesium sulfate purge was administered. The authors noted that particularly with strong solutions, greater magnesium absorption than ordinarily occurs can take place. (4) Collins and Russell (146) reported a fatality in a 4-year-old resulting from magnesium sulfate absorption. They pointed out that the gastrointestinal membrane is quite permeable to magnesium salts. (5) Stevens and Wolff (735) reported two cases of toxicity, one fatal, resulting from rectal administration of magnesium sulfate. The authors observed that normal kidneys may not remove magnesium sulfate from the blood stream fast enough to prevent amounts toxic to the central nervous system from accumulating. (6) Rosler (641) warned against the peroral administration of large doses of magnesium sulfate (epsom salts) after the deaths of two persons, undergoing a tapeworm cure. He pointed out

that under certain physiological conditions magnesium sulfate is absorbed from the intestinal tract which inundates the system with a large amount of magnesium leading to deep narcosis and paralysis of the respiratory center. (7) Colomb et al. (147) discussed the case of lesions on the cheeks and lips resulting from epsom salts spilled on the face. (8) Rees (620) reported the death of a pregnant young woman who dies from the effects of epsom salts used to induce abortion, which had entered the blood stream via the uterus. (9) Willner (827) described the death of a hospitalized woman accidentally administered magnesium sulfate by duodenal tube.

Allison (012) discussed a toxic sulfate effect from water high in  $\text{MgSO}_4$  and other sulfates, on both man and livestock. He urged further experimental research particularly concerning the effect on calcium equilibrium.

Hirschfelder (323) studied the effects of oral administration of magnesium sulfate on normal persons and those with renal disease. He concluded that an ordinary epsom salts purgation of 20-30 grams in nephritic patients can induce a magnesium coma easily mistaken for uremic coma. He found that the maximum increase of plasma magnesium was 0.4 mg/100 cc for normal individuals whereas it was 7-9 mg/100 cc in nephritic patients.

Abramowitz and Russo (001) described the case of a patient who developed a rash on her hands after ingesting magnesium hydroxide either as tablets or in suspension.

Brady and Williams (098), Lipsitz and English (448) and Lipsitz (447) all reported magnesium toxicity in newborns whose mothers were administered magnesium sulfate prior to delivery. Stone and Pritchard (736) on the other hand, reported that they had not found any deleterious effects from this treatment.

Long-Term  
Toxicity

No chronic or long-term toxicity studies could be found in the literature.

Special Studies

Labkovsky (424) found that magnesium salts (chloride or sulfate) administered orally or subcutaneously stimulated the induction of pulmonary adenomas by urethan.

Bazikyan and Akimov (051) studied the antineoplastic effect of magnesium chloride. They found that under the conditions of their experiments and in the amount administered,  $MgCl_2$ : (1) reduced the toxicity of 9,10-dimethyl-1,2-benzanthracene (DMBA) and 1,2,5,6-dibenzanthracene (DBA); and (2) markedly decreased the number of skin papillomas induced by DMBA and subcutaneous sarcomas induced by DBA.

Wies (826) found that subcutaneous doses of magnesium sulfate solution lowered the fertility of the females and produced defects in the brain and spinal cord of some of the fetuses.

Neal and Neal (537) studied the effect of hard water and water containing magnesium sulfate on atherosclerosis.

They observed that the rabbits given magnesium sulfate were the only group which had no atherosclerosis, did not have diarrhea and were larger with more subcutaneous fat. In addition even though the blood serum of all the experimental groups contained the same amount of fat, only the serums from the magnesium sulfate group were clear, the others were turbid. The authors questioned whether there is a protective substance in hard water or whether there is a specific effect of magnesium on fat metabolism. Further research with other ionizing compounds needs to be done.

Ozsoylu (565) examined the effect of giving magnesium sulfate to control blood sugar levels in 3 diabetic children. It was found that the blood sugar levels of the diabetics dropped (116 mg%, 195 mg% and 234 mg%) within one hour following the infusion of magnesium sulfate. There was no effect on the blood sugar of normal controls.

Frewin et al. (237) found that when intraarterial infusions of magnesium sulfate were given into the brachial artery, there was a direct vasodilator action on the blood vessels of the forearm, as measured by forearm or hand blood flow.

Absorption and  
Distribution

Barbour and Winter (038) investigated both the absorability of soluble organic and inorganic magnesium salts

and the subject of magnesium and calcium storage. They performed a number of feeding experiments on dogs from which they concluded: (1) the absorptive capacity of magnesium cannot be regarded as a constant; (2) the form in which magnesium is ingested can determine both its rate and total amount of absorption; (3) both soluble organic and inorganic magnesium salts markedly raise the level of serum magnesium; (4) the retention of calcium, when there is prolonged administration of large amounts of soluble organic magnesium salts, is dependent on the phosphate intake. Adequate phosphate intake leads to no loss of serum calcium while a phosphate intake exceeding 4 mmoles/kg/day leads to retention of additional calcium.

In a review of magnesium metabolism Wacker and Parisi (809) discussed current knowledge of magnesium absorption. The dietary content of anions, particularly phosphate influences the amount of magnesium absorbed. Thus stool content varies both with intake and the anions in the diet. Calcium, by competition for a common absorptive pathway, also effects the amount of absorbed magnesium (Alcock and MacIntyre,(007)).

From animal studies it has been found that magnesium is mainly absorbed in the small intestine. Studies in man with  $^{28}\text{Mg}$  have shown that peak levels of the isotope are measured in the plasma between two and eight hours

after administration of the dose (Graham et al., 271 ). The time course of appearance of orally administered labeled magnesium ( $^{28}\text{Mg}$ ) in human plasma is consistent with the finding in animals that the small intestine is the main site of magnesium absorption. However, the development of hypermagnesemia after rectal enemas is evidence for colonic absorption as well (Stevens and Wolff, 735 ).

Normally about one-third of the ingested magnesium is absorbed and excreted in the urine. Under ordinary circumstances of dietary intake this figure is a good estimation. However, the total dietary intake of this element must be taken into consideration.

In a study (Graham et al., 271 ) of 13 subjects given oral  $^{28}\text{Mg}$ , 44.3% of a diet containing 20 mEq/day was found to be absorbed, whereas 75.8% of a diet containing 1.9 mEq and 23.7% of a diet containing 47 mEq/day were found to be absorbed.

The contention that little endogenous magnesium is eliminated by the fecal route under ordinary circumstances is supported by the observation that when  $^{28}\text{Mg}$  is given intravenously to normal subjects, only 1-2% is recovered in the feces (Aikawa et al., 005 and Silver et al., 701 ).



Metabolism and  
Excretion

Clark (142) studied the effects of orally administered magnesium chloride on calcium and phosphate metabolism in adult rats. The authors concluded that: (1) dietary magnesium ions markedly influenced the absorption and excretion of calcium and phosphorus; (2) renal conservation of phosphorus is responsible for the improvement in phosphorus balance resulting from supplemental dietary magnesium; (3) two possible consequences for human beings might be: (a) in the treatment or prevention of those metabolic bone diseases resulting from inadequate intestinal absorption of calcium and (b) the observation that supplementary dietary magnesium which resulted in a drop in urinary phosphorus could serve as a partial explanation for the reduction of renal calculus formation following oral administration of magnesium salts.

Massry et al. (486) evaluated the characteristics of the renal handling of magnesium in dogs. They found that magnesium excretion is determined by filtration and reabsorption alone without evidence for tubular secretion.

Smith et al. (712) studied the excretion of magnesium ions in dogs following the intravenous injection of magnesium sulfate. They concluded that during the first 3-4 hours the magnesium ions were distributed throughout the extracellular fluid, while subsequently indications were that some of the ions were segregated from the extracellular fluid and thus not excreted.

Aikawa et al. (005) used  $^{28}\text{Mg}$  to investigate the kinetics of magnesium distribution in normal and diseased subjects. Following intravenous administration the isotope rapidly appeared in the urine but only insignificant amounts appeared in the stool. This was taken to indicate that most of the magnesium in the stool is of exogenous origin. Equilibration of the isotope with magnesium in the body was slow and parenterally administered magnesium was excreted very slowly by the kidneys. There was no marked difference between the results obtained from healthy and diseased patients.

Consolazio et al. (151) studied magnesium excretion in sweat. They concluded that under extreme conditions sweat could account for about 25% of the daily magnesium loss.

Figure 1 in Biochemical Section III, D 4 shows those enzyme systems in intermediary metabolism known to require magnesium (Ozsoylu, 565 ).

Drake (200) attempted to ascertain whether magnesium oxide exerts a "protective effect" in calcium oxalate stone formation. The test group (daily oral administration of  $\text{MgO}$ ) was found to excrete significantly more calcium than the controls.

Enzymes and Other  
Biochemical  
Parameters

Both MacIntyre (463) and Wacker and Parisi (809) summarized the in vitro actions of magnesium ions (see main body of text for details). MacIntyre noted that

the reactions involving ATP and ADP which are activated by magnesium are so widespread that magnesium must influence all life processes. Wacker and Parisi (809) concluded that since magnesium is the most abundant divalent intracellular cation, that it would be a tenable hypothesis to assume analogous in vivo function.

LaBarre and Vesselovsky (421) investigated the influence of magnesium sulfate on the functioning of the exocrine portion of the pancreas. They concluded that: (1) magnesium sulfate has a depressive effect on gastric and pancreatic secretions; (2) this effect is dependent on the integrity of the encephalic centers. Thus in those persons whose pancreatic function is curtailed either by secretin administration or the presence of food in the digestive tract, magnesium sulfate causes a reduction in production of pancreatic juice and enzyme formation.

#### Drug Interaction

Barbour and Taylor (039) studied the relations obtained various combinations of sodium barbital and magnesium chloride with respect to toxicity and hypnotic action. They concluded that the combination of magnesium chloride and sodium barbital hastened the onset of narcosis and lessened the persistence of the narcotic effect without significantly increasing toxicity.

Barbour and Winter (037) and Winter and Barbour (831) investigated the effect of magnesium chloride on the antipyretic action of amidopyrine and sodium salicylate.

Their observations with respect to both these experiments were: (1) various combinations of magnesium chloride with amidopyrine showed both higher antipyretic action and lower toxicity than amidopyrine alone; (2) the combination of 150 mg/kg BW magnesium chloride with 50 mg/kg BW sodium salicylate was more effective than 100 mg/kg BW sodium salicylate alone. The authors concluded that magnesium salts potentiate the early effects of antipyretic drugs even when intense or prolonged action tends to be inhibited.

Winter et al. (832) studied the effect of magnesium oxide on the antipyretic action of phenacetin. They found that the antipyretic action of orally administered phenacetin was enhanced by the addition of MgO. Since MgO alone was not antipyretic they considered their results to be strongly indicative of a true synergism.

Gershoff and Prien (253) found that after 5 years, 30 out of 36 patients with recurring calcium oxalate urolithiasis, maintained on a daily oral program of magnesium oxide and vitamin B<sub>6</sub>, showed either no recurrence or decreased recurrence of stone formation. The authors postulated that a possible explanation for the observed prevention of renal stone formation might involve an effect on the solvent characteristics of urine.

### Consumer Exposure

Magnesium is a natural constituent of all vegetables, fruits, meats and seafoods. (See Tables in main body of text.) The magnesium compound most widely used for both food processing and pharmaceutical purposes, is magnesium carbonate (NAS/NRC Tables). In food processing it serves as: an anti-caking and drying agent, a carrier, a disintegrating and dispersing agent, a color fixative, a color retention agent and as an adjunct. Pharmaceutically, magnesium carbonate is used as an antacid and a laxative. Additional uses for other magnesium salts are: magnesium chloride (831) as an emulsifying agent for suspending dried skim milk in water; magnesium oxide (610) for clarifying sugar juice and in molasses and raw sugar processing; magnesium phosphates as nutrients and dietary supplements; and magnesium stearate (405) as a foaming and whipping agent and in plasticizers used for food packaging materials.

Hard water is another source of exposure to magnesium.

Magnesium has been recognized as an essential nutrient since 1932. However, as a major inorganic constituent of living things and as a major intracellular cation, it is not in the strictest sense a nutrient (677).

According to Schroeder et al. (677): (1) except in states of malabsorption and excessive excretion due to other conditions, there is no evidence of widespread dietary magnesium deficiency; (2) the daily dietary

requirement for adult man is probably the obligatory urinary excretion plus losses in sweat and the amount unabsorbed in feces; (3) there is no good evidence that marginal dietary intakes contribute to any chronic disease of unknown etiology.

## CHEMICAL INFORMATION

### MAGNESIUM CARBONATE

#### I. Nomenclature

A. Common names: Landsfordite, Magnesite (Magnesium carbonate minerals)

B. Chemical names: Magnesium carbonate, Magnesium carbonate hydroxide

C. Trade names: none

D. Chemical Abstracts Services Unique Registry Number: 1319502

#### II. Empirical Formula

Approximately  $O_{14}Mg_5C_4H_2 \cdot 5OH_2$

#### III. Structural Formula

Approximately  $(MgCO_3)_4 \cdot Mg(OH)_2 \cdot 5H_2O$

IV. Molecular Weight: 485.69 (based on approximated formula)

#### V. Specifications

The Food Chemicals Codex, Second Edition (148), presents the following specifications for food grade magnesium carbonate:

##### 1. Description

Magnesium carbonate is a basic hydrated magnesium carbonate or a normal hydrated magnesium carbonate. It occurs as light, white, friable masses, or as a bulky, white powder. It is odorless, and is stable in air. It is practically insoluble in water, to which, however, it imparts a slightly alkaline reaction. It is insoluble in alcohol, but is dissolved by dilute acids with effervescence. When treated with diluted

hydrochloric acid T.S., it dissolves with effervescence and the resulting solution gives positive tests for Magnesium.

## 2. Specifications

**Assay.** The equivalent of not less than 40% and not more than 43.5% of MgO.

**Limits of Impurities**

**Acid-insoluble substances.** Not more than 500 parts per million (0.5%).

**Arsenic (as As).** Not more than 3 parts per million (0.0003%).

**Calcium oxide.** Not more than 0.6%.

**Heavy metals (as Pb).** Not more than 30 parts per million (0.003%).

**Lead.** Not more than 10 parts per million (0.001%).

**Soluble salts.** Not more than 1%.

## 3. Tests

**Assay.** Dissolve about 1 gram, accurately weighed, in 30 ml of 1 N sulfuric acid, add methyl orange T.S., and titrate the excess acid with 1 N sodium hydroxide. From the volume of 1 N sulfuric acid consumed, deduct the volume of 1 N sulfuric acid corresponding to the content of calcium oxide in the weight of the sample taken for the assay. The difference is the volume of 1 N sulfuric acid equivalent to the magnesium oxide present. Each ml of 1 N sulfuric acid is equivalent to 20.16 mg of MgO and to 28.04 mg of CaO.

**Acid-insoluble substances.** Mix 5 grams with 75 ml of water, add hydrochloric acid in small portions, with agitation, until no more of the sample dissolves, and boil for 5 minutes.



If an insoluble residue remains, filter, wash well with water until the last washing is free from chloride, ignite, cool, and weigh.

Arsenic. A solution of 1 gram in 10 ml of diluted hydrochloric acid T.S. meets the requirements of the Arsenic Test.

Calcium oxide. Dissolve about 1 gram, accurately weighed, in a mixture of 3 ml of sulfuric acid and 22 ml of water. Add 50 ml of alcohol, and allow the mixture to stand overnight. If crystals of magnesium sulfate separate, warm the mixture to about 50° to dissolve them. Filter through a Gooch crucible containing an asbestos mat that previously has been washed with diluted sulfuric acid T.S., water, and alcohol, and ignited and weighed. Wash the crystals on the mat several times with a mixture of 2 volumes of alcohol and 1 volume of diluted sulfuric acid T.S. Ignite the crucible and contents at a dull red heat, cool, and weigh. The weight of calcium sulfate so obtained, multiplied by 0.4119, gives the equivalent of calcium oxide in the sample taken for the test.

Heavy metals. Dissolve 667 mg in 10 ml of diluted hydrochloric acid T.S., and evaporate the solution to dryness on a steam bath. Toward the end of the evaporation stir frequently to disintegrate the residue so that finally a dry powder is obtained. Dissolve the residue in 20 ml of water, and evaporate to dryness in the same manner as before. Redissolve the residue in 25 ml of water, and filter if necessary. This solution meets the requirements of the Heavy Metals Test, using 20 mcg of lead ion (Pb) in the control (Solution A).

Lead. A solution of 1 gram in 10 ml of diluted hydrochloric acid T.S. meets the requirements of the Lead Limit Test, using 10 mcg of lead ion (Pb) in the control.

Soluble salts. Mix 2 grams with 100 ml of a mixture of equal volumes of *n*-propyl alcohol and water. Heat the mixture to the boiling point with constant stirring, cool to room temperature, add water to make 100 ml, and filter. Evaporate 50 ml of the filtrate on a steam bath to dryness, and dry at 105° for 1 hour. The weight of the residue does not exceed 10 mg.

Packaging and storage. Store in well-closed containers.

Functional use in foods. Alkali; drying agent; color-retention agent; anticaking agent; carrier.

## VI. Description

### A. General characteristics

Magnesium carbonate occurs as an odorless, light, white, friable mass, or white, bulky powder.

### B. Physical properties

Magnesium carbonate is soluble in 3300 parts of CO<sub>2</sub>-free water, to which it imparts an alkaline reaction, and is more soluble in water containing CO<sub>2</sub>. It is insoluble in alcohol and soluble in weak acids with effervescence.

### C. Stability in containers

Not available.

## MAGNESIUM CHLORIDE

### I. Nomenclature

A. Common names: none

B. Chemical names: Magnesium chloride, Magnesium chloride  
(hexahydrate)

C. Trade names: Magnogene

D. Chemical Abstracts Services Unique Registry Number:

7786303 (anhydrous)

7791186 (hexahydrate)

### II. Empirical Formula

$\text{Cl}_2\text{Mg}$  (anhydrous)

### III. Structural Formula

$\text{MgCl}_2$ ;  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$

IV. Molecular Weight: 95.23 (anhydrous), 203.30 (hexahydrate)

### V. Specifications

The Food Chemicals Codex, Second Edition (1948), presents the following specifications for food grade magnesium chloride:

#### 1. Description

Colorless, odorless flakes or crystals. It is very deliquescent. It is very soluble in water and freely soluble in alcohol. A 1 in 10 solution gives positive tests for Magnesium, and for Chloride.

#### 2. Specifications

Assay. Not less than 99% and not more than the equivalent of 105% of  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ .

### Limits of Impurities

**Ammonium.** Not more than 50 parts per million (0.005%).

**Arsenic (as As).** Not more than 3 parts per million (0.0003%).

**Heavy metals (as Pb).** Not more than 10 parts per million (0.001%).

**Sulfate.** Not more than 200 parts per million (0.02%).

### 3. Tests

**Assay.** Dissolve about 450 mg, accurately weighed, in 25 ml of water, add 5 ml of ammonia-ammonium chloride buffer T.S. and 0.1 ml of eriochrome black T.S., and titrate with 0.05 M disodium ethylenediaminetetraacetate until the solution is blue in color. Each ml of 0.05 M disodium ethylenediaminetetraacetate is equivalent to 10.16 mg of  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ .

**Ammonium.** Dissolve 1 gram in 90 ml of water, and slowly add 10 ml of a freshly boiled and cooled solution of sodium hydroxide (1 in 10). Allow to settle, then decant 20 ml of the supernatant liquid into a color comparison tube, dilute to 50 ml with water, and add 2 ml of Nessler's reagent. Any color does not exceed that produced by 10 mcg of ammonium ( $\text{NH}_4$ ) ion in 48 ml of water and 2 ml of the sodium hydroxide solution.

**Arsenic.** A solution of 1 gram in 35 ml of water meets the requirements of the Arsenic Test.

**Heavy metals.** A solution of 2 grams in 25 ml of water meets the requirements of the Heavy Metals Test, using 20 mcg of lead ion (Pb) in the control (Solution A).

**Sulfate.** Any turbidity produced by a 1-gram sample does not exceed that shown in a control containing 200 mcg of sulfate ( $\text{SO}_4$ ).

Packaging and storage. Store in tight containers.

Functional use in foods. Color-retention agent; firming agent.

## VI. Description

### A. General characteristics

Magnesium chloride and its hexahydrate occur as colorless, odorless flakes or crystals which are deliquescent.

### B. Physical properties

1. Anhydrous. The density of the anhydrous form is 2.41 (also reported as 2.325) and the melting point is  $712^{\circ}$  (rapid heating). It is soluble in water with the evolution of much heat and forming clear solutions.

2. Hexahydrate. The density of the hexahydrate form of magnesium chloride is 1.56. It loses  $2 \cdot \text{H}_2\text{O}$  at  $100^{\circ}\text{C}$ , and begins to lose some HCl at  $110^{\circ}\text{C}$ . One gram is soluble in 0.6 ml of water, 0.3 ml of boiling water, and 2 ml of alcohol.

### C. Stability in containers

Not available.

## MAGNESIUM HYDROXIDE


### I. Nomenclature

- A. Common names: Brucite (naturally occurring mineral)
- B. Chemical names: Magnesium hydroxide, Magnesium hydrate
- C. Trade names: Marinco-H, Hydro-Magma (30% suspension of magnesium hydroxide in water)
- D. Chemical Abstracts Services Unique Registry Number: 1309428

### II. Empirical Formula



### III. Structural Formula



### IV. Molecular Weight: 58.34

### V. Specifications

The Food Chemicals Codex, Second Edition (148), presents the following specifications for food grade magnesium hydroxide:

#### 1. Description

A white, bulky powder. It dissolves in dilute acids, but is practically insoluble in water and in alcohol. A 1 in 20 solution in diluted hydrochloric acid T.S. gives positive tests for magnesium.

#### 2. Specifications

Assay. Not less than 95% of  $\text{Mg}(\text{OH})_2$  after drying.

Loss on ignition. Between 30% and 33%.

Limits of Impurities

Alkalies (free) and soluble salts. Passes test.

Arsenic (as As). Not more than 3 parts per million (0.0003%).

Calcium oxide. Not more than 1%.

Heavy metals (as Pb). Not more than 40 parts per million (0.004%).

Lead. Not more than 10 parts per million (0.001%).

Loss on drying. Not more than 2%.

### 3. Tests

Assay. Transfer about 400 mg, previously dried at 105° for 2 hours and accurately weighed, into an Erlenmeyer flask. Add 25 ml of 1 N sulfuric acid, and, after solution is complete, add methyl red T.S. and titrate the excess acid with 1 N sodium hydroxide. From the volume of 1 N sulfuric acid consumed, deduct the volume of 1 N sulfuric acid corresponding to the content of calcium oxide in the sample taken for the assay. The difference is the volume of 1 N sulfuric acid equivalent to the  $\text{Mg}(\text{OH})_2$  in the sample of magnesium hydroxide taken. Each ml of 1 N sulfuric acid is equivalent to 29.16 mg of  $\text{Mg}(\text{OH})_2$  and to 28.04 mg of  $\text{CaO}$ .

Loss on ignition. Transfer about 500 mg, accurately weighed, to a tared platinum crucible, and ignite, increasing the heat gradually, to constant weight.

Alkalies (free) and soluble salts. Boil 2 grams with 100 ml of water for 5 minutes in a covered beaker, then filter while hot. Titrate 50 ml of the cooled filtrate with 0.1 N sulfuric acid, using methyl red T.S. as the indicator. Not more than 2 ml of the acid is consumed. Evaporate 25 ml of the filtrate to dryness, and dry at 105° for 3 hours. Not more than 10 mg of residue remains.

Arsenic. A solution of 1 gram in 25 ml of diluted hydrochloric acid T.S. meets the requirements of the Arsenic Test.

Calcium oxide. Dissolve about 500 mg, accurately weighed, in a mixture of 3 ml of sulfuric acid and 22 ml of water. Add 50 ml of alcohol, and allow the mixture to stand overnight. Warm the mixture to about 50°, if necessary, to dissolve any crystals of magnesium sulfate, and filter through a Gooch crucible containing an asbestos mat which has been previously washed with diluted sulfuric acid T.S., water, and alcohol, and ignited. Wash the crystals on the mat several times with a mixture of 3 volumes of alcohol and 1 volume of water. Ignite the crucible and contents at a dull red heat, cool, and weigh. The weight of calcium sulfate thus obtained, multiplied by 0.4119, gives the equivalent of calcium oxide (CaO).

Heavy metals. Dissolve 1 gram in 10 ml of diluted hydrochloric acid T.S., and evaporate to dryness on a steam bath. Toward the end of the evaporation, stir the residue frequently, disintegrate it to obtain a dry powder, dissolve the powder in 20 ml of water, and filter. A 10-ml portion of the filtrate meets the requirements of the Heavy Metals Test, using 20 mcg of lead ion (Pb) in the control (Solution A).

Lead. A solution of 1 gram in 20 ml of diluted hydrochloric acid T.S., meets the requirements of the Lead Limit Test, using 10 mcg of lead ion (Pb) in the control.

Loss on drying. Dry at 105° for 2 hours.



Packaging and storage. Store in tight containers.

Functional use in foods. Alkali; drying agent; color-retention agent.

VI. Description

A. General characteristics

Magnesium hydroxide occurs as a white bulky amorphous powder.

B. Physical properties

Magnesium hydroxide is soluble in dilute acids, but is practically insoluble in water (1:80,000). The pH of an aqueous slurry is 9.5-10.5.

C. Stability in containers

Not available.

## MAGNESIUM OXIDE

### I. Nomenclature

- A. Common names: Magnesia, Calcined magnesia, Magnesia usta,  
Periclase (naturally occurring mineral)
- B. Chemical names: Magnesium oxide
- C. Trade names: Magcal, Maglite
- D. Chemical Abstracts Services Unique Registry Number: 1309484

### II. Empirical Formula

OMg

### III. Structural Formula

MgO

### IV. Molecular Weight: 40.32

### V. Specifications

The Food Chemicals Codex, Second Edition (148), presents the following specifications for food grade magnesium oxide:

#### 1. Description

A very bulky, white powder known as light magnesium oxide or a relatively dense, white powder known as heavy magnesium oxide. Five grams of light magnesium oxide occupy a volume of approximately 40-50 ml, while 5 grams of heavy magnesium oxide occupy a volume of approximately 10-20 ml. It is practically insoluble in water and is insoluble in alcohol. It is soluble in dilute acids. A solution of magnesium oxide in diluted hydrochloric acid T.S. gives positive tests for Magnesium.

## 2. Specifications

Assay. Not less than 96% of MgO after ignition.

### Limits of Impurities

Acid-insoluble substances. Not more than 0.1%.

Alkalies (free) and soluble salts. Passes test.

Arsenic (as As). Not more than 3 parts per million (0.0003%).

Calcium oxide. Not more than 1.5%.

Heavy metals (as Pb). Not more than 40 parts per million (0.004%).

Lead. Not more than 10 parts per million (0.001%).

Loss on ignition. Not more than 10%.

## 3. Tests

Assay. Ignite about 500 mg to constant weight at 800° to 825° in a tared platinum crucible, weigh the residue accurately, dissolve it in 30 ml of 1 N sulfuric acid, boil gently to remove any carbon dioxide, cool, add methyl orange T.S., and titrate the excess acid with 1 N sodium hydroxide. From the volume of 1 N sulfuric acid consumed deduct the volume of 1 N sulfuric acid corresponding to the content of calcium oxide in the magnesium oxide taken for the assay. The difference is the volume of 1 N sulfuric acid equivalent to the MgO in the portion of magnesium oxide taken. Each ml of 1 N sulfuric acid is equivalent to 20.15 mg of MgO and to 28.04 mg of CaO.

Acid-insoluble substances. Mix 2 grams with 75 ml of water, add hydrochloric acid in small portions, with agitation, until no more dissolves, and boil for 5 minutes. If an insoluble residue remains, filter, wash well with water until the last washing is free from chloride, ignite, cool, and weigh.

Alkalies (free) and soluble salts. Boil 2 grams with 100 ml of water for 5 minutes in a covered beaker, and filter while hot. Add methyl red T.S., and titrate 50 ml of the cooled filtrate with 0.1 N sulfuric acid. Not more than 2 ml of the acid is consumed. Evaporate 25 ml of the filtrate to dryness, and dry at 105° for 1 hour. Not more than 10 mg of residue remains.

Arsenic. A solution of 1 gram in 20 ml of diluted hydrochloric acid T.S. meets the requirements of the Arsenic Test.

Calcium oxide. Dissolve about 400 mg, accurately weighed, in a mixture of 3 ml of sulfuric acid and 22 ml of water. Add 50 ml of alcohol, and allow the mixture to stand overnight. If crystals of magnesium sulfate separate, warm the mixture to about 50° to dissolve them. Filter through a Gooch crucible containing an asbestos mat that previously has been washed with diluted sulfuric acid T.S., water, and alcohol, and ignited and weighed. Wash the crystals on the mat several times with a mixture of 2 volumes of alcohol and 1 volume of diluted sulfuric acid T.S. Ignite the crucible and contents at a dull red heat, cool, and weigh. The weight of calcium sulfate obtained, multiplied by 0.4119, gives the equivalent of calcium oxide in the sample taken for the test.

Heavy metals. Dissolve 500 mg in 20 ml of diluted hydrochloric acid T.S., and evaporate the solution to dryness on a steam bath. Toward the end of the evaporation stir frequently to disintegrate the residue so that finally a dry powder is obtained. Dissolve the residue in 20 ml of water and evaporate to dryness in the

same manner as before. Redissolve the residue in 20 ml of water and filter if necessary. This solution meets the requirements of the Heavy Metals Test, using 20 mcg of lead ion (Pb) in the control (Solution A).

Lead. A solution of 1 gram in 20 ml of diluted hydrochloric acid T.S. meets the requirements of the Lead Limit Test, using 10 mcg of lead ion (Pb) in the control.

Loss on ignition. Weigh accurately about 500 mg in a tared covered platinum crucible. Ignite at between 800° and 825° for 15 minutes, cool and weigh.

Packaging and storage. Store in tight containers.

Labeling. Label magnesium oxide to indicate whether it is light magnesium oxide or heavy magnesium oxide.

Functional use in foods. Alkali; neutralizer.

## VI. Description

### A. General characteristics

Magnesium oxide occurs as a very bulky, white powder (light form) or as a dense, white powder (heavy form). (Five grams of the light form occupies 40-50 ml; 5 grams of the heavy form occupies 10-20 ml.)

### B. Physical properties

Magnesium oxide has a melting point of 2800°, is practically insoluble in water and is insoluble in alcohols. It is soluble in dilute acids. The pH of a saturated aqueous solution is 10.3.

### C. Stability in containers


Not available.

## MAGNESIUM PHOSPHATE, DIBASIC


### I. Nomenclature

- A. Common names: Newbergite, Phosphor-roesslerite (naturally occurring minerals)
- B. Chemical names: Magnesium phosphate, dibasic, Magnesium hydrogen phosphate, Secondary magnesium phosphate
- C. Trade names: none
- D. Chemical Abstracts Services Unique Registry Number: 7782754

### II. Empirical Formula



### III. Structural Formula



### IV. Molecular Weight: 174.36

### V. Specifications

The Food Chemicals Codex, Second Edition (148), presents the following specifications for food grade dibasic magnesium phosphate:

#### 1. Description

A white, odorless crystalline powder. It is slightly soluble in water and insoluble in alcohol, but is soluble in dilute acids.

#### 2. Identification

- A. Dissolve about 200 mg in 10 ml of diluted nitric acid T.S. and add, dropwise, ammonium molybdate T.S. A greenish yellow precipitate of ammonium phosphomolybdate forms which is soluble in ammonia T.S.
- B. Dissolve 100 mg in 0.5 ml of diluted acetic acid T.S. and 20 ml of water. Add 1 ml of ferric chloride T.S., let stand for

5 minutes, and filter. The filtrate gives a positive test for Magnesium.

### 3. Specifications

Assay. Not less than 96% of  $Mg_2P_2O_7$  calculated on the ignited basis.

Loss on ignition. Between 29 and 36%.

#### Limits of Impurities

Arsenic (as As). Not more than 3 parts per million (0.0003%).

Fluoride. Not more than 10 parts per million (0.001%).

Heavy metals (as Pb). Not more than 30 parts per million (0.003%).

Lead. Not more than 5 parts per million (0.0005%).

### 4. Tests

Assay. Weigh accurately about 500 mg of the residue obtained in the test for Loss on ignition, dissolve it in a mixture of 50 ml of water and 1 ml of hydrochloric acid, dilute to 100 ml with water, and mix. Transfer 50 ml of this solution into a 250 ml Erlenmeyer flask, add 10 ml of ammonia-ammonium chloride buffer T.S. and 12 drops of eriochrome black T.S., and titrate with 0.1 M disodium ethylenediaminetetraacetate until the wine-red color changes to pure blue. Each ml of 0.1 M disodium ethylenediaminetetraacetate is equivalent to 22.25 mg of  $Mg_2P_2O_7$ .

Loss on ignition. Weigh accurately about 1 gram, and ignite, preferably in a muffle furnace, at  $800^\circ \pm 25^\circ$  to constant weight.

Arsenic. A solution of 1 gram in 5 ml of diluted hydrochloric acid T.S. meets the requirements of the Arsenic Test.

Fluoride. Transfer 5 grams of the sample into a 200-ml distilling flask connected with a condenser and carrying a thermometer and a dropping funnel equipped with a stopcock. Dissolve the sample in 25 ml of dilute sulfuric acid (1 in 4), add 6 glass beads, and connect the apparatus for distillation, using a 600-ml beaker to collect the distillate. Add 40 ml of the dilute sulfuric acid to the flask through the dropping funnel, then fill the funnel with water, heat the solution to boiling, and continue heating until the temperature reaches 165°. Adjust the stopcock of the dropping funnel so that the temperature is maintained at 165°  $\pm$  5°, and continue the distillation until about 300 ml has been collected. Rinse the condenser and condenser arm with water, collecting the rinsings in the beaker. Add sodium hydroxide T.S. to the distillate to make it alkaline to litmus paper, and then add 5 ml in excess. Add 5 ml of 30% hydrogen peroxide and 6 glass beads to the beaker, boil until a volume of about 30 ml is reached, and cool. Transfer the condensed distillate, including the glass beads, into a 125-ml distilling flask connected with a condenser and carrying a thermometer and a capillary tube, both of which must extend into the liquid. Add 30 ml of perchloric acid, and continue as directed under the Fluoride Limit Test, Method I, beginning with "Connect a small dropping funnel or a steam generator to the capillary tube."

Heavy metals. Suspend 1.33 grams in 20 ml of water, and add hydrochloric acid, dropwise, until the sample just dissolves. Adjust the pH to between 3 and 4, filter, and dilute the filtrate



to 40 ml with water. For the control (Solution A), add 20 mcg of lead ion (Pb) to 10 ml of the filtrate, and dilute to 40 ml. For the sample (Solution B), dilute the remaining 30 ml of the filtrate to 40 ml. Add 10 ml of hydrogen sulfide T.S. to each solution, and allow to stand for 5 minutes. Solution B is no darker than Solution A.

Lead. Dissolve 1 gram in 20 ml of diluted hydrochloric acid T.S., evaporate the solution to a volume of about 10 ml on a steam bath, dilute to about 20 ml with water, and cool. This solution meets the requirements of the Lead Limit Test, using 5 mcg of lead ion (Pb) in the control.

Packaging and storage. Store in well-closed containers.

Functional use in foods. Nutrient; dietary supplement.

## VI. Description

### A. General characteristics

Dibasic magnesium phosphate occurs as a white, odorless crystalline powder.

### B. Physical properties

Dibasic magnesium phosphate has a density of 2.13; it is slightly soluble in water, soluble in dilute acids, and is insoluble in alcohol.

### C. Stability in containers


Not available.

## MAGNESIUM PHOSPHATE, TRIBASIC


### I. Nomenclature

- A. Common names: Bobierrite (naturally occurring mineral)
- B. Chemical names: Magnesium phosphate, tribasic, "Neutral" magnesium phosphate, Tertiary magnesium phosphate, Tri-magnesium phosphate
- C. Trade names: none
- D. Chemical Abstracts Services Unique Registry Number: 10233871

### II. Empirical Formula



### III. Structural Formula



- IV. Molecular Weight: 353.00 (pentahydrate), 262.86 (anhydrous)

### V. Specifications

The Food Chemicals Codex, Second Edition (148), presents the following specifications for food grade tribasic magnesium phosphate:

#### 1. Description

Tribasic magnesium phosphate may contain 4, 5, or 8 molecules of water of hydration. It occurs as a white, odorless, tasteless crystalline powder. It is readily soluble in dilute mineral acids but is almost insoluble in water.

#### 2. Identification

- A. Dissolve about 200 mg in 10 ml of diluted nitric acid T.S. and add, dropwise, ammonium molybdate T.S. A greenish-yellow precipitate of ammonium phosphomolybdate forms which is soluble in ammonia T.S.

B. Dissolve 100 mg in 0.7 ml of diluted acetic acid T.S. and 20 ml of water. Add 1 ml of ferric chloride T.S., let stand for 5 minutes, and filter. The filtrate gives a positive test for Magnesium.

### 3. Specifications

Assay. Not less than 98% and not more than the equivalent of 101.5% of  $\text{Mg}_3(\text{PO}_4)_2$ , calculated on the ignited basis.

Titration value. Passes test.

Loss on heating.  $\text{Mg}_3(\text{PO}_4)_2 \cdot 4\text{H}_2\text{O}$ , between 15 and 23%;

$\text{Mg}_3(\text{PO}_4)_2 \cdot 5\text{H}_2\text{O}$ , between 20 and 27%;  $\text{Mg}_3(\text{PO}_4)_2 \cdot 8\text{H}_2\text{O}$ , between 30 and 37%.

#### Limits of Impurities

Arsenic (as As). Not more than 3 parts per million (0.0003%).

Fluoride. Not more than 10 parts per million (0.001%).

Heavy metals (as Pb). Not more than 30 parts per million (0.003%).

Lead. Not more than 5 parts per million (0.0005%).

### 4. Tests

Assay. Weigh accurately about 200 mg of the sample, and dissolve it in a mixture of 25 ml of water and 10 ml of diluted nitric acid T.S. Filter, if necessary, wash any precipitate, then dissolve the precipitate by the addition of 1 ml of diluted nitric acid T.S. Adjust the temperature to about  $50^\circ$ , add 75 ml of ammonium molybdate T.S., and maintain the temperature at about  $50^\circ$  for 30 minutes, stirring occasionally. Allow to stand for 16 hours or overnight at room temperature. Wash the precipitate once or twice with water by decantation, using from 30-40 ml

each time, and pour these two washings through a filter.

Transfer the precipitate to the same filter, and wash with potassium nitrate solution (1 in 100) until the last washing is not acid to litmus paper. Transfer the precipitate and filter to the precipitation vessel, add 50 ml of 1 N sodium hydroxide, agitate until the precipitate is dissolved, add 3 drops of phenolphthalein T.S., and then titrate the excess alkali with 1 N sulfuric acid. Each ml of 1 N sodium hydroxide is equivalent to 5.714 mg of  $Mg_3(PO_4)_2$ .

Titration value. Ignite about 3 grams at about 425° to constant weight, and dissolve, by warming, 2 grams of the ignited salt in 50 ml of 1 N hydrochloric acid. Cool, add methyl orange T.S., and slowly titrate the excess of 1 N hydrochloric acid with 1 N sodium hydroxide to a yellow color, shaking the mixture vigorously during the titration. Not less than 29 ml and not more than 30.8 ml of 1 N hydrochloric acid is consumed.

Loss on heating. Weigh accurately about 1 gram and heat at about 425° to constant weight.

Arsenic. A solution of 1 gram in 10 ml of diluted hydrochloric acid T.S. meets the requirements of the Arsenic Test.

Fluoride. Determine as directed in the Fluoride Limit Test under Magnesium Phosphate, Dibasic.

Heavy metals. Suspend 1.33 grams in 20 ml of water, and add hydrochloric acid, dropwise, until the sample just dissolves. Adjust the pH to between 3 and 4, filter, and dilute the filtrate to 40 ml with water. For the control (Solution A), add 20 mcg

of lead ion (Pb) to 10 ml of the filtrate, and dilute to 40 ml. For the sample (Solution B), dilute the remaining 30 ml of the filtrate to 40 ml. Add 10 ml of hydrogen sulfide T.S. to each solution, and allow to stand for 5 minutes. Solution B is no darker than Solution A.

Lead. Dissolve 1 gram in 20 ml of diluted hydrochloric acid T.S., evaporate the solution to a volume of about 10 ml on a steam bath, dilute to about 20 ml with water, and cool. This solution meets the requirements of the Lead Limit Test, using 5 mcg of lead ion (Pb) in the control.

Packaging and storage. Store in well-closed containers.

Functional use in foods. Nutrient; dietary supplement.

## VI. Description

### A. General characteristics

Tribasic magnesium phosphate, containing 4, 5, or 8 molecules of water of hydration, occurs as a white, odorless, tasteless crystalline powder.

### B. Physical properties

Tribasic magnesium phosphate is insoluble in water, but readily soluble in dilute mineral acids. It becomes anhydrous at 400°.

### C. Stability in containers

Not available.

## MAGNESIUM STEARATE

### I. Nomenclature

- A. Common names: none
- B. Chemical names: Magnesium stearate
- C. Trade names: none
- D. Chemical Abstracts Services Unique Registry Number: 557040

### II. Empirical Formula

Approximately  $O_4C_{36}H_{70}Mg$

### III. Structural Formula

Approximately  $Mg(C_{18}H_{35}O_2)_2$ , containing 96% stearic acid and some palmitic acid in the article of commerce.

### IV. Molecular Weight: Approximately 590

### V. Specifications

The Food Chemicals Codex, Second Edition (148), presents the following specifications for food grade magnesium stearate:

#### 1. Description

Magnesium stearate is a compound of magnesium with variable proportions of stearic and palmitic acids. It occurs as a fine, white, bulky, powder, having a faint, characteristic odor. It is unctuous, and is free from grittiness. It is insoluble in water, in alcohol, and in ether. It conforms to the regulations of the federal Food and Drug Administration pertaining to specifications for salts of fatty acids and fatty acids derived from edible fats sources.

## 2. Identification

- A. Heat 1 gram with a mixture of 25 ml of water and 5 ml of hydrochloric acid. Fatty acids are liberated, floating as an oily layer on the surface of the liquid. The water layer gives positive tests for Magnesium.
- B. Mix 25 grams of the sample with 200 ml of hot water, then add 60 ml of diluted sulfuric acid T.S., and heat the mixture, with frequent stirring, until the fatty acids separate cleanly as a transparent layer. Wash the fatty acids with boiling water until free from sulfate, collect them in a small beaker, and warm on a steam bath until the water has separated and the fatty acids are clear. Allow the acids to cool, pour off the water layer, then melt the acids, filter into a dry beaker, and dry at 105° for 20 minutes. The solidification point of the fatty acids so obtained is not below 54°.

## 3. Specifications

Assay. Not less than the equivalent of 6.8% and not more than the equivalent of 8% of MgO.

### Limits of Impurities

Arsenic (as As). Not more than 3 parts per million (0.0003%).

Heavy metals (as Pb). Not more than 40 parts per million (0.004%).

Lead. Not more than 10 parts per million (0.001%).

Loss on drying. Not more than 4%.

## 4. Tests

Assay. Boil about 1 gram, accurately weighed, with 50 ml of 0.1 N sulfuric acid for 10 minutes, or until the fatty acid layer

is clear, adding water if necessary to maintain the original volume. Cool, filter, and wash the filter and flask thoroughly with water until the last washing is not acid to litmus. Add methyl orange T.S., and titrate the excess sulfuric acid with 0.1 N sodium hydroxide. Each ml of 0.1 N sulfuric acid is equivalent to 2.015 mg of MgO.

Arsenic. Mix 1 gram of the sample with 10 ml of hydrochloric acid and 8 drops of bromine T.S., and heat on a steam bath until a transparent layer of melted fatty acid forms. Add 50 ml of water, boil down to about 25 ml, and filter while hot. Cool, neutralize with a 1 in 2 solution of sodium hydroxide, and dilute to 35 ml with water. This solution meets the requirements of the Arsenic Test.

Heavy metals. Place 750 mg of the sample in a porcelain dish, place 250 mg of the sample in a second dish for the control, and to each add 5 ml of a 1 in 4 solution of magnesium nitrate in alcohol. Cover the dishes with 7.6 cm short stem funnels so that the stems are straight up. Heat for 30 minutes on a hot plate at the low setting, then heat for 30 minutes at the medium setting, and cool. Remove the funnels, add 20 mcg of lead ion (Pb) to the control, and heat each dish over an Argand burner until most of the carbon is burned off. Cool, add 10 ml of nitric acid, and transfer the solutions into 250-ml beakers. Add 5 ml of 70% perchloric acid, evaporate to dryness, then add 2 ml of hydrochloric acid to the residues, and wash down the inside of the beakers with water. Evaporate carefully to dryness



again, swirling near the dry point to avoid spattering. Repeat the hydrochloric acid treatment, then cool, and dissolve the residues in about 10 ml of water. To each solution add 1 drop of phenolphthalein T.S. and sufficient sodium hydroxide T.S. until the solutions just turn pink, and then add diluted hydrochloric acid T.S. until the solutions become colorless. Add 1 ml of diluted acetic acid T.S. and a small amount of charcoal to each solution, and filter through Whatman No. 2, or equivalent, filter paper into 50-ml Nessler tubes. Wash with water, dilute to 40 ml, and add 10 ml of hydrogen sulfide T.S. to each tube. The color in the solution of the sample does not exceed that produced in the control.

Lead. Ignite 500 mg in a silica crucible in a muffle furnace at 475°-500° for 15-20 minutes. Cool, add 3 drops of nitric acid, evaporate over a low flame to dryness, and re-ignite at 475°-500° for 30 minutes. Dissolve the residue in 1 ml of a mixture of equal parts by volume of nitric acid and water, and wash into a separator with several successive portions of water. Add 3 ml of Ammonium Citrate Solution and 0.5 ml of Hydroxylamine Hydrochloride Solution, and make alkaline to phenol red T.S. with stronger ammonia T.S. Add 10 ml of Potassium Cyanide Solution. Immediately extract the solution with successive 5-ml portions of Dithizone Extraction Solution, draining off each extract into another separator, until the last portion of dithizone solution retains its green color. Shake the combined extracts for 30 seconds with 20 ml of dilute nitric acid (1 in 100), and discard the

chloroform layer. Add to the acid solution exactly 4 ml of Ammonia-Cyanide Solution and 2 drops of Hydroxylamine Hydrochloride Solution. Add 10 ml of Standard Dithizone Solution, and shake the mixture for 30 seconds. Filter the chloroform layer through an acid-washed filter paper into a Nessler Tube, and compare the color with that of a standard prepared as follows: to 20 ml of dilute nitric acid (1 in 100), add 5 mcg of lead ion (Pb), 4 ml of Ammonia-Cyanide Solution and 2 drops of Hydroxylamine Hydrochloride Solution, and shake for 30 seconds with 10 ml of Standard Dithizone Solution. Filter through an acid-washed filter paper into a Nessler tube. The color of the sample solution does not exceed that in the control.

Loss on drying. Dry at 105° to constant weight, using 2-hour increments of heating.

Packaging and storage. Store in well-closed containers.

Functional use in foods. Anticaking agent; binder; emulsifier.

## VI. Description

### A. General characteristics

Commercial preparations of magnesium stearate is a magnesium compound containing various proportions of stearic and palmitic acids. It occurs as a fine, white, bulky powder, having a faint characteristic odor.

### B. Physical properties

Magnesium stearate is insoluble in water, alcohol and ether and is decomposed by dilute acids.

### C. Stability in containers

Not available.

## MAGNESIUM SULFATE

### I. Nomenclature


A. Common names: Bitter salts, Epsom salts, Kieserite (naturally occurring mineral)

B. Chemical names: Magnesium sulfate, Sulfuric acid, magnesium salt, heptahydrate

C. Trade names: none

D. Chemical Abstracts Services Unique Registry Number: 10034998

### II. Empirical Formula



### III. Structural Formula



IV. Molecular Weight: 246.47

### V. Specifications

The Food Chemicals Codex, Second Edition (148), presents the following specifications for food grade magnesium sulfate:

#### 1. Description

Small, colorless crystals, usually needle-like, with a cooling, saline, bitter taste. It is freely soluble in water, slowly soluble in glycerin, and sparingly soluble in alcohol. It effloresces in warm, dry air. Its solutions are neutral. A 1 in 20 solution gives positive tests for Magnesium, and for Sulfate.

#### 2. Specifications

Assay. Not less than 99.5% of  $\text{MgSO}_4$  after ignition.

Loss on ignition. Between 40 and 52%.

### Limits of Impurities

Arsenic (as As). Not more than 3 parts per million (0.0003%).

Heavy metals (as Pb). Not more than 10 parts per million (0.001%).

Selenium. Not more than 30 parts per million (0.003%).

### 3. Tests

Assay. Weigh accurately about 500 mg of the residue obtained in the test for Loss on ignition, dissolve it in a mixture of 50 ml of water and 1 ml of hydrochloric acid, dilute to 100 ml with water, and mix. Transfer 50 ml of this solution into a 250-ml Erlenmeyer flask, add 10 ml of ammonia-ammonium chloride buffer T.S. and 12 drops of eriochrome black T.S., and titrate with 0.1 M disodium ethylenediaminetetraacetate until the wine-red color changes to pure blue. Each ml of 0.1 M disodium ethylenediaminetetraacetate is equivalent to 12.04 mg of  $\text{MgSO}_4$ .

Loss on ignition. Weigh accurately about 1 gram in a crucible, heat at  $105^\circ$  for 2 hours, then ignite in a muffle furnace at  $450^\circ \pm 25^\circ$  to constant weight.

Arsenic. A solution of 1 gram in 10 ml of water meets the requirements of the Arsenic Test.

Heavy metals. A solution of 2 grams in 25 ml of water meets the requirements of the Heavy Metals Test, using 20 mcg of lead ion (Pb) in the control (Solution A).

Selenium. A solution of 2 grams in 40 ml of dilute hydrochloric acid (1 in 2) meets the requirements of the Selenium Limit Test.

Packaging and storage. Store in well-closed containers.

Functional use in foods. Nutrient; dietary supplement.

## VI. Description

### A. General characteristics

Magnesium sulfate, heptahydrate, occurs as small, colorless, usually needle-like crystals. It has a cooling, saline, bitter taste.

### B. Physical properties

The density of magnesium sulfate, heptahydrate is 1.67. It loses approximately one  $H_2O$  (7%) on exposure to air at ordinary temperature; loses four molecules of  $H_2O$  (about 28%) at 70-80°; loses five molecules of  $H_2O$  (36.5%) at 100°; loses 6 molecules of  $H_2O$  at 120° and becomes anhydrous at 250°. Rapidly reabsorbs water on exposure to moist air. One gram dissolves in 0.8 ml of water, 0.2 ml of boiling water, slowly soluble in 1.1 ml of glycerin and is slightly soluble in alcohol. Aqueous solutions are neutral.

### C. Stability in containers

Not available.

Note: Magnesium sulfate may also occur as the trihydrate, an odorless powder, which is used in the preparation of aperients (Morison's paste).

## VII. Analytical Methods

1. Atomic absorption spectroscopy has been widely used as a sensitive method devoid of most interference problems for the detection and estimation of various metals. Applications of this method to the determination of magnesium in biological material have been presented by Dawson and Heaton (171), Horn and Latner (334), and Sunderman and Carrol (743), who compare atomic absorption spectroscopy to the Bohuon method (see below).

The major probable source of interference in magnesium determination in biological material arises from formation of sulfate and phosphate salts which may under some flame conditions be resistant to thermal molecular dissociation. A study of the precision and accuracy of the atomic absorption method (811) showed a mean value of a sample known to contain 0.954 ppm to be  $0.956 \pm 0.035$  ppm with a coefficient of variation of 3.7% based on seventeen determinations. The accuracy was 100.3%.

Other workers have used atomic absorption spectroscopy for the determination of magnesium in biological materials (258,261 ,341 ,346 , and 393 ) and special note is made of the work of Schroeder, Nason, and Tipton (461) who used atomic absorption spectroscopy in the magnesium analysis of foods and water.

2. The Magnon reagent, the sodium salt of 1-azo-2-hydroxy-3-(2,4-dimethoxycarboxyanilino)naphthalene-1-(2-hydroxybenzene-4-sulfonic acid), of Mann and Yoe has been used for the spectrophotometric determination of magnesium in various biological mediums. The reagent is eight times more sensitive than the titan yellow reagent, also used in magnesium determinations. C. Bohuon (089) has presented a technique for the Magnon determination of magnesium which requires 100  $\mu$ l of plasma or cerebrospinal fluid, which gave an accuracy of near 100% (sic) based on recovery of known

amounts of magnesium. Rice and Lapara have modified the Bohuon method to require only 40 ul of sample with increased sensitivity, accuracy and precision.

A discussion of the optimum conditions for the Magnon spectrophotometric determination of magnesium is presented by Burcar, et al. (109).

Phosphate is the major interfering ion causing low results in determinations, but may be removed by precipitation with aluminum at pH 5.5-5.6, being careful to avoid adding large excesses of aluminum which may also interfere with the determination. Gluconate does not interfere as in determinations with titan yellow. Calcium interference which is somewhat inconsistent may be compensated for by adding a known quantity of calcium to a standard curve developed at the time of reaction. Interference due to protein may be compensated for by precipitation of protein with trichloroacetic acid or addition of protein to the standards.

3. Another spectrophotometric method for determining magnesium in biological fluids employs the dye Eriochrome Black T (EBT), 1-(1-hydroxy-2-naphthylazo)-6-nitro-2-naphthol-4-sulfonic acid, sodium salt. The reagent forms metallochromic complexes with several cations, particularly calcium, which may interfere. The response of the calcium to the reagent can be eliminated by incorporating barium-EGTA (the barium salt of ethylene glycol bis( $\beta$ -aminoethyl ether)-N,N,N',N'-tetraacetic acid). The calcium displaces the barium which does not interfere in the EBT reaction. Interferences due to iron, copper and zinc can be overcome by addition of trace amounts of cyanide ion. Connerty and Briggs (150) compared results from serum magnesium analyses by the EBT spectrophotometric method with results obtained from atomic absorption spectroscopy and found no evidence of a systematic difference between the methods.

4. Other methods for the determination of magnesium include flame emission spectroscopy (762, and 790); spectrochemical methods (517 and 568 ); fluorimetric methods (048, 392, and 555); the titan yellow colorimetric method (037, 156 , 219 , 353 , 477 , and 794).

With respect to ease of operation and accuracy and precision of results, none of these above methods offer any significant advantages.

#### VIII. Occurrence

##### A. Plants

Magnesium, which is essential for life, is present in all plants. The magnesium  $2+$  ion is a required co-factor for many enzymes, particularly phosphohydrolases and phosphotransferases. In green leaf plants, magnesium  $2+$  functions as a coordinate ion, stabilizing the porphyrin ring structure of chlorophylls a and b. While it has long been believed that green leafy vegetables are a major source of magnesium, Schroeder, et al. (461) showed that this has been found true only on a caloric basis and not on a weight basis. In their analyses (by atomic absorption spectrophotometry), they obtained mean concentrations of 170 ppm in 9 green, leafy vegetables; 194 ppm in 8 root vegetables; 241 ppm in 3 legumes; and 184 ppm in 11 vinous or fleshy vegetables. Schroeder et al. further found that analyses made on white and green parts of the same plant showed 55-61% more magnesium in the white portion on a weight basis. (For a complete breakdown of magnesium concentrations in various food plants, consult Table 22, in Biological Data, Section VI.)

##### B. Animals

Iain MacIntyre (463) has presented comparative magnesium concentrations for various animals. These values are shown in Table I. (Note:



Table I. Magnesium Content of Tissue and Serum of Various Animals (463)

Animal	Habitat	Tissue (mEq/kg)	Serum (mEq/l)
Man	T	19	1.7
Pig	T	22	
Dog	T	19	1.9
Frog	A	19	
Pike (teleost)	FW	26	
Cod (teleost)	M		5
Dogfish (elasmobranch)	M		12
Crab (limulus)	M		83
Lobster (Homarus)	M		13
Crab (Astacus)	FW		5
Octopod (Eledone)	M	25	114
Mussel (Anodonta)	FW	5	0.7

---

T = Terrestrial

A = Amphibious

M = Marine

FW = Fresh water

These values were collected from sources dated between 1896 and 1950, and therefore, because of less precise methods of analysis, may not be accurate.)

The adult human body contains between 21 and 28 grams (approximately 2000 mEq) of magnesium (809). About half of this amount is contained in bone, while the remainder is distributed evenly between muscle and non-muscular soft tissue. Reported values of serum magnesium have been in the range of 1.4-2.5 mEq/l. (This range may result partially from the imprecise methods of analysis.)

Tables 2 and 3 show values as reported by Schroeder, et al. (677) for magnesium in human tissue (analyzed by flame emission spectroscopy).

#### C. Synthetics

No information available.

#### D. Natural Inorganic Sources

Magnesium, which makes up about 2.1% of the earth's crust, occurs as a constituent of several minerals. These minerals consist of two types: The solubles; magnesite ( $\text{MgCO}_3$ ), dolomite ( $\text{MgCO}_3 \cdot \text{CaCO}_3$ ), and meerschaum, serpentine, talc, and asbestos which are all silicates and the insolubles; carnallite, kainite, schoenite and kieserite.

Magnesium salts are present in hard water, the degree of hardness of the water depending largely on the concentration of these salts plus calcium salts. The median magnesium concentrations of the water supplies of the 100 largest American cities is given as 6.25 ppm (677), with the breakdown in various waters shown in Table 4

Table 2. Magnesium In Human Tissues, U.S. Values,  
150 Accidental Deaths, Mean Values (677)

Organ	Organ weight (g)	Magnesium	
		Content (mg)	Concentration ( $\mu\text{g/g}$ wet wt $\pm$ S.E.M.)
Muscle	28,000	5300	189 $\pm$ 7.2
Fat	12,500	300	24 $\pm$ 0.2
Bone	10,000	11,000	1100 $\pm$ 25.2
Blood	5500	210	38 $\pm$ 0.4
Skin	4900	290	59 $\pm$ 4.0
Connective tissue	2000	300*	150*
Liver	1800	310	172 $\pm$ 5.2
Brain	1400	210	150 $\pm$ 3.4
G.I. tract	1200	150	125 $\pm$ 9.6
Lungs	1000	71	71 $\pm$ 2.2
Heart	350	63	180 $\pm$ 5.5
Kidneys	310	40	129 $\pm$ 3.3
Spleen	180	23	128 $\pm$ 4.2
Pancreas	100	16	160 $\pm$ 4.4
Other	760	114*	150*
Total	70000	18397	285
Soft tissues	60000	9000	150

\*Estimated from mean of all soft tissues. Analyses not done.

Note: Concentrations were obtained from analyses of 150 accidental deaths. Contents are based on organ weights of "standard man", using these concentrations. These 150 cases were selected from the total number analyzed on the basis of their dying accidentally and not of a known disease.

Table 3. Magnesium In Other Human Tissues, Mean Values (677)

	No. Samples	Ash (% of wet weight)	Magnesium	
			g/100 g ash ( $\pm$ S.E.M.)	$\mu\text{g/g}$ wet weight ( $\pm$ S.E.M.)
Adrenal	10	0.5	0.89 $\pm$ 0.10	45 $\pm$ 5.0
Aorta	94	1.7	1.5 $\pm$ 0.06	255 $\pm$ 10.2
Diaphragm	91	0.93	1.7 $\pm$ 0.05	167 $\pm$ 4.9
Esophagus	57	0.89	1.4 $\pm$ 0.06	125 $\pm$ 5.3
Stomach	110	0.82	1.5 $\pm$ 0.04	123 $\pm$ 3.3
Duodenum	60	0.81	1.3 $\pm$ 0.04	105 $\pm$ 3.2
Jejunum	83	0.92	1.3 $\pm$ 0.06	120 $\pm$ 5.5
Ileum	77	0.75	2.1 $\pm$ 0.12	158 $\pm$ 9.0
Cecum	31	0.67	3.1 $\pm$ 0.25	203 $\pm$ 16.8
Sigmoid colon	83	0.68	2.6 $\pm$ 0.22	177 $\pm$ 15.0
Rectum	41	0.85	2.6 $\pm$ 0.19	221 $\pm$ 16.1
Omentum	72	0.25	1.2 $\pm$ 0.06	30 $\pm$ 1.5
Larynx	49	3.3	1.2 $\pm$ 0.06	396 $\pm$ 19.8
Trachea	57	1.9	1.2 $\pm$ 0.05	228 $\pm$ 9.5
Thyroid	14	1.4	0.98 $\pm$ 0.19	137 $\pm$ 26.6
Prostate	40	1.2	1.6 $\pm$ 0.05	192 $\pm$ 7.2
Testis	63	1.1	1.1 $\pm$ 0.04	121 $\pm$ 4.4
Ovary	16	1.0	0.94 $\pm$ 0.06	94 $\pm$ 5.9
Uterus	30	1.0	1.1 $\pm$ 0.05	110 $\pm$ 6.0
Bladder	98	0.78	1.1 $\pm$ 0.05	109 $\pm$ 3.9

Table 4. Magnesium In Some Waters (677)

	ppm
Natural waters	
Sea water, Caribbean	1300
Lake George, N.Y.	2.0
Lake Spofford, N.H.	8.4
Connecticut, River, Brattleboro, Vt.	23.0
Springs	
Brattleboro, Vt.	2.4
West Brattleboro, Vt.	0.8
Silverdale, N.H.	14.1
Chesterfield, N.H.	3.6
Saratoga, N.Y. (State Seal water)	5.3
Well, Putney, Vt.	5.2
Municipal waters	
Bangor, Penn. (Roseto)	0.8
Brattleboro, Vt.	1.4
Bridgeport, Conn.	3.8
Gibsonville, Ohio	16.4
Peterborough, N.H.	1.3
White Pains, N.Y.	1.7

## BIOLOGICAL SECTION

### I. Acute Toxicity

#### A. Aquatic Animals

Dowden and Bennett (198) studied the median tolerance limits ( $TL_m$ ) of magnesium sulfate ( $MgSO_4$ ) and magnesium chloride ( $MgCl_2$ ) with 3 species of aquatic animals: water fleas, Daphnia magna; a fish, the bluegill, Lepomis macrochirus; and snail eggs, Lymnaea sp. in two different support media; standard reference water and University Lake water filtered through glass wool. The results are shown on the acute toxicity Table 5 . The values given for the snail eggs represent the concentration of the magnesium salt at which 50 percent of the eggs hatched.

The authors found that increasing or decreasing the pH of a concentration of a chemical can cause an increase or decrease in its toxicity.

#### B. Mice

Barbour and Winter (1937) studied the acute toxicity of magnesium chloride when administered subcutaneously to white mice (sex and age not given). The observations are shown in Table 6 .

#### C. Rats

1. Selisko and Ackermann (1961) showed that intraperitoneal injections of high doses of magnesium salts (9.75 mg) produced symptoms of tetany in albino rats (inbred,  $95 \pm 5$  g). The salts,  $MgSO_4$  and  $MgCl_2$  were administered in 3 concentrations; 1.5 ml with 6.5 mg Mg/ml, 0.75 ml with 13.0 mg Mg/ml, and 0.55 ml with 17.8 mg Mg/ml.

Many animals developed convulsions and all, including those that survived, developed narcosis. (See Table 7 for results.)

Table 5  
Acute Toxicity of  $\text{MgSO}_4$  and  $\text{MgCl}_2$  in Aquatic Animals (198)

Substance	Animal	Route	Dosage mg/kg body wt.	Measurement
Magnesium chloride	Water fleas <u>Daphnia magna</u>	In standard	3,391 at 25 hr.	$\text{TL}_m$ (mg/l)
		reference water	3,699 at 50 hr.	
		support medium	3,484 at 100 hr.	
Magnesium sulfate	Water fleas <u>Daphnia magna</u>	In glass wool	963 at 24 hr.	$\text{TL}_m$ (mg/l)
		filtered Univer-	929 at 48 hr.	
		sity Lake water	861 at 72 hr.	
		support medium	788 at 96 hr.	
	Water fleas <u>Daphnia magna</u>	In standard	3,803 at 96 hr.	$\text{TL}_m$ (mg/l)
		reference water		
	Fish <u>Lepomis</u> <u>macrochirus</u>	support medium		$\text{TL}_m$ (mg/l)
		In standard	19,000 at 24 hr.	
	Snail eggs <u>Lymnaea</u> sp.	reference water		$\text{TL}_m$ (mg/l)
		support medium		
		In glass wool	10,530 at 24 hr.	
		filtered Univer-	6,525 at 48 hr.	
		sity Lake water	6,300 at 72 hr.	
		support medium	6,250 at 96 hr.	

• Table 6. Acute Toxicity (037)

MgCl <sub>2</sub> (1% solution) mg/kg BW	No. of mice	
	Survived	Died
1000	2	0
1050	1	3
1100	0	4
1200	0	2
1300	0	2
1400	0	2
1500	0	2
2000	0	2

Table 7. Acute Toxicity (681)

Magnesium salt	Conc.	No. Animals	No. with convulsions	Death
Chloride	a	30	11 = 37%	1 = 3%
Chloride	b	30	30 = 100%	25 = 83%
Chloride	c	30	30 = 100%	30 = 100%
Sulfate	a	40	3 = 7%	0 = 0%
Sulfate	b	30	22 = 73%	3 = 10%
Sulfate	c	30	18 = 60%	2 = 7%

Mg dose, 9.75 mg each: (a) 1.5 ml with 6.5 mg Mg/ml, (b) 0.75 ml with 13.0 mg Mg/ml, (c) 0.55 ml with 17.8 mg Mg/ml.

The authors observed that the intensity of convulsions was greater with salts of monobasic than with salts of dibasic acids. Also as Table 7 shows the toxicity of the chloride increased with increased concentration while with the sulfate, the medium concentration was most toxic.

The authors concluded that when aqueous solutions of magnesium salts are administered intraperitoneally, the toxicity is largely dependent on both the anion and the concentration.

2. The same authors, Selisko and Ackermann (682) compared the LD<sub>50</sub>'s of several magnesium salts on Wistar albino rats (40 animals per salt, both sexes, 100  $\pm$  5 g in weight). The salts were given in a single intraperitoneal injection. The corresponding concentrations of the doses given (0.1 to 0.25 ml) had the same magnesium content. The LD<sub>50</sub>'s for magnesium sulfate and magnesium chloride at the three concentrations of Mg tested are shown on Table 8 .

#### D. Rabbits

Barbour and Taylor (039) subcutaneously injected normal, healthy adult rabbits with magnesium chloride. They found that 60% of the animals given a dose of 725 mg/g BW died. All of the ten animals given this dosage showed deep narcosis and six died, while of 5 rabbits given 750 mg/kg BW four showed deep narcosis and three died. Of eleven rabbits given 700 mg/kg BW, only five showed light narcosis and one deep narcosis with death. The figure of 725 mg/kg BW was therefore taken as the medium lethal dose for rabbits.



Table 8

Acute Toxicity of  $\text{MgSO}_4$  and  $\text{MgCl}_2$  in Mice, Rats and Rabbits

Substance	Animal	Sex & No.	Route	Dosage mg/kg body wt.	Measurement	Ref. Bibliogr. No.
Magnesium sulfate	Wistar albino rats	M & F 40	i.p.	150 mg Mg conc. of 6.5 mg/ml	LD <sub>50</sub>	Selisko and Ackerman (682)
	Wistar albino rats	M & F 40	i.p.	130 mg Mg at conc. of 13.0 mg/ml	LD <sub>50</sub>	Selisko and Ackerman (682)
	Wistar albino rats	M & F 40	i.p.	140 mg Mg at conc. of 17.8 mg/ml	LD <sub>50</sub>	Selisko and Ackerman (682)
Magnesium chloride	White mice	Sex not given, 4	s.c.	1050	LD <sub>50</sub>	Barbour and Winter (037)
Magnesium chloride	Albino rats	Sex not given, 30	i.p.	100 mg Mg	LD <sub>100</sub>	Selisko and Ackerman (681)
Magnesium chloride	Wistar albino rats	M & F 40	i.p.	100 mg Mg at conc. of 6.5 mg/ml	LD <sub>50</sub>	Selisko and Ackerman (682)
	Wistar albino rats	M & F 40	i.p.	90 mg Mg at conc. of 13.0 mg/ml	LD <sub>50</sub>	Selisko and Ackerman (682)
	Wistar albino rats	M & F 40	i.p.	66 mg Mg at conc. of 17.8 mg/ml	LD <sub>50</sub>	Selisko and Ackerman (682)
Magnesium chloride	Rabbits	Sex not given, 10	s.c.	725	LD <sub>50</sub>	Barbour and Taylor (039)

## II. Short-Term Toxicity

### A. Rats

Moinuddin and Lee (515) studied whether feeding hydrogogue cathartics such as magnesium sulfate, which exert considerable osmotic effects in the lumen, would increase gastrointestinal growth in the rat.

Three feeding experiments were carried out for a 4-week period, with weanling male Sprague-Dawley rats (6 rats per group). In each experiment varying amounts of additional magnesium sulfate (anhydrous powder) replaced the cornstarch in a basal diet which already contained a salt mixture (3.8% anhydrous magnesium sulfate plus 0.02% manganous sulfate monohydrate).

Experiment a: 0.88 mmole/kg feed of  $\text{MgSO}_4$  was added and fed at this level throughout the experiment. No unusual effects were observed at this low level of feeding.

Experiment b: The  $\text{MgSO}_4$  was added initially at 8.64 mmole/kg feed and increased as follows: doubled on 9th day to 17.28 mmole/kg; redoubled on the 17th day to 34.56 and again on the 25th day to 69.12 where it was maintained for the remaining days of the experiment.

Experiment c: The  $\text{MgSO}_4$  was added and maintained at 138 mmole/kg of feed throughout the experiment.

Controls: Basal diet with original salt mixture and cornstarch instead of added salt.

The observations of the dietary effect of  $\text{MgSO}_4$  were:

- (a) Chronic diarrhea.
- (b) Depressed intake of food, also found when  $\text{MgCl}_2$  is fed at excessive levels.

- (c) At the excessive dose fed in experiment c, there was a large increase in the weight of the cecum which with the rectum was also noticeably increased in length.
- (d) In experiment b, cecal contents were runny. In experiment c, both cecal and colon contents were runny.
- (e) Serum alkaline phosphatase activity was higher in  $\text{MgSO}_4$  fed rats than controls.

The authors note that the diarrhea and cecal distention observed in experiment c also has been reported to occur when  $\text{MgCl}_2$  or  $\text{MgCO}_3$  are fed to rats in large quantity.

#### B. Humans

1. A number of cases have been reported in the literature over the last forty years, describing magnesium sulfate poisoning. Several of these are summarized below chronologically.

- (a) Thatcher in 1928 (766) reported the case of a 26 year old hospitalized male who died one hour after accidentally ingesting a large amount of magnesium sulfate solution (quantity or concentration not stated). The most significant gross pathological changes were a large quantity (approximately 1 liter) of fluid in the stomach and the dark red, hemorrhagic appearance of the stomach lining. The most marked microscopic pathological changes were small hemorrhages on the mucosa of the stomach and intestine. The amount of magnesium sulfate recovered in the stomach (57 g) was more than three times the normal dosage (15.5 g) given. The lungs were markedly congested. The author suggested that considering other reports

of toxicity and death resulting from magnesium sulfate administration, there are individuals for whom even an average dose may be toxic.

- (b) Roller in 1936 (636) described a case in which a young woman (21 years) developed a picture of tetany with respiratory distress following injection of 2 ml of a 20% magnesium sulfate solution. The symptoms disappeared after a calcium salt solution was injected intravenously.
- (c) Byron in 1939 (113) reported the cases of 5 young Tamil children with helminthiasis (age range 2-1/2 to 10 years) who died after the oral administration of a magnesium sulfate purge (mistura alba). All the deaths were characterised by the rapidity with which they occurred following the magnesium sulfate ingestion. The only postmortem abnormality noted was general congestion. The blood serum of one of the dead children contained 8 mg/100 ml serum as contrasted to the normal level of 2-3 mg/100 ml. The authors speculate that greater magnesium absorption than ordinarily occurs can happen in certain cases and particularly with strong solutions.
- (d) Collins and Russell in 1949 (146) described a fatality owing to magnesium poisoning resulting from absorption of magnesium sulfate by the intestine, in a 4-year-old boy with primary megacolon. At autopsy the blood serum contained 30 mg Mg/100 ml. The authors considered the possibility that under certain conditions magnesium sulfate (Epsom salt) may become poisonous owing to its unexpected absorption. They note in

this regard that the gastrointestinal membrane is quite permeable to magnesium salts.

- (e) Stevens and Wolff (735) in 1950 reported two cases of toxicity, one of which was fatal when a 300 ml solution containing 189 g of magnesium sulfate was administered rectally over a 3 hour period by slow Murphy drip. These toxic reactions resulted from significant absorption of the salt from the rectum which produced depressant cerebral effects. The authors observed that normal kidneys may not remove the magnesium sulfate from the blood stream fast enough to prevent amounts toxic to the central nervous system from accumulating.
- (f) Rosler in 1952 (641) mentioned the case of one person (out of 12) treated for tapeworm with 80-120 g epsom salts (magnesium sulfate) dissolved in 200 ml water intubated into the duodenum, who died of central respiratory paralysis and magnesium narcosis. The author also mentioned being informed of another fatality resulting from curing tapeworm with magnesium sulfate. He pointed out that there are physiological conditions under which absorption of magnesium sulfate from the intestinal tract occurs. As a result the system is inundated with a large amount of magnesium leading to deep narcosis and paralysis of the respiratory center. He warned against the peroral use of large doses of epsom salts.
- (g) In 1969, Colomb et al., (147) described the case of a young man who developed caustic lesions on the interior and exterior surfaces of his cheeks and lips, when epsom salts which was used as smelling salts spilled on his face.

- (h) Rees (620) in 1970 reported the case of a pregnant female (aged 30) who apparently died from the effects of epsom salts which had entered the blood stream via the uterus in an attempted abortion. Abnormally high concentrations of  $Mg^{2+}$  ion were found in the heart blood (14.4 mg%) and in the peripheral blood (11.1 mg%). Prior to death, the woman allegedly reported a sensation of great heat passing through her body. This physiological phenomenon had been previously reported in connection with intravenous injections of a 42% solution of magnesium sulfate to volunteers.
- (i) Willner (827) reported in 1971 on the death of a woman accidentally administered magnesium sulfate. The woman (59 years old) was a hospital patient who in the course of diagnostic tests was given 400 ml of 40% magnesium sulfate by duodenal tube. She died 6 hours later. On autopsy it was found that the blood magnesium level was about 33%. The author compared this with other reported blood magnesium levels following death from magnesium poisoning and concluded that magnesium levels about 15% indicated a toxic content. He also considered his finding of 8-10% magnesium in the vitreous fluid as probably 2 to 3 times higher than is normally present.

2. Allison (012) discussed a toxic sulfate effect from water found in about 1/4 the state of Minnesota which is high in  $\text{MgSO}_4$  (as well as other sulfates). In the regions in which this high sulfate containing water (770 ppm to 3,590 ppm) occurs both man and livestock suffer from its drastic cathartic action and weakening effect. Calves are stunted in growth and many of the cattle die prematurely. Cattle from Dakota and Montana where similar water exists have been observed to be similarly affected. The author suggested that people may be affected in the same way but perhaps to a different degree. He urged a continuation of experimental studies in particular those concerning the effect on the equilibrium of calcium in the body.

3. Hirschfelder (323) studied the effects of oral administration of magnesium sulfate. Normal persons showed no rise in the level of magnesium in the blood plasma after ingesting epsom salts and thus had no ill effects. Patients with renal disease had entirely different results. When one or more purgative doses (25 g per dose) was administered orally there was a large rise in plasma magnesium within 4 to 6 hours. Many of the patients showed a syndrome of high plasma magnesium (hypermagnesemia, 9-11 mg/100 ml plasma) which was accompanied by somnolence or light coma. The author believed that epsom salts purgation in nephritic patients can induce a magnesium coma which is easily mistaken for uremic coma. The difference is significant because magnesium coma can be treated by intravenous injection of calcium chloride.

There is another condition, hypomagnesemia or low plasma magnesium characterised by convulsions or muscular twitching. This condition is relieved by oral administration of epsom salts. The author believed

that hypomagnesemia with its accompanying condition of hyperirritability of the neuromuscular system is probably more common than is realized.

4. Abramowitz and Russo (001) described the case of a "fixed" eruption caused by magnesium hydroxide. The patient had previously suffered cutaneous lesions and a rash on her hands after ingesting phenolphthalein and phenobarbital (0.06 g). Subsequently on taking either magnesium hydroxide tablets or magnesia magma (suspension of magnesium hydroxide in water) the eruption recurred.

5. Randall et al. (616) studied the relation of clinical manifestations of hypermagnesemia and serum magnesium level. Marked hypermagnesemia and severe manifestations of toxicity were developed by five patients with renal disease who were administered prescribed magnesium medications. (Table 9 summarizes the results.) Fourteen patients with chronic renal failure ingested 180 ml daily (6 equal 30 ml doses at 3 hour intervals on 3 consecutive days) of one of two magnesium antacids (either the hydroxide or the trisilicate). (The results are summarized on Table 10 ). Similar doses under the same circumstances were given to six normal subjects. (The results are summarized on Table 11.)



Table 9. Clinical and Laboratory Data in Patients With Toxic Manifestations of Hypermagnesemia (616)

Patient No.	Diagnosis	Creatinine Clearance ml/min	Hospital Day	Magnesium Intake		Serum				
				Oral	Paren-teral	Mg mEq/l	Ca mg/100 ml	PO <sub>4</sub> mg/100 ml	Na mEq/l	K mEq/l
1	Bleeding duodenal ulcer	12	5	Mg(OH) <sub>2</sub> <sup>c</sup>	MgSO <sub>4</sub> 40 mEq im <sup>d</sup>			2.1	145	2.6
			6		MgSO <sub>4</sub> 16 mEq iv <sup>d</sup>	7.5	5.1	2.6	148	3.7
			3-7		MgSO <sub>4</sub> 116 mEq im <sup>d</sup> in 5 days		6.7	5.0	126	3.2
2	Acute pyelonephritis with papillary necrosis	30	13			1.4			132	5.4
			14-16		MgSO <sub>4</sub> 220 mEq im <sup>d</sup> in 3 days	7.2	7.2	5.0	138	5.0
			21			3.0	7.4		138	5.0
3	Malignant hypertension	10	6	MgSO <sub>4</sub> (laxative)			6.8	9.6	122	3.8
		<1	10			4.2				7.3
			15			5.4	6.0		140	8.4
4	Cirrhosis with oliguria		1	Mg(OH) <sub>2</sub> <sup>c</sup>					135	3.3
		20	5			9.1		4.3	120	5.5
		10	6			7.6	6.6	3.4	121	5.7
5	Diabetic glomerulo-sclerosis		1	Mg(OH) <sub>2</sub> <sup>c</sup>			8.9	7.4	138	5.3
		7	10			5.0	8.9			7.1
			14			5.4	7.6	7.9	130	8.4

aNPN rather than BUN.

bAll interval measurements are in seconds.

cGiven as Maalox, 120-300 ml daily.

dAdministered as 50% MgSO<sub>4</sub>·7H<sub>2</sub>O as prepared by Lilly and Co.

eArterial pCO<sub>2</sub> 39 mm Hg at this time.

Table 9 (Con't)

Patient No.	Serum		Creatinine mg/100 ml	pH (art:)	Blood		Urine ml/day	Electro- cardiogram	Toxicity
	Cl mEq/liter	CO <sub>2</sub> mM/liter			Urea Na mg/100 ml	Hematocrit Reading %			
1	85	26			62 <sup>a</sup>	43		PR, A6, QRS .09, QT <sub>c</sub> .49, T waves flat	Following injections: coma, areflexia, apnea, hypotension Apnea and death after injection
	80	41	7.0	7.49	100 <sup>a</sup>	48	>1,000		
2	84	14			60	26		Normal	First course of magnesium reduced symptomatology presumably related to magnesium depletion
	112	17			36				
	106	15	3.2		38		>1,000	Nodal rhythm, QRS .16, QT <sub>c</sub> .47, T waves inverted	Second course of injections followed by arrhythmia, transient asystole, hypotension, coma, decreased res- pirations
	107	17			73	26		Normal	Improved as serum magnesium level fell
3	92	17	4.8		90 <sup>a</sup>	29	200	PR .16, QRS .10	
	90	8	53		228 <sup>a</sup>		0	PR .22, P waves flat, QRS .13, T waves tall	P-R prolongation. Coma and death probably related to severe uremia
4	73	26	4.0		81	30	300		
	60	36			130		100		Progressive coma, hypotension, di- minished reflexes and respirations
	60	37 <sup>e</sup>	6.6	7.59	135	36	50		
5	106	15	11.0		90	33		PR .15, QRS .08, QT <sub>c</sub> .41, PR .32,	Progressive drowsiness, confusion, hypotension, dysarthria, difficulty voiding, diminished respirations, P-R prolongation Death
	104	15			110		600	QRS .16, QT <sub>c</sub> .50, ST depressed, T waves tall	
	99	14	12.0		125	33	1,000	Same	

Table 10. Clinical and Laboratory Data in Patients with Renal Disease Ingesting Magnesium-antacids Experimentally (616)

Patient No.	Diagnosis*	Creatinine Clearance	Medication†	Serum Magnesium		Blood Urea N	Serum			Urine Magnesium		Toxicity**
				Before Antacid	After 3 days		Cr	Ca	P <sub>OH</sub>	Before Antacid	14 hr. Day	
		ml/min		mEq/liter		mg/100 ml	mg/100 ml			mg/24 hr		
6	Chronic glomerulonephritis (b)	30	Maalox	2.0	3.0	36	3.2	8.1	4.6	61	170	Nausea
7	Nephrosclerosis (a)	—	Maalox	1.8	4.3	116	9.0	5.0	12.3			Steady, nausea
8	Diabetic glomerulosclerosis (b)	11	Gelasil	2.1	3.4	106	7.0	8.7	—	84	136	M
9	Chronic glomerulonephritis (b)	29	Maalox	2.8	3.6	75	5.2	7.1	4.8			
10	Diabetic glomerulosclerosis (b)	28	Maalox	2.4	3.0	62	2.4	9.7	5.3			
11	Diabetic glomerulosclerosis (c)	8	Maalox	2.4	2.6	150	11.0	8.9	7.6	65	82	
12	Chronic glomerulonephritis; congestive failure (c)	10	Maalox	2.3	3.9	108	5.6	8.1	5.3			Drowsy
13	Nephrosclerosis; nephrocalcinosis (a)	8	Maalox	1.7	3.5	100	5.0	7.0	3.3			
14	Amylloidosis (a)	—	Maalox	2.0	2.7	40		8	4.7			
15	Nephrocalcinosis (Milk-alkali) (c)‡	55	Gelasil	2.2	2.5	37	1.5	13.3	3.0	125	300	
16	Nephrocalcinosis (Milk-alkali) (b)	30	Maalox	2.4	3.5	31	3.2	13.0	2.7			
17	Chronic pyelonephritis (a)	8	Maalox§	2.6	4.9	100	7.9	7.3	6.4	50	120	Nausea, drowsiness, malaise; difficult micturition and defecation; muscle twitches; postural hypotension, slight P-R prolongation
18	Chronic glomerulonephritis (a)	4	Maalox	1.8	3.2	200	16.4	7.9	8.1	32	44	Drowsiness
19	Adenocarcinoma prostate (c)	—	Maalox	1.8	3.6	60		8.3	2.7			

\*Established by autopsy examination (a), renal biopsy (b), or clinically (c).

†See subscript Table 3.

\*\*Toxic manifestations began during the period of medication and subsided upon stopping the medication as the serum magnesium concentration fell.

‡Reported in detail elsewhere (21).

§ Similar results were obtained during the ingestion of a comparable amount of magnesium hydroxide as Milk of Magnesia (USP).

Table 11. Serum and Urine Magnesium in Normal Subjects Ingesting Antacids (616)

Subject No.	Creatinine Clearance	Antacids*	Before Antacid	Serum Magnesium		Urine Magnesium	
				After 3 days	Before Antacid	Third Day	
	<u>ml/min</u>	<u>180 ml/day</u>		<u>mEq/liter</u>		<u>mg/24 hr.</u>	
1	110	Maalox	1.9	2.1	122	234	
2	120	Maalox	2.1	2.0	87	284	
2	120	Gelusil	2.0	2.0	85	400	
3	100	Maalox	2.2	2.3			
4	115	Maalox	1.9	2.0			
5	130	Maalox	1.7	1.8	115	270	
6	140	Gelusil	2.3	2.5	125	306	

\*Provides approximately 2.8 g (235 mEq) of magnesium as  $\text{Mg}(\text{OH})_2$  in Maalox and 2.9 g (240 mEq) of magnesium as magnesium trisilicate in Gelusil.

The authors pointed to several observations for particular emphasis:

- (a) The serum concentrations at which toxic manifestations appeared in patients with renal disease were much lower than those in the normal subjects.
- (b) Patients with renal failure have a limited capacity to excrete magnesium.
- (c) The potential danger of magnesium absorbed from ingested laxatives and antacids in patients with renal disease has not been fully appreciated.
- (d) Patients with severe renal failure who ingest customary doses of standard laxative and antacid preparations, may develop significant deviations of serum magnesium accompanied by manifestations of toxicity within 3 days.

6. Brady and Williams (098) reported the case of a premature infant with hypermagnesemia. The mother was administered magnesium sulfate (a total of 11 gm) prior to the infant's birth. The infant had a high serum magnesium level (15 meq/l at 9 and 16 hours and 9.8 meq/l at 24 hours) and showed severe symptoms of motor and respiratory paralysis. The condition was cured by exchange transfusion. The authors concluded that the magnesium sulfate therapy given to the mother had profound effects on the fetus.

7. Lipsitz and English (448) reported on sixteen cases of hypermagnesemia in newborn infants. All of these infants were born to mothers given magnesium sulfate for toxemia. In both the maternal and fetal serums, magnesium levels were higher than expected. The clinical

manifestations of the newborn were similar to those of adults with hypermagnesemia. The authors urged awareness of this problem.

8. Stone and Pritchard (736) on the other hand reported that they had not found any deleterious effects from magnesium sulfate administered to mothers prior to delivery. They claimed that over a 14 year period, 7000 infants were born of mothers who had received magnesium sulfate therapy. They further claimed that even though the serum level of magnesium in the fetus rapidly approached that of the mother, it could not be correlated with any ill effect. The authors specifically cited the previously reported study of Lipsitz and English.

9. Lipsitz (447) warned again that more detailed study has shown that if continuous intravenous infusion of magnesium sulfate is given to the mother for more than 24 hours, the newborn will manifest all the signs of hypermagnesemia. Two routes of magnesium sulfate administration were studied (27 mothers, intravenously; 8 mothers, intramuscularly after initial intravenous injections). The author concluded that when toxemic mothers are administered a large dose of magnesium sulfate or prolonged magnesium sulfate therapy, there is great likelihood that their infants will manifest the symptoms of magnesium toxicity. These symptoms which are similar to those described in adults (see Randall et al. 616, Biology Section II, B, 5) with hypermagnesemia are unlike those born to toxemic mothers without magnesium sulfate therapy. Magnesium toxicity should be recognized and treated in such newborns.

### III. Long-Term Toxicity

No information available.

#### IV. Special Studies

##### A. Frogs

Chiray et al. (138) studied the effect of magnesium chloride on the frog intestine in vivo. A cannula was introduced via a lateral opening in frogs with previously destroyed medullas (number and sex not given). By making a second opening 2 or 3 cm away to allow the perfusion liquid to exit, the first 2 or 3 cm of the small intestine could be studied. In the first part of the experiment Tyrode's solution at 38° was perfused and the number of drops passing the section of intestine being studied during one or more minutes was counted by means of an electric drop counter. The perfusion was carried out at a constant pressure and temperature (10-15 cm of water at 38°). Then the Tyrode's solution was replaced by water containing magnesium chloride (from Chatel-Guyon). In a few minutes there was an approximate 70 percent decrease in the drop flow from 50-55 per minute to about 15. This reduced flow which was established in a few minutes remained as long as the perfusate contained magnesium chloride. When this was replaced by physiological solution, the flow returned to its former rate after a few drops.

The authors relate this effect, which they ascribe to a strong amplification of normal peristaltic waves, to what happens when magnesium chloride exerts a laxative effect in man.

They point out that the large doses (1500 cc and up daily) which were previously administered therapeutically at Chatel-Guyon have been discontinued because they cause diarrhea. Present dosages of a few hundred

grams which are used to regulate intestinal function apparently play a decontracting role in the case of spasmodics and a contracting role with atonics. The authors suggest it would be of interest to study the effect of magnesium chloride containing water on an intestine contracted by various pharmacological agents.

#### B. Mice

1. Labkovsky (424) studied the effect on the induction of pulmonary adenomas with urethan in mice, when magnesium salts (chloride or sulfate) are administered orally or subcutaneously. All the experiments were carried out on stick-bred white mice (19-20 g, number and sex not given).

The six oral experiments, each with its own controls were as follows:

- (a) Starting 2 weeks prior to first urethan injection (1 g/kg in 5% solution administered intraperitoneally twice per day every 4 days),  $\text{MgCl}_2$  (320 mg/kg) was given in drinking water for 10 weeks.
- (b) The same amount of  $\text{MgCl}_2$  was administered via stomach tube. Administration started one day prior to first urethan injection (same as 1) and continued for 10 days.
- (c) As above.
- (d) As above.
- (e) Everything the same as in 2, 3, and 4 except  $\text{MgSO}_4$  administered.
- (f) Everything the same as in 2, 3, 4, and 5 above except  $\text{MgSO}_4$  administered subcutaneously.



The results showed the following:

Experiment	Percent when average no. of pulmonary adenomas per control mouse taken as 100%
a	135
b	216
c	215
d	112
e	194 and 196
f	173

In another series of four experiments  $\text{MgSO}_4$  was injected subcutaneously daily (5 ml/kg of 0.5% solution) for 10-15 days. These experiments were as follows:

- (1) Urethan (same amount as above) administered 2 days after cessation of  $\text{MgSO}_4$  injections.
- (2)  $\text{MgSO}_4$  injections started 10 days after second urethan injection.
- (3)  $\text{MgSO}_4$  injections started 2.5 months after urethan administration.
- (4) Control.

The results were:

Experiment	Percent when average no. of pulmonary adenomas per control mouse taken as 100%
a	174
b	146
c	114

The author concluded:

- (1) Magnesium salts (chloride or sulfate) stimulated the induction of pulmonary adenomas by urethan in mice.
- (2) The method of administration or the composition of the salt (chloride or sulfate) did not have any effect.

2. Bazikyan and Akimov (051) studied the antineoplastic effect of magnesium chloride in mice under experimental conditions. A total of 259 mature male CC57BR mice were used in all the experiments described below. Skin papillomas were induced by rubbing a 0.1% benzene solution of 9,10-dimethyl-1,2-benzanthracene (DMBA, 50 mg/kg per mouse) on the posterior portion of the spine on alternate days for one month (15 applications). Two series of experiments were carried out as described below:

- (a) In series 1, there were 2 subgroups. In one the mice were given drinking water to which magnesium chloride (60 mg/kg/day) had been added throughout the entire time the carcinogen was applied (30 days). The other group which were controls received ordinary drinking water. (See Table 12 for details and results.)
- (b) In series 2, there were 4 subgroups. One was a control group similar to the one in series 1 above. Subgroup 1 received  $MgCl_2$  (60 mg/kg/day) as above 7 days prophylactically and 30 days with DMBA. Subgroup 2, received  $MgCl_2$  as in subgroup 1 but with DMBA for 53 days. Subgroup 3 received the  $MgCl_2$  from the eighth day after the first application of DMBA and for 48 days thereafter, when the first papilloma appeared. See Table 12 for details and results.

Some observations were:

- (a) Magnesium chloride showed definite antitoxic action in series 2 subgroup 2 in particular, in which  $MgCl_2$  was administered for the longest time. (See Table .)

Table 12 a. Effect of Magnesium Chloride on the Origin of Skin Papillomas in Mice (051)

Series of experiments	Animal subgroups	No. of mice			p
		at start of expt.	at end of expt.	died %	
I	Control	30	22	27	0, 5
	Experimental	30	24	20	
II	Control	29	19	35	0, 3 0, 05
	Experimental 1	30	24	20	
	Experimental 2	31	27	10	
	Experimental 3	29	17	42	

Table 12 b.

Series of experiments	Animal subgroups	No. of mice with tumors		p	Avg. no. of tumors per mouse		p
		abs.	in %		abs.	% of control	
I	Control	18	82	0, 5	1, 1 $\pm$ 0, 16	100	0, 1
	Experimental	16	66	0, 3	0, 80 $\pm$ 0, 11	73	
II	Control	14	74		0, 90 $\pm$ 0, 17	100	0, 1 0, 07 0, 7
	Experimental 1	12	50	0, 1	0, 63 $\pm$ 0, 12	70	
	Experimental 2	11	41	0, 05	0, 55 $\pm$ 0, 14	51	
	Experimental 3	11	65	0, 05	0, 82 $\pm$ 0, 14	90	

- (b) On the other hand in subgroup 3 of series 2, 42% of the mice died. The authors postulated that this was due to the 8-day delay in administration of  $MgCl_2$ .
- (c) In all the groups of both experimental series the first tumors appeared between days 48 and 54. Thereafter there was a different rate of increase.
- (d) The experimental mice in series 1 had fewer tumors than the controls throughout the experiment.
- (e) Subgroup 3 of series 2 had the largest number of animals of all that received magnesium, both with respect to skin papillomas and number of tumors per mouse. The death rate was also highest in this group. The authors conclude from this observation that: (a) magnesium chloride must be administered both for several days before topical application of the carcinogen and concomitantly with it, in order to manifest an antitoxic and antineoplastic action and, (b) that irreversible changes in the cells which later become malignant are produced from the very first application of carcinogen.

In another series of experiments the effect of magnesium chloride on connective tissue tumors was studied. The connective tissue sarcomas were induced in all 3 experimental groups by a single injection of 3 mg of 1,2,5,6-dibenzanthracene (DBA).

The controls received tap water throughout the duration of the experiment (200 days). Group 1 drank water with added magnesium chloride (60 mg/kg) for 30 days and group 2 for the duration of the experiment. (See Table 13 for details and results.)

Table 13. Effect of Magnesium Chloride on the Induction of Subcutaneous Sarcomas in Mice (051)

Groups of animals	No. of mice			p	No. of mice with tumors		p
	at start of expt.	at end of expt.	died %		abs.	in %	
Control	30	18	40	0, 3	15	83	
Experimental 1	19	15	21	0, 02	10	66	0, 2
Experimental 2	21	19	10	0, 05	7	37	0, 01

The observations were:

- (a) The group with the highest mortality and the first animals to die from the toxic effects of DBA were the controls, as in the previous experiments.
- (b) Also as in the previous experiments, the group with the lowest mortality had the smallest number of mice with tumors. The authors concluded from this experiment that the ingestion of  $\text{MgCl}_2$  lowered the number of DBA induced sarcomas.

They concluded from all the above experiments that:

- (a) Under the conditions of the experiments and in the amount administered,  $\text{MgCl}_2$  reduced the toxicity of DMBA and DBA.
- (b) Under their experimental conditions  $\text{MgCl}_2$  markedly decreased the number of skin papillomas induced by DMBA and subcutaneous sarcomas induced by DBA.

#### C. Rats

Wies (826) reported on experiments to determine whether magnesium sulfate is teratogenic in rats. A series of experiments were carried out using 34 albino rats (18 female and 16 male, 120-200 days old). The females were given subcutaneous doses of sterile magnesium sulfate solution ( $0.8 \text{ mg/kg MgSO}_4 \cdot 7\text{H}_2\text{O}$ ) on alternate days (for approximately 8 weeks). The females were all mated at least once.

It was observed that:

- (a) The fertility of females receiving magnesium sulfate was low (15 percent).
- (b) Varying degrees of toxicity were produced by the magnesium sulfate including death.

(c) Defects in the brain and spinal cord of at least one fetus were observed. (It is not stated how many fetuses were produced nor in how many of those produced were the described defects observed.) Specifically the defects consisted of a diminution of the number of neuroblastic cells in a circumscribed area of the midbrain. In the spinal cord the lateral walls had not fused over the central canal leaving it open.

#### D. Guinea Pigs

Chiray et al. (138) carried out the same experiment with guinea pigs in which the action of magnesium chloride containing water on the intestine in vivo was studied as has already been described in detail with frogs, in A, 1 in this section.

Before inserting the cannula the guinea pigs (number and sex not given) were lightly anesthetized with ether. In the guinea pig experiment the section of intestine studied was the end of the ileum and the ileocecal valve. The original perfusion solution was not specifically specified but referred to as physiological perfusion liquid.

The main differences in the response of the guinea pig were that the rate of flow was less slowed down (from 38 drops per minutes to 28 as compared to 50-55 drops per minute to 15 in the frog) and return to the original rate of flow (with physiological solution) took longer to reestablish than with the frog. In all other respects the results with both species were identical. Perfusion of a small section of the small intestine with water containing magnesium chloride caused the flow of the perfusate to be slowed down owing to the contraction of the smooth

fibers of the intestine. This contraction was caused by a strong exaggeration of the normal peristaltic waves.

#### E. Rabbits

1. Serbescu (686) studied the influence of magnesium sulfate on tar-induced carcinomas. Two groups of 30 rabbits each were used (age and sex not given). One served as controls and the other was administered magnesium sulfate by stomach probe (20 mg/kg, dissolved in redistilled water) every day except Sunday. All the animals were painted on the inside of both ears with tar oil three times per week. Animals in both groups which died were immediately replaced, thus in the course of the experiment a total of 123 rabbits were used (75 experimental and 48 controls). Both pathological studies (the digestive tract and the ganglionic system) and histological tumor examination were carried out. The following were noted for the magnesium treated animals:

- (a) A higher mortality rate.
- (b) Cancer developed more easily.

In the same paper Mr. P. Delbet submitted a rebuttal which made essentially the following points:

- (a) His own experience with the magnesium halogen salts has indicated a restraining effect on cancer development.
- (b) Magnesium sulfate would not necessarily behave similarly for two reasons: (a) it is not as well absorbed as the chloride, (b) the sulfate tends to make the urine alkaline whereas the halogen salts reduce its pH. Delbet does not explain why he considers this significant.



- (c) The described method of sulfate administration would probably weaken the animals both by acting as a purge and by preventing them from getting adequate nourishment.
- (d) Delbet's experimental work in which he demonstrated a reduction of tumor formation when animals were injected with a mixture of magnesium halogen salts, was repeated successfully. The experimenter, Braier, additionally determined that the C/N ratio in the urine of cancerous rats was lowered in the controls but remained normal in the magnesium treated animals. Delbet concludes from this that the poor utilisation of fats and carbohydrates is a factor in cancerous cachexia and that large doses of magnesium halogen salts have a therapeutic effect.

2. Neal and Neal (537) studied the effect of hard water and water containing magnesium sulfate on atherosclerosis in rabbits. Four groups of New Zealand rabbits (5 per group, age and sex not given) were given drinking water as follows for 3 months:

Group I - distilled water

Group II - hard sulfur water

Group III - one tablespoon of calcium carbonate suspended in one gallon of distilled water

Group IV - 7 tablespoons of magnesium sulfate (Epsom salt) dissolved in one gallon of distilled water (30,000 ppm).

The diet of all groups was Purina rabbit pellets with 1% crystalline cholesterol and 5% butter. The aortas and hearts of all the rabbits were studied.

The following was observed:

- (a) Group I rabbits (distilled water) had the most atherosclerosis.
- (b) Group II and Group III rabbits had moderate atherosclerosis.
- (c) Group IV rabbits (magnesium sulfate) had no atherosclerosis, did not have diarrhea and were larger and fatter (more subcutaneous fat) than the rest of the rabbits.
- (d) Although the blood serums of rabbits from each group contained the same amount of fat, the serums from the Group IV rabbits (magnesium sulfate) were relatively clear while those from the other 3 groups were turbid or "milky".

The authors raised the questions of whether there is a protective substance in hard water or whether there is a specific effect of magnesium on fat metabolism. They suggested the necessity for further studies using other ionizing compounds.

#### F. Dogs

1. Hazard and Wurmser (308) studied the effect of magnesium chloride on the excitability of the sympathetic nervous system of dogs. Male dogs (6.6 kg, number not given) were administered  $MgCl_2$  intravenously at various strengths (0.01 g/kg to more than 0.10 g/kg).

It was found that:

- (a) Weak doses of  $MgCl_2$  (0.01-0.05 g/kg) slightly increased the heart's response to excitation by accelerators.
- (b) Strong doses (0.10 g/kg and up) had the opposite effect.
- (c) In vagotomized or atropinized dogs large doses of  $MgCl_2$  increased the accelerating effect of adrenaline and lessen the hypertensive action of nicotine.

- (d) A depressant effect on the viscera, particularly the kidney, resulted from large  $\text{MgCl}_2$  doses.

2. Maxwell et al. (490) studied the effects of hypermagnesemia on the hearts of intact dogs. Hypermagnesemia was induced in 6 dogs (13-30 kg, sex not given) by intravenous injection of a 10% solution of magnesium chloride. In an unstated number of other dogs 2 ml/kg of 10% magnesium chloride was administered by infusion. Before and after the induction of hypermagnesemia, cardiac output, coronary flow, and vascular pressures were measured. The main changes which took place were tachycardia and a fall in systemic pressure. The authors postulated that under the conditions of the experiments the resulting tachycardia and hypotension appeared to be associated with variation in the calcium-to-magnesium ratio owing to the increase in magnesium.

#### G. Humans

1. Bernstein and Simkins (076) studied the intravenous effects of therapeutic doses of magnesium sulfate on the heart. One hundred adult patients were used in the study, 34 apparently free of cardiovascular disease (but with various other diseases) and 66 with various forms and degrees of cardiac disease (see original paper for details). Warmed magnesium sulfate solution (10 cc of a 10% aqueous solution) was injected into a large antecubital vein of the patient. Three sets of electrocardiograph tracings (one from each lead) were made for each patient.

The authors considered that the most salient feature of their study was the lack of uniformity in the changes in the heart brought about by the magnesium sulfate injections. In a fairly high percentage of cases there were slight changes in QRS complexes and T waves indicating a direct effect on the

myocardium of the ventricles. The authors concluded that intravenous magnesium injections exerted no deleterious effect on the human heart.

2. Boyd and Scherf (092) studied the therapeutic use of magnesium sulfate in paroxysmal tachycardia. Eleven patients (10 cases of paroxysmal tachycardia and 1 case of flutter) were treated by intravenous injection of magnesium sulfate in concentrations of 10% or 20%. The results were favorable in 3 out of 8 cases for the 10% solution. The effect was immediate. It was more successful at the higher concentration (15 to 20 cc of a 20% solution); 8 out of 8 cases. The authors considered this procedure safe but cautioned against its use in the presence of: marked myocardial damage, marked intraventricular conduction, or gallop rhythm.

3. Ozsoylu (565) examined the effect of giving magnesium sulfate to control blood sugar levels in diabetic children. Three children (from 5-14 years) were given an intravenous infusion of a 3% magnesium sulfate solution (150 mg/kg). The blood sugar levels were determined on the three days prior to magnesium sulfate therapy as well as on the fourth day when the magnesium sulfate was given. The same amount of insulin was given on all four days. Three patients (controls) who did not have diabetes were given the same amount of magnesium sulfate on the fourth day following three days of checking their blood sugar levels. It was found that the blood sugar level of all the diabetic patients dropped considerably (116 mg%, 195 mg% and 234 mg%) within one hour following the infusion of magnesium sulfate. It appeared (see Table in original paper for details) that the higher the blood sugar level the greater effect the magnesium sulfate had on decreasing it. There was no effect on the normal blood sugar levels of the controls, from the magnesium sulfate infusion. The author expressed

the hope other investigators will be interested to gather further data since there are not too many diabetic children in Turkey.

4. Frewin et al. (237) attempted to clarify the vasodilator action of magnesium sulfate. Eight volunteer subjects (7 medical students and one patient with a sympathetically denervated forearm) were used. Intra-arterial infusions of magnesium sulfate (20 and 50 mg/min) were given into the brachial artery at the elbow of one side. Forearm or hand blood flow was measured.

The results of the experiments demonstrated that:

- (a) Magnesium sulfate has a direct local vasodilator action on the blood vessels of the forearm.
- (b) The vessels of both the skin and the muscle are affected.
- (c) The local direct action on the blood vessels was further confirmed by the presence of a vasodilator action in the patient with the sympathetically denervated forearm.
- (d) Administration of propranolol, hyoscine or mepyramine did not modify the response. This indicated that the dilator action is not caused by stimulation of  $\beta$ -receptors, a cholinergic mechanism or the release of histamine.
- (e) The simultaneous administration of an equimolar dose of calcium gluconate, a vasoconstrictor, abolished the dilator action of magnesium sulfate.

## BIOCHEMICAL ASPECTS

### I. Breakdown

The breakdown of the various magnesium compounds in the body is discussed in the following sections. No breakdown in storage, processing, or cooking has been noted in current literature.

### II. Absorption and Distribution

#### A. Rats

Alcock and MacIntyre (007) studied the interrelation of calcium and magnesium absorption by studying the effect of calcium deficiency on magnesium absorption in rats.

Female rats (28 animals, 90-110 g) were tube-fed diets deficient in calcium for 18 days. A control group (26 animals) received a diet (also by tube) with calcium present in a normal amount (see Tables 14 and 15 for details). Balance techniques were used to study the effect of dietary calcium deficiency on the absorption of magnesium. (See original paper for details of method.)

The excretion of magnesium by the calcium deficient animals was markedly diminished. There were no abnormal signs in these animals apart from the fact that the feces were soft and less bulky than normal. Also the fecal dry weight was diminished. The mean daily excretion of magnesium and calcium together with that of the normal control animals is shown below.

Table 14  
The Electrolyte Content of the Diets (007)

Type of diet	Mg	Ca	K	Na	Phosphate
Normal	34	300	145	45	100
Ca deficient	34	3.0	145	45	100

(Magnesium, calcium, potassium and sodium content is expressed in terms of m-equiv/kg of diet; phosphate content is expressed in terms of m-moles/kg of diet.)

Table 15  
The Daily Intake ( $\mu$ -equiv/rat) of Calcium and Magnesium (007)

Diet	Magnesium	Calcium
Normal Control	272	2400
Calcium Deficient	272	2

Mean daily fecal excretion of calcium and magnesium (u-equivalents).  
Results are expressed as mean (number of determinations)  $\pm$  S.E.M.

<u>Excretion</u>			
Experimental group	<u>Magnesium</u>	Experimental group	<u>Calcium</u>
	Normal group		Normal group
84.9(26) $\pm$ 6.4	154.0(28) $\pm$ 6.3	6.2(26) $\pm$ 0.95	1684(28) $\pm$ 57.4

The authors concluded that their results strongly suggested the existence of a common transport mechanism for the transfer of calcium and magnesium across the intestinal wall. They considered the increased magnesium absorption from the gut in calcium deficient diets as evidence in support of this mechanism. They suggested both that a similar mechanism may exist in the kidney and that the possibility exists that such a common absorptive mechanism for calcium and magnesium in both the intestine and renal tubule may exist in many different species.

#### B. Rabbits

Aikawa and Burns (004) investigated the maternal-to-fetal transfer of magnesium and its uptake by various maternal and fetal tissues in the rabbit. The radioactive isotope  $^{28}\text{Mg}$  (as magnesium sulfate solution) used in the study was injected into the marginal vein of the ears of eleven pregnant domestic albino rabbits (between 28-30 days of gestation). A total of 86 fetuses and their placentas were removed from the rabbits at intervals ranging from 7 minutes to 26 hours.

The observations were as follows:

- (a) Maternal tissue uptake of  $^{28}\text{Mg}$  resembled that previously found in nonpregnant adult rabbits with the exception that bone and muscle uptake was slower.



- (b) Placental concentration of  $^{28}\text{Mg}$  rapidly rose above the maternal plasma level.
- (c) By 26 hours the specific activity of magnesium in all the studied fetal tissues reach a fairly constant value.
- (d) The turnover of magnesium in fetal tissues in utero, particularly in bone and muscle, is considerably more rapid than in the respective maternal tissues.

### C. Dogs

1. Smith et al. (712) examined the distribution of magnesium in dogs after injection of magnesium sulfate. Adult female dogs (number not given) were injected in the femoral or jugular vein with an isotonic magnesium sulfate solution (0.154 M at a rate of about 10 cc per min.). All urine and feces were collected for the next 24 hours. The concentration of magnesium in the serum preceding and following injection was compared. The retention of any magnesium in the body was determined from these data and the amount of magnesium still remaining in the extracellular fluid calculated. The authors postulated that if some of the retained magnesium could not be accounted for in the extracellular fluid, it would have to have been deposited elsewhere.

It was observed that:

- (a) In the first 3 or 4 hours after injection the volumes of distribution of both magnesium and sulfate ions were nearly identical. This indicates that both ions are distributed throughout the same portion of the extracellular fluid (20 to 25% of the body weight). Therefore magnesium distribution resembles that of sodium rather than potassium.

(b) All the injected magnesium could not be recovered in the urine and stools within 24 hours. In fact the authors did not believe there was any evidence that any of the injected magnesium was excreted via the feces. Therefore they concluded that sometime between the 4th and 24th hour after injection a variable proportion of the injected magnesium over and above the amount which can be accounted for in urine and stools, may leave the extracellular fluid and be segregated in an unknown form in undetermined places in the body.

2. Barbour and Winter (038) investigated both the absorbability of magnesium administered as soluble organic and inorganic salts (gluconate, lactate and oxide) and the question of magnesium and calcium storage.

In seventeen feeding experiments with 5 dogs (age and sex not given) a uniform increase of about 1 mg% in serum magnesium was observed when doses of 200 mg/kg BW of oxide, lactate or gluconate were given. Smaller doses were less effective (see Table 16). Since the calcium content of the serum was reduced in only one case, the authors considered this an indication that a single feeding of a large amount of magnesium does not displace calcium from the blood. Another observation suggested by these experiments (see Biochemical Table 16) was that gram for gram the three compounds tested were practically equivalent in their capacity to increase serum magnesium. This is despite the fact that the relative amount of magnesium in these compounds; gluconate, lactate and oxide, is about 1:2:12.

As a result of these experiments as well as other experiments performed in their laboratory by co-workers and quoted in this paper, the authors concluded that:

Table 16 Maximum changes in Serum Ca and Mg After Oral Administration of MgO, Mg gluconate or Mg lactate (038)

Amount of dose (mg/kg)	Oral dose of MgO		Oral dose of Mg gluconate		Oral dose of Mg lactate	
	Ca mg%	Mg mg%	Ca mg%	Mg mg%	Ca mg%	Mg mg%
50			10.9-12.0	4.0-4.0		
			10.6-10.8	4.0-4.0		
			10.1-10.8	4.4-4.4		
100	11.0-11.8	3.6-3.7				
	10.3- 9.5	3.8-3.9				
	10.6-10.9	4.0-4.5				
200	10.1-11.0	4.6-5.2	10.7-11.4	4.4-5.5	10.2-12.8	4.2-5.2
	10.8-12.3	4.3-5.4	10.1-11.1	4.2-5.4	11.0-12.5	4.6-5.2
	10.8-11.4	4.8-6.1*	9.1-10.4	4.4-5.0	10.2-10.8	4.1-5.5
			10.6-10.5*	3.8-4.0*	10.4-10.6*	3.7-4.0*

\*Two hours only.

- (a) The absorptive capacity for magnesium cannot be regarded as constant. Ordinarily harmless amounts of substances can for example be absorbed from a diseased or irritated intestine with unexpected results.
- (b) The form in which magnesium is ingested can determine both its rate and total amount of absorption.
- (c) Both soluble organic and inorganic magnesium salts markedly raise the level of serum magnesium. When large amounts of orally administered organic magnesium salts are administered, from one- to two-thirds may be readily absorbed.

A series of feeding experiments with magnesium gluconate and lactate in which the dietary levels of magnesium calcium and phosphorus were varied (adequate to low) was also carried out. From these the authors concluded that:

- (a) When the phosphate intake was adequate, prolonged administration of large amounts of soluble organic magnesium salts was not accompanied by loss of serum calcium.
- (b) When the phosphate intake exceeded 4 mmoles/kg BW/day, additional calcium was retained. (See original paper for tables.)

#### D. Humans

1. Wacker and Parisi (309) summarized some of the research findings on magnesium absorption. Magnesium is the fourth most abundant cation in the body (21-28 g in an adult) and the second most plentiful intracellularly. Bone contains about half the total body magnesium. The remainder is almost equally distributed between muscle and the nonmuscular soft tissues. Liver and striated muscle have the highest concentration (15 to 20 mEq/kg) of the non-osseous tissues. The concentration of magnesium in erythrocytes varies from 4.4-6.0 mEq/liter.

Calcium influences the amount of magnesium absorbed from the dietary intake (ca. 20-25 mEq/day) apparently by competition for a common absorptive pathway. About one-third of the daily ingested magnesium is absorbed.

It has been shown from animal studies that magnesium is absorbed mainly in the proximal part of the small intestine. In man the time course of appearance of orally administered labeled magnesium ( $^{28}\text{Mg}$ ) in the plasma is consistent with this interpretation. Peak levels of the isotope are measured between two and eight hours after the dose. However, after rectal enemas, colonic absorption can occur as evidenced by the development of hypermagnesemia (see Biological Section II B).

2. Silver et al. (701) examined magnesium turnover in man with  $^{28}\text{Mg}$ . The authors noted that both the use of this isotope and interpretation of results obtained with it should be viewed with caution because of its relatively low specific activity.

Ten adults (white, 3 males and 7 females) ranging in age from 38 to 72, were infused intravenously or orally administered a specially prepared  $^{28}\text{Mg}$  solution (pH 7.4 in HCl) in amounts ranging from 20 to 104  $\mu\text{c}$  (total volumes ranged from 25 to 100 ml). Each patient was on a consistent diet for 5 days before and 3 days after administration of the isotope. The magnesium dietary intake was fixed at about 50-60 mEq/day for each patient.

The chief observations following intravenous administration were that the isotope appeared in the urine rapidly but only insignificant amounts appeared in the stool. The authors interpreted this as indicating that most of the magnesium in the stools is of exogenous origin.

At 40-60 hours only about 10-25% of the body's total magnesium had equilibrated with the isotope and at 90 hours about one-third had. Thus indicating that equilibration of the isotope with the magnesium in the body is slow.

The authors concluded that their studies tended to confirm earlier data from magnesium turnover studies with nonisotopic techniques which showed:

- (a) There are several relatively small rapidly equilibrating compartments. In man there are at least three compartments in the magnesium body pool which turnover with half-times of 1, 3 and 14 to 35 hours respectively.

(b) There is one or more compartments in which turnover is very slow.

In man probably 25-50% of the magnesium has a turnover rate of less than 2 percent per day.

(c) Studies with  $^{28}\text{Mg}$  in man suggest that gastrointestinal absorption of  $\text{Mg}^{2+}$  is very limited.

However, the authors pointed out that these turnover values must be interpreted with caution because of possible effects from the significant amounts of dietary magnesium in addition to the effect of that administered with the low specific activity  $^{28}\text{Mg}$ .

3. Hirschfelder (323) studied the effects in humans of taking a single ordinary purgative dose of epsom salts. First they administered this dose of magnesium sulfate (within 24 hours after ingestion) to 7 normal men. They found that within 24 hours, 40-44% of the ingested dose appeared in the urine. However, despite the large quantity of absorbed magnesium the concentration of magnesium in their blood plasma remained relatively constant.

However, when a similar oral dose of magnesium sulfate was given to a series of patients (13) with renal disease, a tremendous rise in plasma magnesium occurred within 4-6 hours (from a normal 2 mg/100 cc to as high as 9-11 mg/100 cc) or to about two thirds of the concentration at which coma sets in. The authors concluded that when magnesium sulfate is administered by mouth to patients with renal disease it is absorbed continuously from the intestine. Therefore dangerously high magnesium levels are reached in the blood.

4. Stevens and Wolff (735) attempted to correlate the serum level of magnesium owing to rectal absorption with the clinical state of the subject.

Six volunteer male subjects were administered hypertonic solutions of magnesium sulfate by rectum (maximum of 300 cc). Four of the subjects received a one-half saturated solution and two a saturated solution.

From the results of their experiment as well as reports in the literature, the authors stated that it was obvious magnesium sulfate is absorbed from the gastrointestinal tract including the rectum in sufficient amounts to raise the blood level significantly (see individual curves in original paper) and to produce depressant cerebral effects. They additionally noted that even normally functioning kidneys may not remove the salt from the blood stream fast enough to prevent the accumulation of amounts toxic to the central nervous system.

Normal serum magnesium appeared to be in the range of 1.5-2.5 mg/100 cc.

#### Relation of serum magnesium level and effects noted

Serum Magnesium Level (mg/100 cc)	Nature of effect
4	sensations of warm skin and dry mouth
7	drowsiness noted
12	marked drowsiness
12-16	diaphragmatic respirations
20.1	respiratory arrest

In another experiment, three subjects were each given a saturated solution of magnesium sulfate orally (40 cc, 25 g). All showed some increase in their serum magnesium (3.6, 4.0 and 4.3 g/100 cc respectively).

#### Total amount of administered magnesium sulfate and serum magnesium levels

Magnesium sulfate administered (g)	Serum magnesium level (g/100 cc)
25 (orally)	4 (average)
94 (per rectum)	7.0
126 (per rectum)	9.0
158 (per rectum)	16.2
189 (per rectum)	20.1 (post mortem)



The authors concluded that magnesium sulfate can be absorbed from the intact bowel in large amounts and may cause dangerous depression of the central nervous system even when renal function is normal.

5. Graham et al. (271) used  $^{28}\text{Mg}$  to obtain further information on the effect of variation of dietary intake on magnesium absorption and excretion in humans.

Ten subjects (8 male and 2 female, 25-64 years old) were fed a control diet (20 mEq Mg/day) and 5 ml of  $^{28}\text{Mg}$  (as  $\text{MgCl}_2$ , specific activity 5  $\mu\text{C}/\text{mEq}$ ) taken with breakfast. Three other subjects (1 female and 2 males) were fed abnormal diets; low Mg, 1.9 mEq/day and high Mg, 47 mEq/day.

Absorption of the  $^{28}\text{Mg}$  was studied by both fecal collection and analysis of the plasma radioactivity.

The observations were:

- (a) Within an hour following ingestion, absorption began, reached a steady level at 2 to 3 hours and continued relatively constantly for the next 4-6 hours. By this time 80% of the total absorption had taken place.
- (b) After 12 hours there was very little absorption.
- (c) 44.3% of the ingested radioisotope was absorbed with the average diet (20 mEq Mg/day).
- (d) 75.8% was absorbed on the low magnesium diet (1.9 mEq/day).
- (e) 23.7% was absorbed on the high magnesium diet (47 mEq/day).

The authors concluded that:

- (a) There is fairly uniform absorption throughout the small intestine.
- (b) There is little or no absorption from the large bowel.

- (c) Absorption is affected by the amount presented to the intestinal mucosa but the relationship is not linear.
- (d) Absorption is apparently unaffected by depletion or repletion of body magnesium.

### III. Metabolism and Excretion

#### A. Rats

1. Clark (142) studied the effects of orally administered supplemental, non-toxic amounts of magnesium chloride on calcium and phosphate metabolism in adult rats. A basic diet containing the same amount of magnesium (0.05%) was fed to 144 adult male Holtzman rats (275-295 g, divided into 12 groups). The diet of each group contained different percentages of calcium and phosphorus (40% of diet) as mixtures of various salts of these elements (see Table 17 ) after 5 days of habituation to the diets, each of the groups was subdivided into two subgroups of 6 animals. A 10% (w/v) glucose solution was added to the drinking water of all the rats. One group of each of the subgroups was also given magnesium chloride in the drinking water (final concentration 2% w/v).

A balance study was carried out for 12 days. During this time the control animals consumed 50 mg of magnesium (as  $MgCl_2$ ) while the experimental animals consumed 800-1000 mg. As a result of the data developed from analysis of both the food and fluid consumption and the urine and feces the following conclusions were made:

- (a) Provided the diets contained at least 0.4% calcium, supplemental magnesium ions decreased fecal calcium significantly.
- (b) Supplemental magnesium ions significantly increased urinary calcium in all groups.

Table 17. Dietary calcium-to-phosphorus ratios (142)

		Percent Dietary Calcium				
		0.2	0.4	0.8	1.6	
Percent Dietary Phosphorus	0.8	KH <sub>2</sub> PO <sub>4</sub>	14.5	14.5	8.0	8.0
		Na <sub>2</sub> HPO <sub>4</sub>			3.7	3.7
		NaH <sub>2</sub> PO <sub>4</sub> ·H <sub>2</sub> O	7.1	7.1		
		CaH(PO <sub>4</sub> ) <sub>2</sub> ·H <sub>2</sub> O	12.6	12.6	23.5	
		Ca <sub>3</sub> (PO <sub>4</sub> ) <sub>2</sub>				28.9
		CaCO <sub>3</sub>		5.0	10.7	12.0
	0.4	K <sub>2</sub> HPO <sub>4</sub>			8.0	8.0
		KH <sub>2</sub> PO <sub>4</sub>	10.5	10.5		
		KHCO <sub>3</sub>	1.5	1.5		
		Na <sub>2</sub> HPO <sub>4</sub>			3.7	3.7
		NaH <sub>2</sub> PO <sub>4</sub> ·H <sub>2</sub> O	7.2	7.2		
		CaH(PO <sub>4</sub> ) <sub>2</sub> ·H <sub>2</sub> O			7.3	7.2
		CaCO <sub>3</sub>	5.0	10.0	17.1	37.3
	0.2	K <sub>2</sub> HPO <sub>4</sub>	5.2	5.2	6.7	6.7
		KHCO <sub>3</sub>	5.4	5.4		
		KCl			1.1	
		NaH <sub>2</sub> PO <sub>4</sub> ·H <sub>2</sub> O	3.6	3.6		
		NaHCO <sub>3</sub>	2.2	2.2		
		NaHPO <sub>4</sub>			3.7	3.7
		CaCO <sub>3</sub>	5.0	10.0	20.0	40.0

- (c) Supplemental magnesium ions affected fecal phosphorus as follows: (a) no effect when dietary phosphorus was low (0.2%), (b) caused a decrease when dietary phosphorus was 0.4%, (c) caused an increase when dietary phosphorus was high (0.8%) and calcium low (0.2%).

Magnesium ions were found to lower serum calcium. While at low dietary levels of phosphorus (0.2%) the magnesium-phosphorus interaction had the effect of lowering serum calcium.

The conclusions drawn from this study were:

- (a) Dietary magnesium ions markedly influenced the absorption and excretion of calcium and phosphorus. These effects of magnesium were also dependent on the dietary levels of calcium and phosphorus, i.e., calcium absorption was increased only when the diet contained 0.4% or more calcium.
- (b) The increase in intestinal absorption of calcium was reflected in elevated urinary calcium. The authors noted that other studies have shown that alkaline magnesium salts such as the oxide or hydroxide are as effective as the acidic chloride in producing hypercalciuria.
- (c) The improvement in phosphorus balance as a result of supplemental dietary magnesium appears to be a consequence of renal conservation of phosphorus.

Two possible consequences for human beings were pointed out by the authors:

- (a) Since divalent magnesium has also been shown to increase calcium balance and absorption in man, it might be of value in the treatment or prevention of those metabolic bone diseases which result from inadequate calcium absorption from the intestines.

- (b) A partial explanation for the reduction of renal calculus formation in man following the oral administration of magnesium ions may be the observation in this study that supplementary dietary magnesium ions resulted in a fall in urinary phosphorus. (See also this section D, 6 and Biochemical Section V, C, 1.)

2. MacIntyre et al. (464) studied the effect of parathyroid hormone on the urinary excretion of magnesium ions in the rat. Pure bovine parathyroid hormone (2000 units/mg potency) was infused (10  $\mu$ g) into the tail vein of male parathyroidectomized rats (150 g) along with an infusion solution containing: glucose, 44 g/liter; sodium chloride 19 mEq/liter; calcium lactate, 10 mEq/liter; magnesium chloride, 10 mEq/liter; and sodium heparin, 0.1 g/liter. A marked fall in magnesium excretion was observed. An effect was noted on magnesium excretion when as little as 1  $\mu$ g of the hormone was infused. The authors claimed that these experiments conclusively show that parathyroid hormone causes marked conservation of urine magnesium. They attribute this to a direct renal effect owing to a change in the tubular handling of the magnesium cation.

#### B. Dogs

1. Speranskaja-Stepanova (725) studied the influence of magnesium on renal activity in dogs. Dogs with chronic fistulas of the ureter (3, sex and age not given) were injected with crystalline  $\text{MgSO}_4$  in 20% and 50% solutions. The doses ranged between 0.08 and 0.4 g/kg BW. Most of the doses administered were a 50% solution of 0.08 g/kg BW. At this dosage no effect on urinary excretion was observed. Doses higher than 0.4 g/kg caused a distinct reduction in the quantity of urine.

2. Haury and Cantarow (304) studied the effect of magnesium sulfate on calcium, magnesium and inorganic phosphorus in the serum, and peritoneal

fluid of dogs. Of the four adult dogs used in the experiment, three were injected intramuscularly and one subcutaneously with 50 cc of a 25% magnesium sulfate solution.

The effects observed on the serum and peritoneal fluid were:

- (a) A fall in serum Ca but a relatively insignificant change in peritoneal fluid Ca concentration.
- (b) A fall in serum inorganic P and a simultaneous fall in peritoneal fluid P.
- (c) Tetany occurred in 3 dogs in which the serum Ca concentration fell below 8 mg%. It was relieved by intravenous injection of a calcium compound. The authors noted that an explanation for the hypocalcemic tetany was not apparent particularly in the presence of marked Mg concentration in both the serum and peritoneal fluids (also presumably, in tissue fluids).

3. Massry et al. (486) evaluated the characteristics of the renal handling of magnesium in dogs. A variety of test procedures were used (32 experiments) with 30 female mongrel dogs (12-27 kg) in which magnesium chloride or magnesium sulfate added to 15% dextrose in water was infused to deliver 1.0-3.0  $\mu\text{g}$  Mg/min./kg BW over 3-6 hours as follows:

- (a) Magnesium chloride or magnesium sulfate alone (8 dogs, 10 experiments).
- (b) Simultaneous infusion of 0.9% NaCl and  $\text{MgCl}_2$  (3 anesthetized dogs).
- (c) Simultaneous infusion of 0.9% NaCl,  $\text{CaCl}_2$  and  $\text{MgCl}_2$  (3 anesthetized dogs).

- (d) Simultaneous infusion of 0.9% NaCl and MgCl<sub>2</sub> in dogs pretreated with deoxycorticosterone acetate (DCA) (2 anesthetized dogs).
- (e) Infusion of MgCl<sub>2</sub> in dogs pretreated with DCA (5 anesthetized dogs).
- (f) Simultaneous infusion of CaCl<sub>2</sub> and MgCl<sub>2</sub> (5 anesthetized dogs).
- (g) Parathyroid extract administered during MgCl<sub>2</sub> infusion (4 anesthetized dogs). (See original paper for details of these experiments.)

The results of these experiments indicated the following:

- (a) Magnesium excretion is determined by filtration and reabsorption alone without evidence for tubular secretion.
- (b) There is a maximal tubular reabsorptive capacity (T<sub>m</sub>) for magnesium of approximately 140 µg/min/kg BW.
- (c) Extracellular volume expansion produced by saline infusion, calcium infusion and chronic treatment with DCA are each associated with a decrease in T<sub>m</sub> for magnesium.
- (d) Parathyroid hormone may directly enhance magnesium reabsorption.

#### C. Pigs

Miller et al. (508) determined both the effects of dietary magnesium on the utilization of calcium, phosphorus and magnesium by the baby pig and the magnesium requirement of the baby pig. When the baby pigs were 4-6 weeks of age, 29 mineral balance studies were made.

Calcium, phosphorus and magnesium balance studies were conducted with pigs receiving 75, 225, 325, 425 or 825 ppm magnesium in their diet. Dietary levels of Ca and P were 0.8 and 0.6% respectively. Distilled water was the sole source of drinking water. Data obtained from this study indicated that increasing the dietary level of magnesium did not significantly affect daily Ca or P retention but increased daily Mg retention. The balance data indicated that the dietary Mg requirement is no higher than 325 ppm.

#### D. Humans

1. Chesley and Tepper (136) studied some of the effects of magnesium acetate and sulfate on the renal excretion of electrolytes in women. The subjects of the study were 34 women, 5 nonpregnant and 29 who were pregnant with a wide range of inulin clearances. Magnesium acetate (100 ml of 10% solution; 93.2 mEq Mg) or magnesium sulfate (16.2 mEq Mg in 10% solution) were injected in various ways (see original paper for details). The results of these experiments were:

- (a) The magnesium clearance increased as a roughly linear function of the serum magnesium concentration. The acetate or sulfate anion had little or no apparent effect on magnesium excretion.
- (b) The corrected ratio of magnesium clearance (corrected for protein binding of serum magnesium) to inulin clearance approached 1 at the highest levels of magnesium studied. This ratio was roughly a linear function of the serum "filterable" magnesium.
- (c) The injection of magnesium salts was followed by an increased urinary calcium excretion. There was a direct relation between



the increased calcium excretion and the serum magnesium concentration.

- (d) As the serum magnesium concentration rose, sodium and chloride excretion was stimulated. The excretion of sodium and chloride dropped back to or below control levels when the serum magnesium became constant or fell.
- (e) The effect of magnesium on potassium excretion varied from subject to subject.

2. Aikawa et al. (1965) used  $^{28}\text{Mg}$  to explore the kinetics of magnesium distribution in normal subjects and in patients with various diseases. A total of 27 studies were performed on 9 normal subjects and 16 patients with various diseases. The  $^{28}\text{Mg}$  was administered intravenously in 12-30 mEq of stable magnesium in 250-350 ml of 5% aqueous dextrose. Urine samples and blood samples from the brachial vein were collected at intervals (most over 24 hours; a few blood samples over 37 hours and urine samples over 67 hours).

The observations in normal subjects were:

- (a) Plasma disappearance was rapid within the first several hours.
- (b) Within 24 hours, a mean of 19.8% of the injected radioactivity was accounted for in the urine.
- (c) Within 18 hours equilibration of  $^{28}\text{Mg}$  in the bile occurred but fecal excretion was negligible.
- (d) The specific activities of plasma and urine stabilized by the 18th hour and thereafter showed only a gradual decrease.
- (e) Exchangeable magnesium contents ranged between 2.6 and 5.3 mEq/kg BW (less than 16% of the estimated total magnesium content of the body).

- (f)  $^{28}\text{Mg}$  exchanged very slowly with the stable ion in bone, muscle and erythrocytes.

There were no striking differences between these results with normal patients and those obtained from patients with diabetes mellitus and hepatic diseases.

The authors summarized the knowledge concerning magnesium metabolism in human beings at the time they wrote their paper (1960) as follows:

- (a) The fraction of magnesium absorbed when it is introduced into the body by the gastrointestinal route is considerably smaller than had been thought.
- (b) Magnesium metabolism appears to be related to the metabolism of phosphorus and glucose in which the liver appears to play a role.
- (c) Whether or not there is a specific hormonal regulatory mechanism is not known.
- (d) The labile pool of magnesium in the body, a small fraction of the total body content of magnesium, is confined primarily to the soft tissues.
- (e) Magnesium exchange in bone, muscle, and erythrocytes is extremely slow.
- (f) Parenterally administered magnesium is excreted very slowly by the kidneys. The over-all balance however, is maintained by a compensatory rapid excretion of endogenous magnesium.

3. Consolazio et al. (151) studied magnesium excretion in sweat. The study was conducted on 3 healthy young men for a total of 32 days.


The men were exposed on a balance study for 16 days to a 37°C temperature for 7.5 hours. It was found that from 10.1 to 15.4 percent of the total output of magnesium was via sweat. The magnesium loss amounted to about 17 mg per day. Acclimatization did not occur as it did in the cases of sodium and potassium. The authors concluded that under extreme conditions sweat could account for about 25% of the magnesium lost daily. When intakes are low this could be important.

4. Ozsoylu (565) included a discussion of magnesium metabolism in his paper. Briefly he noted:

(a) As an intracellular cation known to be an activator in several pathways of carbohydrate metabolism, magnesium is second in abundance only to potassium.

(b) Figure shows some of the pathways in which the magnesium ion has some role such as Embden-Meyerhof, glucose storage and pentose shunt.

(c) The magnesium ion is also required in the citric acid cycle:




(d) The magnesium ion is known to be necessary for the activation of ATP requiring enzymes as well as phosphorolase, phosphatase, enolase, peptidase and decarboxylase.

(e) An absolute requirement for this ion, has however, not been proven.

5. Jones et al. (364) investigated the relationship between the functional status of the thyroid gland and magnesium metabolism. Hyperthyroid and hypothyroid patients were studied by determinations of serum and erythrocyte magnesiums, exchangeable magnesiums, and by complete

Figure 1.  
Enzymatic Systems Requiring Mg



balance studies during therapy with propylthiouracil and trifiodothyronine, respectively. Eight hyperthyroid and eight hypothyroid patients were used in this study which was carried out with  $^{28}\text{Mg}$ . At zero time 50-150  $\mu\text{c}$  of  $^{28}\text{Mg}$  was injected intravenously over a 2-3 minute period. The results of serum urine and stool analyses showed that:

- (a) Plasma magnesium values tended to be lowered in hyperthyroidism and elevated in hypothyroidism.
- (b) Erythrocyte magnesium levels were significantly reduced in only one hyperthyroid patient before therapy and varied predictably after therapy.
- (c) Hyperthyroid patients were found to have decreased plasma magnesium, increased urinary excretion of  $^{24}\text{Mg}$  and  $^{28}\text{Mg}$  and normal total and cellular exchangeable magnesium before therapy.
- (d) Hypothyroid patients had elevated plasma magnesium, decreased urinary  $^{24}\text{Mg}$  and  $^{28}\text{Mg}$  excretion, increased fecal magnesium excretion, and strikingly reduced total and cellular exchangeable magnesium before therapy.
- (e) Erythrocyte magnesium values were normal in both groups.

The authors concluded that the data suggest that a defect in magnesium transport occurs in thyroid hormone deficiency states.

6. Drake (200) followed the effects of orally administered magnesium oxide ( $\text{MgO}$ ) on urinary Ca and P excretion and attempted to ascertain whether  $\text{MgO}$  exerts a "protective effect" in calcium oxalate stone formation. Forty healthy young men on a submarine patrol were the experimental subjects. Urinary samples were taken at the beginning and end of the patrol. During the intervening period twenty-one of the men ingested

310 mg MgO daily, while nineteen served as controls. The test group was found to excrete significantly more calcium than the controls. The author postulated that magnesium oxide exerts its protective influence on renal stone formation by means of a mechanism whereby the rate of calcium excretion is altered. He also noted that this finding does not agree with the "decreased urinary excretion" theory for the protective influence, but rather lends some support to the "magnesium-oxalate balance" theory. (See also Biochemical Section V, C, 1 and this section A, 1.)

#### IV. Enzymes and Other Biochemical Parameters

##### A. In Vitro

1. MacIntyre (463) summarized the in vitro actions of magnesium ion in a review. He noted however, that so many enzymes are likely to be affected that predicting in vivo effects of magnesium deficiency from the in vitro actions of the ion is quite impossible.

The enzyme systems for which magnesium is important are:

- (a) Magnesium activates the alkaline phosphatases, pyrophosphatases and prostatic acid phosphatases.
- (b) Magnesium activates all enzymes transferring phosphate from ATP or to ADP. The author noted that the reactions involving ATP and ADP are so widespread that magnesium must influence all life processes.

Examples of the phosphate-transferring enzymes activated by magnesium are: hexokinase, fructokinase, creatine transphosphorylase, phosphopyruvic transphosphorylase, diphosphopyridine nucleotide phosphorylase and phosphoglucomutase.

- (c) Magnesium ions are necessary whenever diphosphothiamine is required. In this group are: yeast carboxylase, mammalian heart carboxylase and the pyruvic oxidase system of brain.
- (d) Enolase, which catalyses the interconversion of 2-phosphoglyceric and phosphoenol-pyruvic acids, generating an energy-rich phosphate bond in the process, requires high concentration of magnesium.
- (e) Some peptidases, such as leucine aminopeptidase, require magnesium.
- (f) Many of the reactions of glycolysis are activated by magnesium ions.

2. Wacker and Parisi (809) summarized the information on the enzyme systems critical to cellular metabolism for which magnesium has been found to be an activator in vitro. It is involved as follows:

- (a) The enzymes that hydrolyze and transfer phosphate groups:  
e.g. the phosphatases and the enzymes concerned in reactions involving adenosine triphosphate (ATP) are activated by magnesium.
- (b) It is a co-factor for oxidative phosphorylation.
- (c) The highly ordered organization of DNA, RNA and ribosomes is stabilized by the presence of magnesium.
- (d) The aggregation or disassociation of component particles in ribosomes obtained from Escherichia coli, baker's yeast, pea seedlings, rat liver and rabbit reticulocytes is critically dependent on the ambient magnesium concentration.


The authors noted in this regard that the fundamental importance of magnesium in cellular function is suggested by its analogous role in plant and animal systems and the generality of this phenomenon.

- (e) It is also required for the in vitro synthesis and degradation of DNA.
- (f) The first step in all the amino acid activating systems, in which a specific amino acyl sRNA synthetase forms a complex with its amino acid in the presence of ATP, is known to be magnesium dependent. The second or transfer step in which a specific sRNA accepts its amino acid from the enzyme complex is not ordinarily magnesium dependent but becomes so if the sRNA is denatured in a manner which removes the metals it contains naturally. The two steps in the formation of aminoacyl sRNA are:

I. Activation



II. Transfer



The authors concluded from all of the above observations that since magnesium is the most abundant divalent intracellular cation that it would be a tenable hypothesis to assume analogous in vivo function.



3. Lebduska and Cervinka (434) examined whether colloidal magnesium hydroxide causes an increase in phagocytosis in vitro. White corpuscles from horse venous blood were used. The experiments were performed in test tubes containing: 2 cc of the separated and washed white cells (largely polynuclear); 2 cc of the solutions being studied (colloidal  $\text{Mg(OH)}_2$  in four different concentrations as shown on Table 18 ); and 0.25 cc of a B. coli culture in peptone containing broth. One tube without  $\text{Mg(OH)}_2$  acted as a control. The method used was first to count 300 polynuclear cells at each concentration and then determine the number of bacteria phagocytised. Table 18 shows the results. The numbers indicate the phagocytoses per 100 leucocytes.

Colloidal magnesium hydroxide was shown to be a strong stimulator in vitro of phagocytosis.

Table 18. Magnesium Effect on Phagocytosis (434)

Experiment	Substance studied	Concentration of the solution				
		Control	N/10	N/100	N/1000	N/10,000
1	$\text{Mg(OH)}_2$	20	30	49	42	57
2	$\text{Mg(OH)}_2$	90	46	164	210	176
3	$\text{Mg(OH)}_2$	42	29	51	59	66
4	$\text{Mg(OH)}_2$	25	26	53	59	59
5	$\text{Mg(OH)}_2$	65	--	100	164	140

## B. Rats

MacIntyre et al. (464) studied the relation of parathyroid hormone to magnesium homeostasis. The details of this experiment are given in Biochem Section III, A, 2.

An important conclusion drawn by the authors was that parathyroid hormone causes marked conservation of urine magnesium and calcium. They further suggested the hypothesis that the mechanism by which plasma magnesium is maintained in the body is the variation in secretion of parathyroid hormone in response to changes of plasma magnesium.

## C. Guinea Pigs

Frommel et al. (239) studied the changes in serum cholinesterase when magnesium chloride was injected in guinea pigs. Their experiment is summarized in Table 19.

The authors pointed out that the liver is the most important organ in the regulation of serum cholinesterase. Furthermore, they noted that in producing toxicity in the guinea pig, by injecting inorganic ions, the various functions of the glandular parenchyme have been severely disturbed resulting in anatomical problems.

As can be seen from the tabulated results, magnesium chloride activates cholinesterase. The authors found this effect of interest in contributing to the understanding of the pharmacodynamics of magnesium chloride as an ionic mineral medicine.

## D. Dogs

1. LaBarre and Vesselovsky (421) continued their research on the action of hypnotics on the digestive glands by determining the influence of magnesium sulfate on the functioning of the exocrine portion of the pancreas.

Table 19. Magnesium Effect On Serum Cholinesterase (239)

Ions	Guinea pig	Weight	*Dose received in g/kg	CHE before	CHE after	Result in %	Average
Magnesium chloride	m	410	4 x 1,0	2,40	3,60	+33%	+34.5%
	m	430	5 x 1 g	2,05	2,75	+34%	
	m	570	2 x 1 g	1,40	1,90	+36%	

\*The injected dose was calculated according to Aberhalden (Handbook of Biological Work Methods 1, 7) to obtain the chronic toxicity for the ion studied.

The experiments were carried out on dogs kept constantly secreting by two means: (1) by continuous intrasaphenous injection of a diluted secretin solution or (2) by administering a test dose of secretin both before and after injecting the magnesium sulfate.

In the first series, there were four experiments in which secretin was injected in chloralosed dogs with a pancreatic fistula (previously ligatured to prevent the passage of chyme into the duodenal region). Once secretion was stabilized 100-150 mg% magnesium sulfate in 20 cc distilled water was injected intrasaphenously.

The results showed the following:

- (a) Weak doses of magnesium sulfate (100 mg/kg) resulted in a slight but brief stimulation of pancreatic secretion.
- (b) The reduction of pancreatic secretion became more marked at elevated doses.
- (c) There was very little variation of lipase content compared to the amount secreted per cc. Generally speaking, the enzyme content of the pancreatic juice varied in proportion to the quantity of juice secreted in a given time.

In a second series of experiments there were two observations:

- (a) When the secretin (10 mg/kg) was injected twice at 60 minute intervals in chloralosed dogs with a pancreatic fistula, the secretory responses in a given dog were constant.
- (b) After an intrasaphenous dose of secretin when magnesium sulfate was injected, weak doses (100 mg) did not particularly modify the pancreatic secretory response. When the magnesium

sulfate doses reached 150 mg there was a definite decrease of the pancreatic secretory response in two cases.

In this experimental series, the lipase content of the collected juice was reduced.

A third series of experiments were designed to gain understanding of the cause of the decrease in pancreatic secretory ability observed after magnesium sulfate administration. Secretin was administered intravenously to vagotomised dogs with a medullary section at the level of the third cervical vertebra. When constant pancreatic secretion was established, magnesium sulfate was administered in doses of 100-150 mg/kg. The results of four experiments indicated that when the pancreas is not under the control of the central nervous system, magnesium sulfate does not reduce pancreatic secretion.

The authors concluded from their experiments that:

- (a) Magnesium sulfate has a depressive effect on gastric and pancreatic secretions.
- (b) The inhibition of postinsulin gastric hypersecretion can be attributed to a postmagnesium paralysis of the thalamic centers.
- (c) The observed blocking of the pancreatic exocrine function apparently is dependent on the integrity of the encephalic centers.
- (d) In those persons in whom pancreatic function is cut-off by administering secretin or who have food in the digestive tract, magnesium sulfate reduces pancreatic juice production and enzyme formation. This action is similar to that of the opiate hypnotics.

2. LaBarre and Kettenmeyer (419) studied whether magnesium sulfate modifies postinsulin hyperadrenalinemia by paralysing the thalamic centers. Lightly chloralosed dogs (number, age, and sex not given) were intravenously injected with 100-150 mg/kg BW of magnesium sulfate. In order to evaluate the adrenaline secretion quantitatively, a fistula was made in the suprarenal vein. Samples of blood were taken and adrenalinemia was evaluated by studying the comparative effect of various blood samples on intestinal motility. (See Figures 1 and 4 in original paper for further details.)

The authors concluded from these experiments, that moderate doses of magnesium sulfate (100-150 mg/kg) inhibit postinsulin hyperadrenalinemia by paralysing the thalamic centers.

## V. Drug Interaction

### A. Rats

Haggard and Greenberg (292) investigated the potential of magnesium sulfate as an antidote for strychnine poisoning. In the experiment recorded in Figure 2, 23 rats (age, sex and strain not given) were injected peritoneally with strychnine in varying amounts from 0.001 to 0.004 mg/g BW. Magnesium sulfate solution was injected intraperitoneally in varying dosages (unconsciousness was caused at 0.8 mg/g BW and death at 0.9 mg/g BW and above). As can be seen in Figure 2 the antispasmodic, magnesium sulfate, did not either prevent or even diminish strychnine convulsions in rats. Magnesium sulfate is therefore not an antidote for strychnine.




Fig. 2. Magnesium sulfate administered to rats poisoned with strychnine: The rats represented by the black dots died; those represented by circles lived. The dose of strychnine and apomorphine given any one of the rats is indicated by the points on the abscissa and ordinate corresponding to the rat. Magnesium sulfate is not an antidote for strychnine. (292)

## B. Rabbits

1. Barbour and Taylor (039) studied the relations obtained between various combinations of sodium barbital and magnesium chloride with respect to toxicity and hypnotic action. Both compounds were administered subcutaneously to normal, healthy adult rabbits. Table 20 shows the toxicity of the two drugs separately and combined.

Table 21 shows the narcotic effects of combining sodium barbital with magnesium chloride as compared to those obtained with sodium barbital alone. From this table it can be seen that it took about one-third longer for the onset of either light or deep narcosis with sodium barbital alone than with a 1:2 combination of sodium barbital and magnesium chloride. Recovery as well as narcosis began much more slowly with the 1:2 combination. Partial recovery required about one-third as long and complete recovery about two-thirds as long as with sodium barbital alone. The phenomenon of accelerated onset and accelerated recovery from barbital narcosis occurred over a wide range of dosage and combination ratios. These different ratios are indicated on Figure 3 .

The authors speculated that the observation that when magnesium chloride is added to sodium barbital recovery is hastened, is probably due largely to the relative rapidity of both absorption and excretion of the smaller magnesium chloride molecule. They further suggested that the "protective antagonism" observed could be accounted for both by the difference in time of peak effect as well as by differences in the anatomical loci of toxicity.



Table 20. Toxicity For Rabbits (039)

Na BARBITAL	MgCl <sub>2</sub>	SURVIVED	DIED	FATALITY
<i>mgm. per kgm.</i>	<i>mgm. per kgm.</i>			<i>per cent</i>
350				0
400				100
				Roemer (9)
	700	11	1	9
	725	10	6	60
	750	2	3	60
	775	0	2	100
188	+	376	2	0
192	+	385		Predicted m.l.d.
196	+	392	2	0
212	+	424	4	0
220	+	440	2	50
230	+	460	1	50
240	+	480	3	25

Table 21. Na-Barbital Narcosis With and Without MgCl<sub>2</sub> (039)

NUMBER OF EXPERI- MENTS	AMOUNT INJECTED		M.L.D.	AVERAGE TIME REQUIRED TO PRODUCE			
	Barbital	MgCl <sub>2</sub>		Light narcosis	Deep narcosis	Partial recovery	Complete recovery
	<i>mgm. per kgm.</i>	<i>mgm. per kgm.</i>		<i>minutes</i>	<i>minutes</i>	<i>minutes</i>	<i>minutes</i>
12	200	0	50	51 ± 2.9	123 ± 22.1	> 167	> 600
10	300	0	75	55 ± 2.6	84 ± 3.8	> 696	> 810
12	150	300	68	41 ± 1.9	62 ± 2.0	233†	560†
7	100	400	81*	41 ± 4.5	66 ± 2.7	281	475

\* Of predicted.

† Eight rabbits only.




Fig. 3. Rabbit Narcotic and Toxic Dosage of Sodium Barbitol, Magnesium Chloride and Their Combinations (039)

Ordinates: Sodium barbitol milligrams per kilogram; abscissa: magnesium chloride mg/kg. A, threshold dosage for deep narcosis; B. toxicity line (predicted or additive); C, toxicity line based on findings with Na-barbital plus  $MgCl_2$  in 1:2 ratio; small dots, results from individual rabbits; crosses, average result from two rabbits; large dots, average results well established by numerous experiments (see tables 20 and 21). Line D represents all sub-lethal doses of a combination of one part sodium barbitol to two parts of magnesium chloride. Small figures above points indicate minutes required to produce light narcosis; figures below points, minutes required to produce deep narcosis; (infinity = no deep narcosis). Long dash line, threshold dose for complete recovery within ten hours (shaded area longer persistence of action).

The main conclusions of this study were:

- (a) A combined single injection of magnesium chloride and sodium barbital when given to rabbits can both hasten the onset of narcosis and lessen the persistence of the narcotic effect without evidence of significantly increasing toxicity.
- (b) When two parts of magnesium chloride are combined with one part sodium barbital, the minimum lethal dose is considerably higher than would be predicted on the basis of simple summation of effects.

2. Barbour and Winter (037) investigated the antipyretic action of magnesium chloride alone and combined with amidopyrine (pyramidon) in rabbits.

Seventeen healthy adult rabbits were made feverish (1.5°F increase in 2 hours) by subcutaneously injecting 1 cc/kg BW of hay-infusion. Doses of magnesium chloride in 5% solution ranging from 150-300 mg/kg BW were administered to 12 of the rabbits with 5 as controls. It was found that definite magnesium antipyresis was produced with a dose of about 200 mg/kg BW and above of the chloride. The authors noted that this was well below the minimum lethal dose of 700 mg/kg BW previously determined by them.

In another series of experiments magnesium chloride was administered in varying combinations with amidopyrine to 18 rabbits. (See original paper for details.)

The authors found that the combination of a weak amidopyrine dose (100 mg) with a weak magnesium chloride dose (150 mg) produces a profound drop in temperature. The maximum decrease in 9 rabbits was 3.9°F. This

was a greater effect than was produced by separate doses of either 200 mg/kg amidopyrine or 300 mg/kg magnesium chloride.

The authors concluded from their study that combinations of magnesium chloride with amidopyrine showed both higher therapeutic efficiency and lower toxicity than amidopyrine alone.

3. Winter and Barbour (831) studied the effect of magnesium chloride on the antipyretic action of sodium salicylate. When 150 mg/kg BW magnesium chloride (ineffective alone) was subcutaneously administered (5% solution) combined with 50 mg/kg BW sodium salicylate (5% solution) to 13 fevered rabbits and combined with 100 mg/kg BW sodium salicylate to 8 fevered rabbits, rapid reduction in temperature was noted. Control fevered rabbits (17) were used to judge the temperature difference.

It was observed that:

- (a) The combination of 50 mg/kg BW sodium salicylate with 150 mg/kg BW magnesium chloride was more effective than 100 mg/kg BW sodium salicylate administered alone.
- (b) The slow, extended temperature fall produced by salicylate was modified to a rapid but briefer effect with the addition of magnesium chloride.

The authors concluded that magnesium salts potentiate the early effects of antipyretic drugs. This synergistic effect takes place even when intense or prolonged action tends to be inhibited.

#### C. Dogs

1. Winter and Barbour (831) investigated the effect on the antipyretic action of aspirin when combined with magnesium chloride or magnesium oxide.

They found that the combination of 150 mg/kg BW magnesium chloride and 100 mg/kg BW aspirin reduced the fever of 4 dogs as effectively as 150 mg/kg BW aspirin alone.

The combination of magnesium oxide with aspirin (100 mg/kg BW aspirin plus 100 mg/kg BW MgO with 5 fevered dogs) was not as effective as an antipyretic as 150 mg/kg BW aspirin alone. A study of magnesium oxide alone showed that it had a weaker antipyretic action than magnesium chloride (see original paper for curves). The authors attributed this to slower absorption of the oxide as compared to the chloride.

2. Winter et al. (832) studied whether the antipyretic action of phenacetin in dogs would be increased by simultaneous oral administration of magnesium oxide. The dogs were made feverish by injection with fresh hay infusion.

When MgO in doses from 75-300 mg/kg BW was orally administered to 15 dogs, 6 of them showed an antipyretic effect which was not dose related. When 200 mg/kg BW phenacetin was orally administered combined with 100 mg/kg BW MgO to 11 fevered dogs, the antipyretic effect was as great as 300 mg/kg BW phenacetin given alone.

The authors concluded that the antipyretic effect of orally administered phenacetin was enhanced by the addition of MgO. They considered their results as strongly indicative of synergism since large doses of MgO alone were not antipyretic.

#### D. Humans

Gershoff and Prien (253) investigated the effects of the daily administration of magnesium oxide (MgO) and vitamin B<sub>6</sub> (pyridoxine) on patients with histories of recurring calcium oxalate urolithiasis (see also Biochemical Section III, D, 6).

Male and female adult patients with a two-year history of forming two or more calcium oxalate renal stones yearly, were used in this study. The subjects were asked to take two 100 mg MgO tablets and one 10 mg pyridoxine tablet daily. Thirty-six patients remained in the program for 5 years.

After one year urine and serum constituents showed the following effects:

- (1) The urines of 45 of 51 patients showed an increased ability to hold oxalate in solution.
- (2) Urinary calcium and citric acid were raised.
- (3) Urinary xanthurenic acid was lowered.
- (4) Urinary magnesium, oxalic acid, phosphate-P or pyrophosphate-P showed no significant changes in values.
- (5) Serum magnesium was lowered.
- (6) Serum beta-lipoproteins decreased in 25 of the 51 subjects.

After 5 years, 30 out of 36 patients maintained on the program showed either no recurrence or decreased recurrence of stone formation.

The authors postulated that a possible explanation for the protective action of MgO and vitamin B<sub>6</sub> administration in preventing renal stone formation might involve an effect on the solvent characteristics of urine. They also pointed out that since there is no reason to believe there is a common etiology in all cases of calcium oxalate urolithiasis, a variation in individual response to prophylactic treatment would be expected.

## VI. Consumer Exposure

As discussed in the Chemical Section VIII, magnesium is a required co-factor for many enzymes, and is also required by plants in the course of photosynthesis. By virtue of these facts, magnesium is found at some level in all vegetables, fruits, meats and seafoods. Table 22 from Schroeder et al. (677) gives magnesium values for a wide variety of foods on both a weight and a caloric basis. (Analyzed by atomic absorption spectrophotometry.)

In addition to exposure to magnesium as a natural constituent of fruits, vegetables, dairy products, meats, and seafood, magnesium may also be encountered as a result of the usage of various magnesium compounds in food processing. Of these compounds, magnesium carbonate is produced in the greatest quantity, with almost 500,000 pounds being produced in the United States in 1970, as shown by an NAS/NRC survey (Table 23).

Magnesium carbonate, in addition to its antacid and laxative uses, is employed in the processing of a variety of foods and may serve as an anti-caking and drying agent, a carrier, a disintegrating and dispersing agent, a color fixative, a color retention agent, and as an adjunct. Magnesium carbonate, in conjunction with sodium phosphate and an emulsifier, has been used in attempts to reduce the amount of chlorophyll degradation in green plants (038). Magnesium carbonate has reportedly been used as an adulterant in pepper improving the appearance of the pepper (206). Also, not too infrequently, it has been added to baking powders for the purpose of adding deceptive bulk.

Table 22. Magnesium in Foods (wet weight) (677)

	µg/g	mg/100 kcal
<b>Fish and sea food</b>		
Blue Gill, raw, frozen	375	31.7
Cod, raw, frozen (3)	249 (207-285)	30.4
Haddock, raw, frozen (2)	484 (436-532)	63.2
Haddock, cooked, chowder (2)	359 (328-389)	37.0
King Crab, cooked, frozen	530	41.7
Lobster, cooked, frozen	376	31.6
Oyster, raw, frozen	154	30.8
Scallops, cooked, frozen (6)	335	31.9
Smelt, raw, frozen	237	28.2
Tuna, canned	334	34.4
Mean	348	36.6
<b>Meat</b>		
Beef and fat	402	14.7
Beef Liver	203	14.2
Chicken leg	255	12.6
Chicken wing	195	9.6
Ground round steak	258	8.9
Lamb chop (2)	249 (239-258)	13.3
Lamb liver	234	16.4
Pork loin	209	7.4
Top round steak	383	15.4
Gelatin, unflavored	283	35.8
Mean	267	14.8
<b>Dairy products</b>		
Cheese, cheddar (2)	268 (266-270)	6.3
Egg, whole	113	6.9
Egg, white	103	27.8
Egg, yolk (2)	101 (100-101)	2.9
Milk, whole, homog. vit. D	102	15.5
Milk, skim, powdered	1067	32.7
Milk, human (2)	29 (28-29)	3.8
Milk, fat free	162	46.3
Milk "cereal cream" 12% butter fat plus milk solids	280	6.9
Mean, less dry milk	157	18.2
<b>Vegetables, fresh</b>		
<b>Leafy</b>		
Broccoli	321	229.3
Brussels sprouts	233	145.6
Cabbage	166	63.8
Chicory	94	104.4
Chives	241	219.1
Endive	160	145.5
Escarole	89	80.9
Lettuce, green, outer	139	126.4
Lettuce, pale, inner	85	77.3
Parsley	566	269.5
Mean, less parsley	170	146.2



Table 22. (Continued)

	µg/S	mg/100 kcal
<b>Vegetables, fresh</b>		
<b>Roots</b>		
Carrots	185 (111-296)	80.4
Leek	264	101.6
Leek greens	478	191.2
Onion (2)	89 (75-103)	38.7
Parsnips	456	93.1
Potatoes (2)	167 (155-179)	19.2
Radish	162	103.0
Scallions	137	33.1
Turnip	95	52.3
Mean	226	80.8
<b>Vegetables, fresh</b>		
<b>Legumes</b>		
Beans, green, string	297	92.8
Beans, wax, yellow	241	89.3
Beans, Lima, dry (2)	1697 (1445-1948)	49.2
Beans, kidney, dry	775	22.4
Peas, green	185	22.0
Pea, green, dry	1182	34.0
Mean, fresh	241	51.4
<b>Vegetables, fresh</b>		
<b>Fleshy</b>		
Artichoke, globe	487	334.7
Cauliflower	229	208.2
Celery	66	73.3
Cucumber	107	118.9
Egg plant	82	54.7
Mushroom	122	174.3
Pepper, green (3)	102 (92-114)	46.4
Squash, butternut	165	30.5
Squash, yellow	137	72.1
Squash, zucchini	237	139.4
Tomato (3)	177 (121-241)	125.4
Mean	174	124.4
<b>Fruits and fruit juices</b>		
Apple	58	12.9
Pear	64	16.0
Grape juice, canned	80	12.1
Orange juice	108	28.4
Papaya	0.1	< 0.1
Mean, less papaya	78	17.4
<b>Nuts</b>		
Brazil nut	3175	49.3
Walnut	1315	24.0
Hazel nut	1739	29.0
Almond	2927	43.9
Peanut	1583	26.3
Pecan	1078	18.0
Mean	1970	32.6

Table 22. (Continued)

	µg/g	mg/100 kcal
<b>Grains and cereal products</b>		
Barley	1505	41.8
Buckwheat	2526	70.2
Corn, hybrid	664	69.2
Corn, late sunshine	393	41.0
Corn, meal, yellow	269	7.6
Corn starch	22	0.6
Cream-filled doughnut	185	4.6
Kellogg's Special K	412	12.2
Kellogg's concentrate	574	15.6
Millet	1670	47.7
Egg noodles	702	19.9
Oats	1705	42.2
Puffed rice	150	4.3
Pumpernickel	550	22.0
Wheat, flour, patent	299	8.5
Wheat, gluten	18	0.5
Wheat bread	204	7.7
Whole wheat bread	340	13.8
Rice, brown	1477	40.9
Rice, wild	906	25.1
Rice, Japanese (composite 204 samples)	251	7.0
Rye, whole	1333	39.8
Tapioca, minute	37	1.0
Wheaties	1032	23.9
Mean	805	23.6
<b>Oil and fats</b>		
Butter, lightly salted (2)	24 (20-27)	0.3
Butter, unsalted	6	< 0.1
Cod liver oil	1	—
Cottonseed oil	1	< 0.1
Peanut oil	8	0.1
Pork fat	5	< 0.1
Sunflower oil	6	< 0.1
Tallow, beef (2)	5 (2-8)	< 0.1
Mean	7	< 0.1
<b>Condiments, spices, etc.</b>		
Allspice, ground	1783	71.3
Cloves, whole	2549	101.9
Chili powder	3329	97.9
Garlic, fresh	230	16.8
Iodized salt	1667	—
Mustard, ground	4225	91.3
Pepper, black, ground	2323	75.2
Thyme, ground	615	24.6
Mean	2598	71.7
<b>Beverages</b>		
Beer	100	23.8
Cocoa, powdered	4289	94.9
Coffee, 97% caffeine free, dry powder	6362 (5624-7143)	—
Coffee, infusion	48	96.0
Gin	0.3	< 0.1
Grape wine, white	98	11.5

Table 22.(Continued)

	ppb	ngg/100 kcal
Tea, black, infusion	11	110.0
Tea, black, leaves	2183	---
Tea, Japanese green, infusion	3	30.0
Tea, Japanese green, leaves	1666	---
Vermouth, Italian	135	9.9
Whisky, bourbon	1.3	< 0.1
Whisky, scotch	4.5	0.2
Sugars and syrups		
Corn syrup (3)	17 (14-23)	0.6
Honey	15	0.5
Jelly, mild cider	215	8.3
Jelly, wild cherry	32	1.2
Jelly, thyme-grape	33	1.3
Maple syrup, Vermont fancy (2)	150 (135-164)	5.1
Molasses, beet	110	4.3
Molasses, cane (4)	2093 (1311-2526)	81.4
Raw sugar, Columbian	122	3.1
Raw sugar, Phillipine	66	1.7
Sugar cane, green	190	---
Sugar crystals, cane, England	3	< 0.1
Sugar, white, crystals, U.S.	3	< 0.1
Sugar, white, France	2	< 0.1
Sugar, white, granulated, U.S.	2	< 0.1
Mean, less cane and molasses	59	2.0

Table 23. Annual Poundage Data for NAS Appendix A Substances (Groups I and II)

Substance name (Survey no.)	# Reports to NAS 1960/1970	Poundage reported to NAS (matching reports for both years)		Total 1970 poundage reported to NASA <sup>a</sup>
		1960	1970	
Magnesium carbonate NAS 0110	12/14	250,983	490,577	498,127
Magnesium hydroxide NAS 0111	---	---	---	170,116
Magnesium oxide NAS 0112	5/6	99	4,287	4,291
Magnesium phosphate (di) NAS 0113	---	4,400	12,855	12,855
Magnesium stearate NAS 0116	14/16	17,100	225,876	233,474
Magnesium sulfate NAS 0117	0/4	4,600	4,242	15,242

<sup>a</sup>There were no additional poundages reported to FEMA.

Other magnesium compounds and some of their uses include: magnesium chloride, as a color fixative, color retention agent, firming agent, and as an emulsifier for suspending dried skim milk in water (831); magnesium hydroxide, as an antacid, laxative, color fixative, color retention agent, and adjunct; magnesium oxide, as an antacid, laxative, buffer, neutralizing agent, and as a clarifier of sugar juice and reductant of molasses formation in raw sugar processing (610); magnesium phosphate (di- and tri-basic) and magnesium sulfate, as nutrients and dietary supplements. Magnesium sulfate has also been used in depression of free fatty acid formation in cold-stored fish muscle (677). Magnesium stearate is used as an anticaking and anti-drying agent, an emulsifier, a foaming and whipping agent, and as a binder, filler, or stabilizer for some plasticizers in the preparation of food-packaging materials. Magnesium stearate has also been used as a tablet lubricant, giving protection against the browning of spray-dried lactose in tablets (405). Table 24 shows usage levels, taken from an NAS/NRC survey, of some of these magnesium compounds. Table 25 also from an NAS/NRC survey, presents possible daily intake of these compounds.

Another source of exposure to magnesium is in water, where magnesium salts are partially responsible for the hardness of water. For a breakdown of magnesium concentrations in some American waters, consult Table 4 Chemical Section, VIII, Occurrence.

Table 24. Usage Levels Reported for NAS Appendix A Substances (Group I) Used in Regular Foods (R)

Substance name (Survey no.)	Food no.	Category name	# Firms reporting	- Usual use - WTD mean, %	- Maximum use - WTD mean, %
Magnesium carbonate NAS 0110	01	Baked goods (R)	-	.23393	.23394
	03	Other grain (R)	-	.00018	.00036
	07	Frozn dairy (R)	-	.01100	.01100
	10	Meat prods (R)	-	.00098	.00196
	11	Poultry (R)	-	.00100	.00200
	13	Fish prods (R)	-	.00200	.00200
	14	Procsd vegg (R)	-	---	---
	21	Soups (R)	-	.00100	.00200
	22	Snack foods (R)	-	.00100	.00200
	23	Bev type I (R)	-	.00641	.00641
	27	Gravies (R)	-	.02000	.02500
	30	Hard candy (R)	-	.00060	.00060
	48	Seas flavrs (R)	-	.90963	.96977
Magnesium hydroxide NAS 0111	05	Milk prods (R)	-	.07200	.09600
Magnesium oxide NAS 0112	06	Cheese (R)	-	.01000	.02000
	14	Procsd vegg (R)	-	.00380	---
	22	Snack foods (R)	-	.02000	.02000
	28	Imit dairv (R)	-	.00900	.00900
	30	Hard candy (R)	-	.08333	.08333
Magnesium phosphate di NAS 0113	16	Soft candy (R)	-	.03000	.03000
Magnesium stearate NAS 0116	01	Baked goods (R)	4	.00022	.00043
	16	Soft candy (R)	6	.66721	1.55070
	17	Conf frost (R)	-	.60404	.61129
	31	Chewing gum (R)	-	---	---
	48	Seas flavrs (R)	-	1.00000	1.00000

Table (Continued)

Substance name (Survey no.)	Food no.	Category name	# Firms reporting	- Usual use - WTD mean, %	- Maximum use - WTD mean, %
Magnesium sulfate	15	Condm relsh (R)	-	.00400	.00500
NAS 0117	23	Bev type I (R)	-	.01092	.01092
Malic acid	01	Baked goods (R)	7	.00997	.01193
NAS 0118 FEMA 2655	02	Break cerls (R)	4	.12082	.14355

Table 25. Possible Daily Intake of NAS Appendix A Substances (Groups I and II), per Food Category and Total Dietary, Based on Food Consumption by Total Sample -- See Explanatory Notes in Exhibits Section

Substances names (survey no.)	Food no.	Category name	# of Firms	Age	Possible daily intake, mg		
					Average	High A	High B
Magnesium carbonate NAS 0110	01	Baked goods (R)	-	0-5 mo.	7.953620	10.526850	7.953960
				6-11 mo.	59.418220	121.175740	59.420760
				12-23 mo.	127.491850	210.069140	127.497300
				2-65+ yr.	320.951960	476.749340	320.965680
Magnesium carbonate NAS 0110	03	Other grain (R)	-	0-5 mo.	.000900	.003000	.001800
				6-11 mo.	.017460	.051430	.034920
				12-23 mo.	.029520	.068220	.059040
				2-65+ yr.	.050040	.110520	.100080
Magnesium carbonate NAS 0110	07	Frozen dairy (R)	-	0-5 mo.	.110000	.451000	.110000
				6-11 mo.	1.045000	2.904000	1.045000
				12-23 mo.	1.584000	3.718000	1.584000
				2-65+ yr.	2.816000	6.787000	2.816000
Magnesium carbonate NAS 0110	10	Meat prods (R)	-	0-5 mo.	.010780	.028420	.021560
				6-11 mo.	.202860	.546840	.405720
				12-23 mo.	.295960	.508620	.591920
				2-65+ yr.	.768320	1.274980	1.536640
Magnesium carbonate NAS 0110	11	Poultry (R)	-	0-5 mo.	.005000	.023000	.010000
				6-11 mo.	.039000	.132000	.078000
				12-23 mo.	.066000	.184000	.132000
				2-65+ yr.	.129000	.328000	.258000
Magnesium carbonate NAS 0110	13	Fish prods (R)	-	0-5 mo.	.002000	.006000	.002000
				6-11 mo.	.026000	.098000	.026000
				12-23 mo.	.108000	.270000	.108000
				2-65+ yr.	.248000	.618000	.248000



Table 25. (Continued)

Substances names (survey no.)	Food no.	Category name	# of Firms	Age	Possible daily intake, mg		
					Average	High A	High B
Magnesium carbonate NAS 0110	14	Procsd vegg (R)	-	0-5 mo.	---	---	---
				6-11 mo.	---	---	---
				12-23 mo.	---	---	---
				2-65+ yr.	---	---	---
Magnesium carbonate NAS 0110	21	Soups (R)	-	0-5 mo.	.002000	.015000	.004000
				6-11 mo.	.233000	.727000	.466000
				12-23 mo.	.348000	.961000	.696000
				2-65+ yr.	.317000	.845000	.634000
Magnesium carbonate NAS 0110	22	Snack foods (R)	-	0-5 mo.	---	.001000	---
				6-11 mo.	.004000	.011000	.008000
				12-23 mo.	.011000	.031000	.022000
				2-65+ yr.	.013000	.037000	.026000
Magnesium carbonate NAS 0110	23	Bev type I (R)	-	0-5 mo.	.153840	.230760	.153040
				6-11 mo.	1.455070	4.980570	1.455070
				12-23 mo.	3.474220	10.416250	3.474220
				2-65+ yr.	6.666400	17.800570	6.666400
Magnesium carbonate NAS 0110	27	Gravies (R)	-	0-5 mo.	.020000	.060000	.025000
				6-11 mo.	.280000	.780000	.350000
				12-23 mo.	.720000	2.040000	.900000
				2-65+ yr.	1.660000	4.260000	2.075000
Magnesium carbonate NAS 0110	30	Hard candy (R)	-	0-5 mo.	.000000	.000000	.000000
				6-11 mo.	.000600	.001800	.000600
				12-23 mo.	.001800	.005400	.001800
				2-65+ yr.	.003600	.010200	.003600

Table 25 (Continued)

Substances names (survey no.)	Food no.	Category name	# of Firms	Age	Possible daily intake, mg		
					Average	High A	High B
Magnesium carbonate NAS 0110	48	Seas flavrs (R)	-	0-5 mo.	---	---	---
				6-11 mo.	---	.090963	---
				12-23 mo.	---	.181926	---
				2-65+ yr.	.090963	.454815	.096977
Magnesium carbonate NAS 0110		All categories	13	0-5 mo.	8.258140	11.345090	8.282160
				6-11 mo.	62.721210	131.499393	63.290070
				12-23 mo.	134.130350	228.453556	135.066280
				2-65+ yr.	333.714283	509.275425	335.426377
Magnesium hydroxide NAS 0111	05	Milk prods (R)	-	0-5 mo.	3.888000	2.880000	5.184000
				6-11 mo.	44.928000	216.072000	59.904000
				12-23 mo.	39.240000	125.568000	52.320000
				2-65+ yr.	28.440000	86.832000	37.920000
Magnesium hydroxide NAS 0111		All categories	-	0-5 mo.	3.888000	2.880000	5.184000
				6-11 mo.	44.928000	216.072000	59.904000
				12-23 mo.	39.240000	125.568000	52.320000
				2-65+ yr.	28.440000	86.832000	37.920000
Magnesium oxide NAS 0112	06	Cheese (R)	-	0-5 mo.	---	.010000	---
				6-11 mo.	.270000	.970000	.540000
				12-23 mo.	.780000	2.220000	1.560000
				2-65+ yr.	.940000	2.360000	1.880000
Magnesium oxide NAS 0112	14	Procsd vgs (R)	-	0-5 mo.	.053200	.159600	---
				6-11 mo.	.912000	2.128000	---
				12-23 mo.	1.482000	2.481400	---
				2-65+ yr.	3.230000	5.441600	---

Table 25 (Continued)

Substances names (survey no.)	Food no.	Category name	# of Firms	Age	Possible daily intake, mg		
					Average	High A	High B
Magnesium oxide NAS 0112	22	Snack foods (R)	-	0-5 mo.	---	.020000	---
				6-11 mo.	.080000	.220000	.080000
				12-23 mo.	.220000	.620000	.220000
				2-65+ yr.	.260000	.740000	.260000
Magnesium oxide NAS 0112	28	Imit dairy (R)	-	0-5 mo.	.000000	.000000	.000000
				6-11 mo.	.126000	.207000	.126000
				12-23 mo.	.072000	.306000	.072000
				2-65+ yr.	.081000	.135000	.081000
Magnesium oxide NAS 0112	30	Hard candy (R)	-	0-5 mo.	.000000	.000000	.000000
				6-11 mo.	.083330	.249990	.083330
				12-23 mo.	.249990	.749970	.249990
				2-65+ yr.	.499980	1.416610	.499980
Magnesium oxide NAS 0112	83	Formulas (B)	-	0-5 mo.	30.145860	55.227000	30.145860
				6-11 mo.	6.142320	29.265820	6.142320
				12-23 mo.	1.975600	.556760	1.975600
Magnesium oxide NAS 0112		All categories	7	0-5 mo.	30.199060	55.416600	30.145860
				6-11 mo.	7.613650	33.040810	6.971650
				12-23 mo.	4.779590	6.934130	4.077590
				2-65+ yr.	5.010980	10.093210	2.720980
Magnesium phosphate di NAS 0113	16	Soft candy (R)	-	0-5 mo.	.060000	.600000	.060000
				6-11 mo.	.660000	2.040000	.660000
				12-23 mo.	1.050000	2.790000	1.050000
				2-65+ yr.	1.740000	5.280000	1.740000

Table 25. (Continued)

Substances names (survey no.)	Food no.	Category name	# of Firms	Age	Possible daily intake, mg		
					Average	High A	High B
Magnesium phosphate di NAS 0113	83	Formulas (B)	-	0-5 mo.	1106.903610	2027.839500	1106.903610
				6-11 mo.	225.535320	1074.590070	225.535320
				12-23 mo.	72.540600	20.443260	72.540600
Magnesium phosphate di NAS 0113		All categories	-	0-5 mo.	1106.963610	2028.439500	1106.963610
				6-11 mo.	226.195320	1076.630070	226.195320
				12-23 mo.	73.590600	23.233260	73.590600
				2-65+ yr.	1.740000	5.280000	1.740000
Magnesium silicate NAS 0115	01	Baked goods (R)	-	0-5 mo.	.029920	.039600	.116620
				6-11 mo.	.223520	.455840	.871220
				12-23 mo.	.479600	.790240	1.869350
				2-65+ yr.	1.207360	1.793440	4.705960
Magnesium silicate NAS 0115	03	Other grain (R)	-	0-5 mo.	.012500	.042500	.015000
				6-11 mo.	.242500	.715000	.291000
				12-23 mo.	.410000	.947500	.492000
				2-65+ yr.	.695000	1.535000	.834000
Magnesium silicate NAS 0115	16	Soft candy (R)	-	0-5 mo.	.001400	.014000	.001400
				6-11 mo.	.015400	.047600	.015400
				12-23 mo.	.024500	.065100	.024500
				2-65+ yr.	.040600	.123200	.040600
Magnesium silicate NAS 0115	31	Chewing gum (R)	-	0-5 mo.	---	---	---
				6-11 mo.	---	---	---
				12-23 mo.	---	---	---
				2-65+ yr.	---	---	---

Table 2, (Continued)

Substances names (survey no.)	Food no.	Category name	# of Firms	Age	Possible daily intake, mg		
					Average	High A	High B
Magnesium silicate NAS 0115		All categories	5	0-5 mo.	.043820	.096100	.133020
				6-11 mo.	.481420	1.218440	1.177620
				12-23 mo.	.914100	1.802840	2.385850
				2-65+ yr.	1.942960	3.451640	5.580560
Magnesium stearate NAS 0116	01	Baked goods (R)	4	0-5 mo.	.007480	.009900	.014620
				6-11 mo.	.055880	.113960	.109220
				12-23 mo.	.119900	.197560	.234350
				2-65+ yr.	.301840	.448360	.589960
Magnesium stearate NAS 0116	16	Soft candy (R)	6	0-5 mo.	1.334420	13.344200	3.101400
				6-11 mo.	14.678620	45.370280	34.115400
				12-23 mo.	23.352350	62.050530	54.274500
				2-65+ yr.	38.698180	117.428960	89.940600
Magnesium stearate NAS 0116	17	Conf frost (R)	-	0-5 mo.	---	.604040	---
				6-11 mo.	.604040	1.208080	.611290
				12-23 mo.	1.208080	4.228280	1.222580
				2-65+ yr.	1.812120	4.832320	1.833870
Magnesium stearate NAS 0116	31	Chewing gum (R)	-	0-5 mo.	---	---	---
				6-11 mo.	---	---	---
				12-23 mo.	---	---	---
				2-65+ yr.	---	---	---
Magnesium stearate NAS 0116	48	Seas flavrs (R)	-	0-5 mo.	---	---	---
				6-11 mo.	---	.100000	---
				12-23 mo.	---	.200000	---
				2-65+ yr.	.100000	.500000	.100000

Table 25. (Continued)

Substances names (survey no.)	Food no.	Category name	# of Firms	Age	Possible daily intake, mg		
					Average	High A	High B
Magnesium stearate NAS 0116		All categories	14	0-5 mo.	1.341900	13.958140	3.116020
				6-11 mo.	15.338540	46.792320	34.835910
				12-23 mo.	24.680330	66.676370	55.731430
				2-65+ yr.	40.912140	123.209640	92.464430
Magnesium sulfate NAS 0117	15	Condm relsh (R)	-	0-5 mo.	---	.004000	---
				6-11 mo.	.032000	.088000	.040000
				12-23 mo.	.112000	.304000	.140000
				2-65+ yr.	.352000	.848000	.440000
Magnesium sulfate NAS 0117	23	Bev type I (R)	-	0-5 mo.	.262080	.393120	.262080
				6-11 mo.	2.478840	8.484840	2.478840
				12-23 mo.	5.918640	17.745000	5.918640
				2-65+ yr.	11.356800	30.324840	11.356800
Magnesium sulfate NAS 0117		All categories	-	0-5 mo.	.262080	.397120	.262080
				6-11 mo.	2.510840	8.572840	2.518840
				12-23 mo.	6.030640	18.049000	6.058640
				2-65+ yr.	11.708800	31.172840	11.796800

MAGNESIUM SALTS BIBLIOGRAPHY

- 1. Abramowitz, E.W., and J.J. Russo. 1940  
Fixed eruption from magnesium hydroxide:  
poly-sensitivity  
Arch. Dermatol. Syph. 41:707-710
2. Aiezzati, L., and R. Prando. 1967  
Humoral regulation of lipolysis and liposynthesis  
(Ital.)  
Metabolismo 3(2):192-201
3. Adlersberg, D. 1929  
Über das Wesen der sogenannten  
gallenblasenreflexe. (So-called gall bladder  
reflex)  
Am. J. Physiol. 90(2):259
- 4. Aikawa, J.K., and P.D. Bruns. 1960  
Placental transfer and fetal tissue uptake of  
Mg<sup>28</sup> in the rabbit  
Proc. Soc. Exp. Biol. Med. 105:95-98
- 5. Aikawa, J.K., G.S. Gordon, and E.L. Rhoades. 1960  
Magnesium metabolism in human beings: Studies  
with Mg<sup>28</sup>  
J. Appl. Physiol. 15:503-507
6. Aisawa, Mituo. 1939  
The effect of the anterior pituitary on tissue  
respiration, especially the influence of various  
pharmaceutical agents on this effect. I. The  
influence of sodium luminal, strychnine nitrate,  
magnesium sulfate and chloral hydrate  
Folia Endocrinol. Japon. 15:10-11
- 7. Alcock, N., and I. MacIntyre. 1962  
Inter-relation of calcium and magnesium absorption  
Clin. Sci. 22:185-193
8. Ali, M. 1970  
Degranulation of beta-cells of the islets of  
Langerhans following intragastric administration  
of magnesium chloride  
Indian J. Physiol. Pharmacol. 14:11-14
9. Alieva, S.G. 1967  
Kliniko-fiziologicheskoe izuchenie narushenii  
perifericheskogo krovoobrashcheniya  
i effektivnosti kompleksnoi terapii pozdnikh  
toksikozov beremennykh. (Clinical  
and physiological study of disorders in the  
peripheral blood circulation and the  
effectiveness of complex therapy of late toxemia  
during pregnancy)  
Akush. Ginekol. 43(11):27-33
10. Allcroft, W.M. 1947  
The fate of calcium and magnesium in the blood  
stream of the ruminant  
Vet. J. 103(5):157-170
11. Allen, Shannon C., and Victor E. Hall. 1952  
Response of rat temperature and respiratory rate  
to thermal stress, magnesium chloride and  
tetrazolium salts  
Fed. Proc. 11(1):4
- 12. Allison, Ira S. 1930  
The problem of saline drinking waters  
Science 71(1848):559-560
13. Amano, T., K. Kato, and R. Shimizu. 1952  
Studies on the role of plakin  
Med. J. Osaka Univ. 3(2/3):293-311
14. Amorosi, Osvaldo. 1930  
I sali alogeni di calcio e di magnesio nei  
processi di cicatrizzazione delle  
ferite. Ricerche sperimentali. (Ca and Mg  
halogen salts in cicatrization processes of  
wounds)  
Riv. Patol. Sper. 5(4/5):327-342
15. Anderson, N.G. 1953  
Studies on isolated cell components. V. The  
effects of various solutions on the  
nuclear envelope of the isolated rat liver  
nucleus  
Exp. Cell Res. 4(2):306-315
16. Anderson, Norman G., and Earl M. Wilbur. 1952  
Studies on isolated cell components. IV. The  
effect of various solutions on the isolated rat  
liver nucleus  
J. Gen. Physiol. 35(5):781-796
17. Anderson, W.W. 1921  
Magnesium sulphate poisoning in children  
J. Med. Assoc. Ga. 10:826
18. Andrews, B.F., D.R. Campbell, and P. Thomas. 1965  
Effects of hypertonic magnesium sulfate enemas on  
newborn and young lambs  
Lancet 1965-II(7402):64-65
19. Annenkov, B.N. 1965  
Vliyaniye sul'fata magniya na udalenie strontsiya-  
90 iz organizme nekotorykh laboratornykh  
zhivotnykh. (Effect of magnesium sulfate on  
strontium-90 excretion from the body of some  
laboratory animals)  
Radiobiologiya 5(4):620-621
20. Anon. 1943  
El sulfato de magnesio en obstetricia.  
(Magnesium sulfate in obstetrics; review)  
Sem. Med. 2:1284-1287
21. Anon. 1964  
Food additives. Deforming agents  
Fed. Regist. 29:12515
22. Anon. 1967  
Frozen desserts. Ice cream, identity standard;  
order listing neutral and mineral salts as  
optional ingredients  
Fed. Regist. 32:12750-12751
23. Anon. 1971  
Ice cream identity standard; calcium carbonate  
and magnesium carbonate as optional ingredients  
Fed. Regist. 36(8, Jan. 13):422-423
24. Arditti, R.R. 1962  
Effetto riattivante del MnCl<sub>2</sub> su una mutazione  
indotta da azoto iprite. (Renewed effect of  
magnesium chloride on a mutation induced by  
nitrogen mustard in *Penicillium chrysogenum*)  
Atti. Ass. Genet. Ital. 7:80
25. Arents, Johannes. 1965  
Meat preservative  
Neth. Pat. 299,216 issued Aug. 10, 1965
26. Askev, E.W. 1971  
Metabolism of fatty acids by mammary glands of  
cows fed normal, restricted roughage,  
or magnesium oxide supplemented rations  
J. Dairy Sci. 54:854-862
27. Bachrach, E. 1946  
Facteurs chimiques biothermiques. (Chemical  
biothermic factors)  
Arch. Int. Physiol. 54(1):19-29
28. Bachrach, Eudoxie, and Noel Guillot. 1941  
Influence of ionic conditions upon thermal optima  
for physiological functions  
Compt. Rend. 212:929-932
29. Bachromejew, Iw.R., and Lydia Pawlova. 1935  
Über das physikalisch-chemische blutbild bei der  
narkose  
Naunyn-Schwiedberg's Arch. Exp. Path. Pharmacol.  
178:390-396
30. Bacivarov, Crum, Gheorghe Velescu, and Constantin  
Apreotesei. 1969  
Cynara scolymus extract  
Rom. Pat. 51,592 issued Mar. 21, 1969
31. Bajpai, P.K., and K.I. Brown. 1964  
The effect of potassium and magnesium chlorides  
on the metabolic activity and fertility of  
turkey spermatozoa  
Poult. Sci. 43(2):459-461



- 32 Baldaev, S.N., and S.A. Kirillov. 1964  
Effect of magnesium sulfate on biochemical indexes in the blood of hogs during fattening (Russ.)  
Mikroelem. Sib. , Inf. Byul. 1964(3):65-67
- 33 Baldaev, S.N., and S.N. Tyumentsev. 1966  
Magnesium deficiency in calves (Russ.)  
Tr. Buryat. S-Kh. Inst. 18:298-300
- 34 Ballantyne, E.E. 1957  
Drinking waters toxic for livestock  
Can. J. Comp. Med. Vet. Sci. 21(7):254-257
- 35 Ballow, F.H., Jr. 1951  
Limestone in the California beet-sugar industry  
Calif. J. Mines Geol. 47:9-16
- 36 Barac, G., and F.G. Sulman. 1963  
Influence of alkaline earth ions on blood flow in dog kidneys  
Compt. Rend. Soc. Biol. 157(12):2351-2354
- \* 37 Barbour, H.G., and J.E. Winter. 1928  
Antipyretic action of magnesium chloride alone and combined with amidopyrine  
Proc. Soc. Exp. Biol. Med. 25:582-586
- \* 38 Barbour, H.G., and J.E. Winter. 1931  
Magnesium absorption in dogs and its effect upon the metabolism of calcium  
J. Pharm. Exp. Ther. 43(4):607-620
- \* 39 Barbour, H.G., and W.P. Taylor. 1931  
The influence of magnesium chloride upon the narcotic and toxic effects of sodium barbital  
J. Pharmacol. Exp. Ther. 42(3):321-331
- 40 Barlow, O.W., and M.S. Biskind. 1928  
Influence of daily oral administration of magnesium sulphate or lactose on blood volumes of normal, beriberi and fasting pigeons  
Am. J. Physiol. 86:594-598  
  
Barlow, O.W., and R.W. Whitehead. No date  
in vergleich des korpergewichts, der erythrocytenzahl und des gesamten blutvolumens bei gesunden ratten, bei an beriberi kranken und hungernden ratten. Der einfluss von milchzucker, mineralol und magnesiumcarbonat auf die anemie bei der reiskrankheit  
Am. J. Physiol. 89:548-554
- 42 Barnes, Richard, Helmut Vogel, and Irving Gordon. 1969  
Temperature of compensation: significance for virus inactivation  
Proc. Natl. Acad. Sci. U.S. 62(1):263-270
- 43 Baroni, V., and J. Comsia. 1932  
Chloride, effect on development of experimental sarcoma  
Cluj. Med. 13:531-533
- 44 Barreras, Robert F. 1970  
Acid secretion after calcium carbonate in patients with duodenal ulcer  
New Engl. J. Med. 282(25):1402-1405
- 45 Bassin, Alexander L., and Lester R. Whitaker. 1930  
Pharmacodynamic effects upon the gall bladder  
New Engl. J. Med. 202(7):311-318
- 46 Basu, A.K., and A.W. Woodruff. 1966  
Effect of acetazolamide and magnesium therapy on erythrocyte survival in sickle-cell anemia and sickle-cell-hemoglobin disease  
Trans. Roy. Soc. Trop. Med. Hyg. 60(1):64-69
- 47 Batko, B. 1966  
Shock after intraduodenal administration of magnesium sulphate  
Wiad. Lek. 19:1637-1640
- \* 48 Batsakis, J.G., F. Madera-Orsini, D. Stiles, and R.O. Briere. 1964  
Fluorometric estimation of magnesium in serum and urine  
Am. J. Clin. Path. 42(5):541-545
- 49 Baumann, J. 1931  
Effects  
Prog. Med. pp. 105-110
- 50 Baysal, Firuz. 1966  
Effects of some pharmacological agents on the isolated guinea pig gallbladder (Turk. )  
Ankara Univ. Tip Fak. Mecm. 19(1):97-106
- \* 51 Bazikyan, K.A., and A.A. Akimov. 1968  
Anticarcinogenic action of magnesium  
Vopr. Onkol. 14(7):57-61
- 52 Bazikyan, K.A., and A.A. Akimov. 1970  
Effect of magnesium on the toxicity and antineoplastic activity of cyclophosphane (Russ.)  
Zh. Eksp. Klin. Med. 10(1):40-44
- 53 Behchuk, L.S. 1940  
Influence of various analgesics on contraction behavior of the isolated uterus. II. Plant narcotics, antipyrine and magnesium sulfate  
Farmakol. Toksikol. 3(3):17-23
- 54 Becka, J. 1929  
As regulator of calcium metabolism  
Z. Ges. Exp. Med. 67:253-259
- 55 Becka, J. 1932  
Theory of action of antiserum; maximum effect from combination with polysan (magnesium hydroxide preparation)  
Cas. Lek. Cesk. 71:161-162
- 56 Becka, Jan. 1933  
Veränderungen der pH-zahl des blutes und des mineralstoffwechsels während der narkose. II. Mitt. Veränderungen der pH-zahl des blutes  
Mauvyn-Schmiedebergs Arch. Exp. Pathol. Pharmacol. 171:244-249
- 57 Becka, Jan. 1933  
Veränderungen der pH-zahl des blutes und des mineralstoffwechsels während der narkose. I. Mitt. Der kalk- und phosphorspiegel im blute der kaninchen während der narkose  
Mauvyn-Schmiedebergs Arch. Exp. Pathol. Pharmacol. 170:377-383
- 58 Becka, Jan. 1934  
Veränderungen der pH-zahl und des mineralstoffwechsels während der narkose. III. Der einfluss der alkalose und azidose auf den verlauf der narkose  
Arch. Exp. Pathol. Pharmacol. 174(1/2):173-181
- 59 Becka, Jan. 1935  
Studie über die bedingungen der assimilation der mineralionen im tierischen organismus. II. Einfluss der magnesiumverbindungen auf die phosphorausscheidung durch die niere und durch den darm  
Sb. Cesk. Akad. Zemed. 10:530-537
- 60 Becka, Jan. 1935  
Studie über die bedingungen der assimilation der mineralionen im tierischen organismus. I. Einfluss der magnesiumverbindungen auf die calciumausscheidung durch die niere und durch den darm  
Sb. Cesk. Akad. Zemed. 10:368-377
- 61 Becka, Jan, and Frantisek Hoch. 1932  
Konservierung von nahrungsmitteln mit magnesiumhydroxyd  
D.B. Pat. 549,017 issued Apr. 22, 1932
- \* 62 Beckman, Harry. 1925  
Der angebliche synergismus von magnesiumsulfat und morphin  
J. Am. Med. Assoc. 85:332-336

- 63 Beidler, Lloyd M. 1953  
Properties of chemoreceptors of tongue of rat  
*J. Neurophysiol.* 16(6):595-607
- 64 Belits, P.A., and M.I. Rudnyev. 1965  
Vmist sul'fidryl'nykh hrup i bilkiv u syvorottsi krovi vahitnykh py piznikh toksykozakh ta yikh zminy v protsesi likuvannya. (Content of sulfhydryl groups and proteins in the blood serum of pregnant women in the presence of late toxicoses and their changes in the process of treatment)  
*Pediatr. Akush. Ginekol.* 27(4):42-43
- 65 Benbough, J.E. 1969  
Factors affecting the toxicity of oxygen towards airborne coliform bacteria  
*J. Gen. Microbiol.* 56(Pt. 2):241-250
- 66 Bennett, T. Izod. 1923  
Die beeinflussung der magensaftabsonderung durch arzneistoffe  
*Arch. Exp. Pathol. Pharmacol.* 96:158-179
- 67 Benson, J.D. 1972  
Metabolism of fatty acids by adipose tissue and liver of cows fed normal, restricted roughage or MJO supplemented rations  
*J. Dairy Sci.* 55:83-92
- 68 Bergamini, L. 1943  
The action of salts on the activity of mucinolytic enzymes  
*Arch. Sci. Biol. (Italy)* 28:293-310
- 69 Berger, Eva. 1929  
Unterschiedliche wirkungen gleicher ionen in und auf verschiedene tierarten  
*Pflügers Arch. Physiol.* 223:1-39
- 70 Bergh, G.A., and J.A. Layne. 1942  
Effect upon sphincter of Oddi of man  
*Am. J. Dig. Dis.* 9:162-165
- 71 Bergman, A. 1929  
Action on intestine; radiologic study  
*Rev. Especialid.* 4:1125-1144
- 72 Bergman, A. 1932  
Action of sodium chloride and magnesium sulphate on experimental polyuria (modified Volhard test)  
*Prensa Med. Argent.* 19:132-143
- 73 Bergman, Adolfo. 1929  
Sulfato de magnesio  
*Rev. Especialidades* 4(6):47-66
- 74 Berka, J. 1931  
Role in metabolism of calcium  
*Rev. Med. Fr. C* 12:227-232
- 75 Berliner, M.D. 1971  
Studies on protoplast induction in the yeast phase of *Histoplasma capsulatum* by magnesium sulfate and 2-deoxy-D-glucose  
*Mycologia* 63:1164-1172
- \* 76 Bernstein, M., and S. Simkins. 1939  
Effects of intravenous injections on human heart  
*J. Lab. Clin. Med.* 25:131-141
- 77 Bhattacharyya, N.K., and D.N. Mullick. 1963  
Effect of magnesium sulfate injection in experimental hypercholesteremia in rabbits. I. Serum and tissue cholesterol  
*Ann. Biochem. Exp. Med. (Calcutta)* 23(12):515-522
- 78 Bhattacharyya, N.K., Brahm Dutt, and D.N. Mullick. 1967  
Effect of magnesium sulfate injection on histopathological changes in aorta and coronary artery in experimental hypercholesteremia in rabbits  
*Indian J. Exp. Biol.* 5(4):248-250
- 79 Bianchi, Camillo, and Ernesta Marazzi-Uberti. 1969  
Acquisition and retention of a conditioned avoidance response in mice as influenced by pemoline, by some of its derivatives and by some CNS stimulants  
*Psychopharmacologia* 15(1):9-18
- 80 Bianchi, Giovanni. 1960  
Chemical and physical constitution of curdled milk  
*Mondo Latte* 14:30-33
- 81 Bieter, Raymond M. 1935  
Die wirkung von diuretica, die in eine niere des aglomerularen seeteufels injiziert sind  
*J. Pharmacol. Exp. Ther.* 53:347-349
- 82 Billheimer, E.C., and F.W. Nitardy. 1936  
Das haltbarmachen von magnesiamilch durch citronensaure  
*J. Am. Pharm. Assoc.* 25:36-38
- 83 Binet, L., J. Dry, and R. Molinard. 1960  
Effects of choline and magnesium chloride on compensatory hepatic hypertrophy following excision of two-thirds of the liver in white rats  
*Compt. Rend. Acad. Sci. (Paris)* 250:782-785
- 84 Biro, N.A., and M. Balint. 1966  
Studies on proteins and protein complexes of muscle by means of proteolysis. I. Influence of ionic milieu on the proteolysis of myosin  
*Acta Biochim. Biophys.* 1(1):13-28
- 85 Blumenthal, Ferd. 1921  
Chlormagnesium im Speiswasser  
*Z. Angew.* 34:534
- 86 Bogicevic, J., D. Jokanovic, and P. Stuiar. 1961  
Fatal cases of magnesium sulfate poisoning in the treatment of taeniasis  
*Srp. Arh. Celok. Lek.* 89:1413-1423
- 87 Bohdanecky, Z., and J. Necina. 1963  
Course of some pharmacological tests in functionally decorticated animals  
*Physiol. Bohemoslov.* 12:55-61
- 88 Bohdanecky, Z., and T. Weiss. 1964  
The electrophysiological manifestations of narcotic state evoked by magnesium-sulfate  
*Acta Nerv. Super.* 6(1):42
- \* 89 Bohuon, C. 1962  
Microanalysis of magnesium in various biological mediums  
*Clin. Chem. Acta* 7:811-817
- 90 Bolaffi, Ada. 1930  
Il trattamento col magnesio dell'adenocarcinoma del topo con speciale considerazione del contenuto in calcio, magnesio e potassio del tumore. (Treatment of adenocarcinoma of the mouse with magnesium, with special reference to the Ca, Mg, and K content of the tumor)  
*Tumori* 16(3):201-211
- 91 Bowen, William J. 1957  
Adenosinetriphosphate and the shortening of muscular models  
*J. Cell. Comp. Physiol.* 49(Suppl. 1):274-290
- \* 92 Boyd, L.J., and D. Scherf. 1943  
Magnesium sulfate in paroxysmal tachycardia  
*Am. J. Med. Sci.* 206:43-48
- 93 Boyden, E.A., and C.L. Birch. 1928  
Entleerung der menschlichen gallenblase nach salinischen abfuhrmitteln  
*Proc. Soc. Exp. Biol. Med.* 25:840-842
- 94 Boyden, Edward A., George S. Bergh, and John A. Layne. 1943  
An analysis of the reaction of the human gall bladder and sphincter of Oddi to magnesium sulfate  
*Surgery* 13:723-733
- 95 Boyland, E., E.R. Busby, C.E. Dukes, P.L. Grover, and D. Manson. 1964  
Further experiments on implantation of materials into the urinary bladder of mice  
*Brit. J. Cancer* 18(3):575-581

- 96 Bozler, Emil, and Daniel Lavine. 1958  
Permeability of smooth muscle  
Am. J. Physiol. 195(1):45-49
- Bradway, Winifred. 1936  
The experimental alteration of the rate of metamorphosis in the tunicate, *Clavelina huntsmani* (Van Name)  
J. Exp. Zool. 72(2):213-224
- \* 98 Brady, J.P., and H.C. Williams. 1967  
Magnesium intoxication in a premature infant  
Pediatrics 40:100-103
- 99 Brannon, C.H. 1934  
Observations on the blow-fly *Lucilia sericata*, Meig  
J. Parasitol. 20(3):190-194
- 100 Brewing Industries Research Institute. 1958  
Stabilized malt beverages  
U.S. Pat. 2,857,279 issued Oct. 21, 1958
- 101 Bringe, A.N. 1969  
Studies on prevention of milk fat depression in the ruminant  
Diss. Abstr. Int. Sect. B. Sci. Eng. 29(12, Pt. 1):4455-4456
- 102 Brinley, Floyd John. 1928  
Der einfluss von chemikalien auf die viscositat des protoplasmas von anochen, angezeigt durch die Brownsche bewegung  
Protoplasma 4:177-182
- 103 Brookfield, Ronald Winston. 1934  
The effects of injection of certain salts on the calcium, magnesium and inorganic phosphorus of the serum of the rabbit  
Biochem. J. 28(2):725-733
- 104 Brovkin, D.P. 1934  
Effect on diuresis  
Sovet. Vrach. Gaz. pp. 1511-1513
- \* 105 Bruck, Samuel, and Joseph M. Fruchter. 1942  
Influence of a single dose of commonly used laxatives on gastro-intestinal motility: a comparative study  
Radiology 38:145-153
- 106 Drugsch, Theodor, and Hans Horsters. No date  
Cholagoga und cholagogie. I. Mitt. II. Die resorptionsgrosse der gallenblase. III. Über den einfluss peroral zugeführter salzlosungen auf die gallenblase; nach experimentellen untersuchungen  
Arch. Exp. Path. Pharmacol. 118:267-291, 292-304, 305-312
- 107 Buckner, G. Davis, and Amanda H. Harms. 1935  
The influence of calcium carbonate in the feed of laying hens, upon digestion  
Poult. Sci. 14(6):360-375
- 108 Budayjan, P.E., and W.P. Pawlow. 1929  
Einfluss der nebensalze des kochsalzes auf die butter  
Z. Unters. Lebensm. 58:600-608
- \* 109 Burcar, P., A.J. Boyle, and R.E. Mosher. 1964  
Spectrophotometric determination of magnesium in blood serum using wagon  
Clin. Chem. 10(11):1028-1038
- 110 Burge, W.E., and A.M. Estes. No date  
Eine untersuchung über die anorganischen bestandteile des körpers in hinsicht auf den reizeffekt, den sie auf den zuckerstoffwechsel ausuben  
Am. J. Physiol. 85:103-105
- 111 Bus, Willem C., Johannes Muetgeert, and Pieter Nienstra. 1958  
Fractionating starch into branched and linear chain components  
U.S. Pat. 2,829,990 issued Apr. 8, 1958
- 112 Busacca, Attilio. 1923  
Experimentelle untersuchungen über die von arsenobenzolen bewirkten "nitritoiden krisen." III.-V. Rkk. in vitro: verss. mit neosalvarsan und einigen löslichen natriumsalzen  
Arch. Farmacol. Sper. 36:129-136, 156-160, 166-171, 186-190
- \* 113 Byron, F.E. 1939  
Fatal results following the administration of magnesium sulfate  
J. Malaya Br., Brit. Med. Assoc. 3:100-101
- 114 Calderone, Frank A. 1935  
Studien über atherdosierung nach praanasthetischer medikation von narkotica (barbiturate, magnesiumsulfat und morphin)  
J. Pharmacol. Exp. Ther. 55:24-39
- 115 Caldini, Paolo, and Baruch Bromberger-Barnea. 1958  
Effects of magnesium and tetraethylammonium chloride on the hypothermic heart  
Science 128(3336):1435
- 116 Callison, W.E., J. Lander, and F.P. Underhill. 1930  
The potassium and calcium content of the brain under MgSO<sub>4</sub> anesthesia  
J. Pharmacol. Exp. Ther. 38(4):385-388
- 117 Camacho, J.B. Jimenez. 1942  
Obstetrics in highlands of Peru; case of eclampsia after labor, treated with magnesium sulfate and notes on native practices  
Actual. Med. Peru. 7:318-320
- 118 Carlu, L. 1934  
Ionization of magnesium chloride in therapy of keloids and keloid cicatrices  
Ann. Dermatol. Syph. 5:162-169
- 119 Carnot, P., and H. Gaehlinger. No date  
Wirkung von MgCl<sub>2</sub> und MgCl<sub>2</sub>-haltigen mineralwassern auf die anhangsdruesen des darmes  
Compt. Rend. Soc. Biol. 88:1129-1131
- 120 Carubelli, R., W.O. Smith, and J.F. Hammarsten. 1959  
Determination of magnesium and calcium in urine  
Clin. Chem. 5(1):45-49
- 121 Casey, John A. 1969  
Reduction and control of calcium salts  
Sugar J. 31(11):20-24
- 122 Caso, R., and E.R.J. Ferloni. 1943  
Intravenous or intramuscular injections of magnesium sulfate in therapy of pruritus during pregnancy, with report of cases  
Bol. Inst. Matern. 12:148-153
- 123 Caso, Rogelio, and E.R.J. Ferloni. 1943  
Intravenous magnesium sulfate during pregnancy  
Dia Med. 15:1252-1253
- 124 Cavston, F.G. 1931  
Lesser known uses  
Practitioner 126:466-469
- 125 Centre d'Expansion de Dietetique Animale "CEDIA". 1968  
Minerals for animal feed  
Fr. Addn. 91,217 issued May 3, 1968
- 126 Centre D'Expansion de Dietetique Animale "Cedia" 1967  
Cattle fodder supplement  
Fr. Pat. 1,476,548 issued Apr. 14, 1967
- 127 Cesari, G., and P.A. Bertacchini. 1966  
Learning and memory in the operatively conditioned rat after administration of Mg penoline  
Boll. Soc. Ital. Biol. Sper. 42(22):1594-1596

- 128 Chabrol, E., and M. Maximin. 1928  
Inhibitory action on bile secretion in liver  
Bull. Mem. Soc. Med. Hop. Paris 52:1693-1698
- 129 Chabrol, Etienne, and Maurice Maximin. 1929  
Recherches experimentales sur les cholagogues  
administrees par voie veineuse  
Presse Med. 37(41):666-669
- 130 Chandler, Asa C., and R.N. Chopra. No date  
Die Wirkungen der Zucker-, Magnesiumsulfat-,  
Natriumcitrat- und verdünnter saure Zufuhr auf  
die Leberschädigungen, hervorgerufen durch  
Tetrachlorkohlenstoff  
Indian J. Med. Res. 14:219-226
- 131 Chandra, Satya V., and Satya P. Srivastava. 1970  
Experimental production of early brain lesions in  
rats by parenteral administration of manganese  
chloride  
Acta Pharmacol. Toxicol. 28(1):177-183
- 132 Chang, Su-Yin, Pe-Yung Mao, and Pin Hsu. 1963  
Antitumor drugs. XVI. The effect of combination  
therapy of 5b-71 with some metal salts on  
Ehrlich ascites carcinoma in mice  
Yao Hsueh Hsueh Pao 10(10):594-599
- 133 Chao, Iping. 1939  
Action of the chlorides of the alkali earth  
metals on muscular contraction  
Chin. J. Physiol. 14:449-455
- 134 Chemische Fabrik Kalk G.m.b.H. 1968  
Feed additive containing sodium, calcium, and  
magnesium phosphates  
Fr. Pat. 1,546,505 issued Nov. 22, 1968
- 135 Chen, S.P., C.H. Kau, S.U. Mu, and K.K. Li. 1959  
Suggestions on the inspection and prevention of  
occupational diseases caused by magnesium  
carbonates  
Shandong Yikao 18:25
- \* 136 Chesley, Leon C., and Ira Tepper. 1958  
Some effects of magnesium loading upon renal  
excretion of magnesium and certain other  
electrolytes  
J. Clin. Invest. 37(10):1362-1372
- 137 Chinosolfabrik Akt.-Ges. 1930  
Darstellung von Darms desinfektionsmitteln  
D. R. Pat. 513,127 issued Nov. 22, 1930
- \* 138 Chiray, M., I. Justin-Besancon, and C. Debray.  
1948  
Action of waters containing magnesium chloride on  
intestine in vivo; experimental study  
Bull. Acad. Natl. Med. 132:76-78
- 139 Cho, M.H., and L.B. Jaques. 1956  
Heparinase. III. Preparation and properties of  
the enzyme  
Can. J. Biochem. Physiol. 34(4):799-813
- 140 Christensen, P. Agerholm, and Maria de Waal. 1947  
Magnesium sulfate and carbolic soap as antidotes  
in snake bite  
S. Afr. Med. J. 21:680-681
- 141 Chura, Alojz. 1927  
Magnesium a detsky vegetativny system nervovy.  
(Magnesium and its influence on the autonomic  
nervous system in children)  
Bratisl. Lek. Listy 7(3):121-135
- \* 142 Clark, I. 1968  
Effects of magnesium ions on calcium and  
phosphorus metabolism  
Am. J. Physiol. 214(2):348-356
- 143 Clark, Irwin, and Raymond Geoffroy. 1958  
Studies in calcium metabolism  
J. Biol. Chem. 233(1):203-205
- 144 Clissuinis, N., and A. Paradelis. 1964  
Comparative experimental research on laxatives  
(Fr.)  
Therapie 19(6):1581-1586
- 145 Codrescu, Ion. 1968  
Increasing the magnesium and calcium content of  
drinking waters  
Rom. Pat. 51,333 issued Dec. 14, 1968
- \* 146 Collins, E.N., and P.W. Russell. 1949  
Fatal magnesium poisoning following magnesium  
sulfate, glycerin, and water enema in primary  
megacolon  
Cleve. Clin. Q. 16:162-166
- \* 147 Colomb, D. 1969  
Burns followed by erythematous-pigmented lesions  
due to harsh resuscitation with Epsom salt  
Bull. Soc. Fr. Derm. Syph. 76:133
- \* 148 Committee on Specifications. 1972  
Food chemicals codex  
Comm. Food Prot., Natl. Res. Council, Natl. Acad.  
Sci. Wash., D.C. pp. 469-478; 481-484
- 149 Condorelli, L. 1968  
Effect of magnesium ions on the bathmotropism of  
the ventricular myocardium (Ital.)  
Boll. Soc. Ital. Cardiol. 13(4):381-418
- \* 150 Connerty, H.V., and A.R. Briggs. 1968  
Spectrophotometric determination of serum  
magnesium by eriochrome black T  
Am. J. Clin. Path. 50(4):499-504
- \* 151 Consolazio, C.F., L.O. Matoush, R.A. Nelson, B.S.  
Harding, and J.E. Canham. 1963  
Excretion of sodium, potassium, magnesium, and  
iron in human sweat and the relation of each to  
balance and requirements  
J. Nutr. 79:407-415
- 152 Contreras, Marin Ramos. 1936  
Die Behandlung von chorea sydenhami durch  
intramuskuläre Magnesiumsulfatinjektionen  
Presse Med. 44:228-229
- 153 Cook, James R., John W. Pollard, and Jack C.  
Cooley. 1967  
Myocardial contractile force in the treatment of  
ouabain intoxication with disodium EDTA,  
potassium chloride, and magnesium sulfate  
J. Lab. Clin. Med. 69(2):292-303
- 154 Cooke, W.R. 1942  
Important advances (intravenous magnesium sulfate)  
J. Mo. N. A. 39:303-306
- 155 Cortis-Jones, B. 1970  
Sugar colour improvement  
U.S. Pat. 3,522,092 issued in 1970
- \* 156 Craig, P., B. Zak, L.T. Iseri, A.J. Boyle, and  
G.B. Myers. 1951  
The spectrophotometric determination of magnesium  
in human serum  
Am. J. Clin. Path. 21:394-398
- 157 Crema, A. 1965  
Pharmacology of the extrahepatic biliary tract  
Arch. Ital. Sci. Pharmacol. 15(1-2):5-27
- 158 Crisler, George. 1928  
The effect of withdrawal of drinking water on the  
susceptibility of rats to certain drugs  
Am. J. Physiol. 86(3):552-555
- 159 Cristol, P., P. Passouant, C. Benezech, and  
Jeanne Dutarte. 1945  
Specificity of cholinesterase. Cholinesterase  
activity and phosphatase activity  
Compt. Rend. Soc. Biol. 139:312-313
- 160 Critelli, G., C. Cernigliaro, G. Bellisario, A.  
Santucci, G. Calcagnini, S. Rizzo, and P.  
Splendore. 1968  
Use of magnesium in bathmotropic disturbances of  
the ventricular myocardium induced by acute  
coronary occlusion (Ital.)  
Gazz. Int. Med. Chir. 73(24 Pt. 2):4609-4622

- 161 Crockett, Franklin Smith. 1927  
Further observations on the effect of magnesium sulphate on phenolsulphonophthalein elimination  
J. Urol. 17(2):139-145
- 162 Crodel, W., and H. Kahler. 1931  
Friedrichshall mineral water in place of duodenal irrigation with magnesium sulphate solution  
Ther. Gegenw. 72:53-57
- 163 Czepa, A. 1929  
Purgative effect of Epsom salts; roentgenologic observations  
Wien. Klin. Wchnschr. 42:537-538
- 164 D'yakov, A.M. 1965  
Changes in the content of high-energy phosphates in animal tissues as a result of combined administration of alpha-tocopherol and magnesium compounds (Russ.)  
Tr. Ryazan. Med. Inst. 26:25-29
- 165 Dabney, J.M., J.B. Scott, and C.C. Chou. 1967  
Effects of cations on ileal compliance and blood flow  
Am. J. Physiol. 212(4):835-839
- 166 Daniel, J., and F. Hogler. 1923  
Über veränderungen im quellungszustande der blutkolloide nach peroraler zufuhr von lösungen verschiedener elektrolyte und anelektrolyte  
Wien. Arch. Inn. Med. 6:355-372
- 167 Danilov, A.N. 1941  
Combined therapy of toxic diphtheria with antiserum and magnesium sulfate  
Sov. Med. 5(17-18):25
- 168 Dastidar, S.K. Ghosh. 1941  
Magnesium sulfate  
Indian Med. Gaz. 78:73-79
- 169 Davies, D.T.P., Aneurin Hughes, and R.S. Tonks. 1968  
Experimental and clinical lung and heart lesions resulting from intravascular platelet clumping and some factors in their prevention  
Proc. Int. Pharmacol. Meet., 3rd 1966(6):75-84
- 170 Davies, J.G. 1969  
Some aspects of raw sugar clarification. (A review)  
Sugar Technol. Rev. 1(1):77-83
- \* 171 Dawson, J.B., and F.W. Heaton. 1961  
The determination of magnesium in biological materials by atomic absorption spectrophotometry  
Biochem. J. 80:99-106
- 172 De la Llane, J. Lopez. 1942  
Use of halogen salts in biology  
Cron. Med., Valencia 36:236-243
- 173 De Almeida, M. Ozorio. 1933  
Action on contraction of posterior extremities of spinal rabbit  
Compt. Rend. Soc. Biol. 114:834-837
- 174 De Man, Th. J., B. Sjoelless, and J. Grasbuis. 1946  
Addition of minerals to the feed of grazing milch cows  
Meded. Inst. Mod. Veevoeding "De Schothorst," Hoogland Amersfoort 1946:9 pp.
- 175 De Meyts, P., A. Cession-Fossion, and G. Cession. 1966  
Effect of magnesium and calcium ions on adrenaline-induced arterial hypertension in the rat (Fr.)  
Arch. Int. Physiol. Biochim. 74(5):733-744
- 176 De Micheli, Alfredo, Gustavo Medrano, Mario Testelli, and Demetrio Sodi-Pallares. 1961  
Modifications of the electrocardiogram of a dog heart system by certain chemicals  
Arch. Inst. Cardiol. Mex. 31:751-760
- 177 De Moor, H., and H. Hendrickx. 1971  
Effect of MgO and Ca3(PO4)2 on the flow and hygroscopic properties and bulk density of spray-dried skim-milk and whole milk  
Meded. Van Fac. Landbouwwet. Rijksuniv. Gent 36(2):731-742
- 178 De Ropp, Robert S. 1967  
Metabolism of alpha- and gamma-aminobutyric acid by rat brain  
J. Neurochem. 14(6):693-694
- 179 De Siqueira, A. 1932  
With magnesium sulphate  
Rev. Assoc. Paulista Med. 1:138-141
- 180 Delbert, Pierre, and Palios. 1928  
Action des sels halogenes de magnesium sur l'evolution des cancers greffes. (Action of halogen salts of magnesium upon the growth of cancer grafts)  
Bull. Assoc. Fr. Etud. Cancer (Paris) 17(8):525-536
- 181 Delbet, P. 1929  
Effects, biological action of halogen salts of magnesium  
Med. Press 128:47
- 182 Delbet, P., and C. Palios. 1928  
Effect on development of epitheliomata grafted on mice  
Bull. Assoc. Fr. Etud. Cancer 17:315-323
- 183 Delbet, P., and Franicevic. 1932  
Effect of halogen salts on urinary hydrogen ion concentration  
Bull. Acad. Med., Paris 108:1470-1475
- 184 Delbet, P., and Palios. 1928  
Effect on mice on deficient diet  
Compt. Rend. Soc. Biol. 99:390-392
- 185 Delbet, P., and Palios. 1928  
Chloride, action on guinea-pigs on scurvy-producing diet  
Compt. Rend. Soc. Biol. 98:1534-1535
- 186 Delbet, Pierre, and A. Beauvy. 1931  
Du magnesium et du calcium dans la bile  
Bull. Acad. Med. (Paris) 105(24):987-995
- 187 Demas-Marsalet, P., Lafon, and Faure. 1942  
Intravenous administration of magnesium sulfate in therapy  
J. Med. Bord. 119:209-214
- 188 Dent, W.E., H.B. Howell, F.W. Adams, and J.P. Mehlig. 1956  
Growth performance and blood and liver copper values in Hereford calves offered certain mineral elements free choice  
J. Anim. Sci. 15(4):1103-1111
- 189 Despres, M. 1931  
Chloride, "curarizing" action  
Compt. Rend. Soc. Biol. 107:602-604
- 190 Deutsche, M. 1921  
Effect of magnesium sulphate in various intoxications  
Dtsch. Med. Wchnschr. 47:103
- 191 Deysach, L.J., and T.W. Bay. 1949  
Absorption of manganese by the thyroid gland of the guinea pig  
Proc. Soc. Exp. Biol. Med. 171(2):188-189
- 192 Dickens, Frank, and Guy Drummond Greville. 1935  
Der stoffwechsel normalen und krebsgewebes. XIII. Neutralsalzwirkungen  
Biochem. J. 29:1468-1483
- 193 Dienes, L., and John T. Sharp. 1956  
The role of high electrolyte concentration in the production and growth of L forms of bacteria  
J. Bact. 71(2):208-213

- 194 Dion, C. 1960  
Comparative study of chloralhydrate and hypertonic NaSO<sub>4</sub> in the alcohol withdrawal syndrome  
Laval. Med. 39:222-231
- 195 DiMaccio, G. 1929  
On blood pressure; antagonism of calcium; experiments  
Riv. Pat. Sper. 4:97-108
- 196 Doctor, V.M., Tad L. Patton, and Jorge Awapata. 1957  
Incorporation of serine-3-C<sup>14</sup> and formaldehyde-C<sup>14</sup> into methionine in vitro. I. Role of folic acid  
Arch. Biochem. Biophys. 67(2):404-409
- 197 Dodel, P., and G. Distogues. 1943  
The influence of some crystalloids on the toxicity of saponins on the larvae of toads  
Arch. Phys. Biol. 16(5):1-3
- 198 Dowden, Bobby F., and Harry J. Bennett. 1965  
Toxicity of selected chemicals to certain animals  
J. Water Pollut. Control Fed. 37(9):1309-1316
- 199 Drake, Clayton T. 1970  
Effects of oral magnesium oxide on renal calcium excretion  
Mil. Med. 135(2):122-132
- 200 Drake, Clayton T. 1970  
The effects of oral administration of magnesium oxide on renal calcium excretion in crew members during a submarine patrol  
U.S. Nav. Submar. Med. Cent. Rep. 601:1-5
- 201 Drews, Joergen, and Lutz Wagner. 1970  
Effect of prednisolone injected in vivo on RNA-polymerase activities in isolated rat thymus nuclei  
Eur. J. Biochem. 13(2):231-237
- 202 Dreyer, N.B. 1930  
Die Wirkung der salinischen abfuhrmittel auf den dunndarm  
Arch. Int. Pharmacodyn. Ther. 16:435-448
- 203 Dryanovska-Noninska, L., I. Gaganov, and B. Gotov. 1966  
Quantitative determination of bismuth, calcium, and magnesium ions and sodium bicarbonate in pharmaceuticals  
Farmatsiya (Sofia) 16(6):36-39
- 204 Dufour, H. 1932  
Magnesium chloride and pyroformine  
J. Med. Paris 52:257-259
- 205 Duliere, W., and L. De Bontgraef. 1924  
Irritability of isolated frog's heart under influence of magnesium chloride  
Compt. Rend. Soc. Biol. 98:843-845
- 206 Dunn, J.T., and H.C.L. Bloxam. 1937  
An unusual adulterant for pepper  
Analyst 62:121
- 207 Dybing, Fred. 1957  
The effect of adrenaline on neuromuscular transmission during partial block produced by magnesium  
Acta Pharmacol. Toxicol. 13(1):60-85
- 208 Dzhamusova, T.A. 1969  
Rate of contracture development and loss of excitability during the action of various concentrations of potassium, sodium, calcium, and magnesium chlorides on the muscle (Puss.)  
Sov. Rab. Inst. Tsitol., Akad. Nauk S.S.S.R. 1969(13):223-231
- 209 Ehrenberg, Rudolf, and Hans-Joachim Guttes. 1949  
Über die Wirkung von rhodaniden und sulfaten auf die Schwellenwerte des geschmacks  
Pflüger's Arch. Ges. Physiol. 251(6/7):664-671
- 210 Fittenmiller, R.R., J.R. Vakil, and K.M. Shahani. 1970  
Production and properties of Penicillium roqueforti lipase  
J. Food Sci. 35(2):130-133
- 211 El-Mangouri, H.A., and H.M. Nour-El-Dine. 1960  
Analytical methods for the evaluation of the iron laxative mixture of the Egyptian National Formulary 1958  
Egypt. Pharm. Bull. 42:319-324
- 212 Elmslie, W.P., and H. Steenbock. 1929  
Calcium and magnesium relations in animal  
J. Biol. Chem. 82:611-632
- 213 Emelin, Yu.A. 1968  
Ob effektivnosti antatsidov (po dannym radiotelemetrii). (Effectiveness of antacids (according to radiotelemetric data))  
Sov. Med. 31(2):21-25
- 214 Endo, Chuzaburo. 1936  
The effect of various chemicals on oxygen consumption by rabbit kidney  
Sei-i-kai Med. J. 55(7):1435-1451
- 215 Endo, Chuzaburo. 1937  
The oxygen consumption of experimental nephritic kidney and the effect of diuretics on it (Ger.)  
Sei-i-kai Med. J. 56(3):329-358
- 216 Engelhardt, Werner. 1934  
Über die resorption des magnesiumsulfates bei der mischnarkose mit avertin-magnesiumsulfat-narkophin nach martin  
Dtsch. Z. Chir. 243(4/5):344-349
- 217 Englander, Martin. 1923  
Zur wirkungsstärke der neutralisalz: Na-, K-, Ca und Mg-Chlorid auf oxydative vorgange  
Wien. Klin. wchschr. 36:845-846
- 218 Erdmann, W.D., and G. Wortmann. 1964  
The additive or superadditive activity of neuromuscular paralyzing compounds with various loci of action (Ger.)  
Arzneimittel-Forschung 14(11):1179-1182
- 219 Erickson, Y. 1955  
Simplified methods for determination of calcium and magnesium in the saliva  
J. Dental Res. 34(1):104-112
- 220 Etzelber, J.G., G.W. Clayton, Jr., A.H. Tuttle, and C.T. Emery. 1952  
Renal function studies in pediatrics; influence of magnesium sulfate on renal hemodynamics in normal children  
Am. Med. Assoc. Am. J. Dis. Child. 83:301-305
- 221 Faquet, Michel. 1969  
Effect of various mineral salts on the growth of Escherichia coli cultures subjected to the action of streptomycin or other antibiotics (French)  
Compt. Rend. Acad. Sci., Ser. D 269(12):1136-1139
- 222 Fejer, A. 1980  
Therapeutic effects of small doses of heavy metal-salts  
Ouv. Med. 84(10):101-103
- 223 Feldman, Felix, and Herbert C. Lichtman. 1967  
Delta-aminolevulinat synthetase activity in the human reticulocyte  
Biochim. Phys. Acta 141(3):653-655
- 224 Fernandez, P.C. 1972  
Metabolic acidosis reversed by the combination of magnesium hydroxide and a cation-exchange resin  
New Engl. J. Med. 286:23-24
- 225 Ferrara, Manfredi. 1921  
Über die pharmacologische wirkung des magnesiumsulfats und über seine anwendung bei der strychninvergiftung  
Arch. Pharmacol. Sper. 32:91-96

- 226 Pilla, Milan, and George E. Kenny. 1967  
Effect of magnesium on replication of rhinovirus  
HGP  
J. Virol. 1(3):439-493
- 227 Fink, Hermann, and Felix Just. 1942  
Über den vitamin B-gehalt verschiedener hefen und  
seine beeinflussung. V. Die vitamin B<sub>1</sub>-verluste  
bei der zucht von hefen im luftungsverfahren.  
(The vitamin B<sub>1</sub> content of various yeasts and  
factors which modify it. V. The losses of vitamin  
B<sub>1</sub> during culture of the yeast by the aeration  
method)  
Biochem. Z. 311(1/3):61-72
- 228 Finney, Karl F., John P. McCammon, and William G.  
Schrenk. 1949  
Effect of varying concentrations of certain  
metals and their salts on gas production  
and loaf volume  
Cereal Chem. 26:140-148
- 229 Pirina, K. 1941  
The influence of some inhibiting fluids and  
special anticoagulant preparations on blood  
coagulation  
Arch. Exp. Path. Pharmacol. 198:292
- \* 230 Fleischman, D.L. 1969  
Stability of chlorophyll and derivatives in  
processed green vegetables  
Diss. Abstr. Int. Sect. B. Sci. Eng.  
30(2):696-697
- 231 Flentje, Martin E. No date  
Calcium- und magnesiumhydrate als flockungsmittel  
J. Am. Water Works Assoc. 17:253-260
- 232 Fodere. 1930  
Contra-indication of use in prevention of cancer  
Bull. Acad. Med., Paris 104:734
- 233 Fontes, J., Freitas S., and P. Da Cunha. 1932  
Influence on contractions of uterus of guinea-pig  
previously treated with ergotin (ergot  
preparation)  
Compt. Rend. Soc. Biol. 109:501-502
- 234 Food Machinery Corp. 1933  
Verfahren zum einkochen von gewuse  
A. Pat. 1,908,795 issued May 16, 1933
- 235 Fordtran, John S. 1968  
Acid rebound  
New Engl. J. Med. 279(17):900-905
- 236 Franke, Hans. 1934  
Magnesium und kohlehydratstoffwechsel  
Naunyn-Schmiedeberg's Arch. Exp. Pathol. Pharmacol.  
174:727-741
- \* 237 Frewin, D.B. 1971  
The effect of magnesium sulphate on the blood  
vessels of the upper limb in man  
Austral. J. Exp. Biol. Med. Sci. 49:101-109
- 238 Fricker, A. 1959  
Ripening of milk  
Int. Dairy Congr., Proc. 15th Congr., Lond.  
3:1276-1283
- \* 239 Frommel, E., A.D. Herschberg, and J. Piquet. 1944  
The effects of inorganic ions on the activity of  
serum cholinesterase. II. In vivo studies with  
the guinea pig  
Helv. Physiol. Pharmacol. Acta 2:193-201
- 240 Frommel, Ed., A.D. Herschberg, and J. Piquet.  
1944  
Effects of inorganic ions on activity of serum  
cholinesterase. I. In horse serum in vitro (Fr.)  
Helv. Physiol. Pharmacol. Acta 2:169-191
- 241 Fuhner, Hermann. 1926  
Über die wirkung des bariuschlorids am  
skelettmuskel  
Arch. Exp. Pathol. Pharmacol. 119:56-65
- 242 Gaebelein, Klaus. 1970  
Salt composition for food  
Ger. Offen. 1,812,452 issued Jun. 18, 1970
- 243 Gajatto, Sante. 1938  
Action of magnesium chloride on the diuresis  
produced by intravenous injection of hypertonic  
solutions of sodium sulfate  
Arch. Pharmacol. Sper. Sci. Aff. 66(1):19-40
- 244 Gantt, W. Horsley, and G. V. Volborth. 1926  
The influence of magnesium sulphate on the  
expulsion of bile from the gall bladder  
J. Lab. Clin. Med. 11(6):542-547
- 245 Gantt, W.H. 1930  
Influence on biliary secretion  
Am. J. Med. Sci. 179:380-385
- 246 Gantt, W.H., and G.V. Volborth. 1930  
Influence on gastric, intestinal and pancreatic  
secretions  
Am. J. Med. Sci. 179:375-380
- 247 Gaponenkow, T.K., and W.N. Myrikowa. 1936  
Der einfluss mineralischer stoffe auf den  
gelatinierungsprozess des systems: pektin-zucker-  
weinsäure  
Colloid J. 2:47-50
- 248 Garcia, V. Canicio. 1944  
Biliary ileus following ingestion of concentrated  
solution; case  
Rev. Clin. Espan. 13:254-255
- 249 Garibdzhanyan, B.T., and R.A. Zakharyan. 1968  
Effect of magnesium ions on the toxicity and  
antiblastic activity of thio-TEPA (tris(1-  
aziridinyl)phosphine sulfide) (Russ.)  
Dokl. Akad. Nauk Arm. SSR 46(1):30-33
- 250 Garman, Philip. 1956  
Experiments in pest control, 1955  
Proc. 65th Ann. Meeting Conn. Entomol. Soc.  
1955:38-42
- 251 Gellhorn, Ernst. 1926  
Action of cations on smooth musculature  
Pflüger's Arch. Ges. Physiol. 213(5/6):789-799
- 252 Gerken, H.J., Jr., and J.P. Fontenot. 1967  
Availability and utilization of magnesium from  
dolomitic limestone and magnesium oxide in steers  
J. Anim. Sci. 26(6):1404-1408
- \* 253 Gershtoff, Stanley N., and Edwin L. Prien. 1967  
Effect of daily MgO and vitamin B<sub>6</sub> administration  
to patients with recurring calcium oxalate  
kidney stones  
Am. J. Clin. Nutr. 20(5):393-399
- 254 Ghosh, B.P., and N. Sarkar. 1959  
Effect of some inorganic salts on water  
absorption by rice during cooking  
Ann. Biochem. Exp. Med. 19:83-86
- 255 Gigon, Alfred. 1937  
Zur blutzuckerfrage  
Schweiz. Med. Wschr. 67:434-436
- 256 Gilberg, Yngvar. 1953  
The keeping quality of light salted herring cured  
without vinegar  
Fiskefidir. Skr. 2(12):3-23
- 257 Gilbert, Albert J. 1938  
Microscopic observations of pulmonary artery  
reactions  
J. Pharmacol. 62:228-235
- \* 258 Giamblet, E.G., A.F. Marney, and R.W. Bonsnes.  
1967  
Determination of calcium and magnesium in serum,  
urine, diet, and stool by atomic absorption  
spectrophotometry  
Clin. Chem. 13(3):204-214

- 259 Giordano, G., M. Marujo, N. Scopinaro, and C.E. Caraceni. 1968  
Urinary excretion of mono- and bivalent cations following the intravenous administration of magnesium  
Arch. "E. Maragliano" Patol. Clin. 24(2):157-174
- 260 Gittleman, H.J., S. Kukolj, and L.G. Welt. 1968  
Inhibition of parathyroid gland activity by hypermagnesemia  
Am. J. Physiol. 215(2):483-485
- \* 261 Gochman, N., and H. Giveller. 1970  
Automated, simultaneous microdetermination of calcium and magnesium by atomic absorption  
Clin. Chem. 16(3):229-234
- 262 Goldstein, Burton J., and Johnathan Braunstein. 1968  
Clinical evaluation of penolide and magnesium hydroxide as a mild stimulant in geriatric patients  
Curr. Ther. Res. Clin. Exp. 10(9):457-460
- 263 Gorhunova-Nikolaeva, M.M. 1933  
K analizu sekretornoj raboty privratnika. (Analysis of pyloric secretion) (Russ.)  
Fiziol. Zh. SSSR 16(1):199-204
- 264 Gordon, A. 1931  
Intraspinal injections of magnesium sulphate for relief of intolerable pain  
Med. J. Sec. 133:399-400
- 265 Gordon, E.M. 1960  
Minerals and spot tests  
Offic. Proc. Annu. Meet., Am. Assoc. Feed Microsc. 8:33-34
- 266 Gorn, L.F. 1963  
Toxic substances in industry and adrenal cortex function (Russ.)  
Gig. Tr. 11:44-51
- 267 Gorodetskii, F.E. 1937  
Intensifying the insulin reaction by the use of magnesium sulfate  
Sov. Vrach. Zh. 41:1485-1490
- 268 Goroq, P., and L. Szporny. 1966  
Effect of magnesium asparaginate on glucose and fat metabolism  
Arzneim. Forsch. 16(11):1486-1488
- 269 Gotsev, T., and L. Atanassova. 1956  
Circulatory effects of isotonic and hypertonic solutions; effect of magnesium sulfate on blood vessels of lower extremities of man  
Arch. Exp. Path. Pharmacol. 227:400-408
- 270 Grabowski, Tadeusz. 1965  
The effect of magnesium sulfate on the radiological behavior of the duodenal bulb  
Gastroenterologia 104(5):261-271
- \* 271 Graham, L.A., J.J. Caesar, and A.S.V. Burgen. 1960  
Gastrointestinal absorption and excretion of Mg<sup>28</sup> in man  
Metabolism 9:646-659
- 272 Gray, I., and W.J. Jordan. 1955  
Magnesium protection in mechanical trauma  
Arch. Biochem. Biophys. 57(2):521-522
- 273 Greeff, K., and B. Knippers. 1964  
Investigation of potassium-magnesium aspartate in animal experiments. Compatibility, elimination, and influence on toxicity of cardiac glycosides  
Arzneimittel-Forschung 14(10):1128-1134
- 274 Green, Arda Alden. 1931  
Studies in the physical chemistry of the proteins. VIII. The solubility of hemoglobin in concentrated salt solutions. A study of the salting out of proteins. IX. The effect of electrolytes on the solubility of hemoglobin in solutions of varying hydrogen ion activity with a note on the comparable behavior of casein  
J. Biol. Chem. 93(2):495-516
- 275 Greenberg, D.M., and M.A. Mackey. 1932  
The determination of magnesium in blood with 8-hydroxyquinoline  
J. Biol. Chem. 96:419-429
- 276 Grignani, Eugenio. 1937  
Zusammensetzung von traubensaftkonzentraten sizilianischer herkunft  
Ann. Chim. Appl. 27:209-212
- 277 Grinda, Jesus. 1950  
Effect of strongly sulfated waters on the toxicity of intestinal-muscle fibers  
Arch. Med. Exp. 11:63-78
- 278 Grishenko, V.I. 1963  
Izmeneniya PKG ploda pri lechenii pozdnikh toksikozov beremennykh fenotiazinovymi preparatami i sernokislou magneziei. (Changes in the phonocardiogram of the fetus in the treatment of late toxicoses of pregnancy with phenothiazine drugs and magnesium sulfate)  
Akush. Ginekol. 6:43-48
- 279 Griswold, R.L., and M. Pace. 1956  
Microdetermination of calcium and magnesium in tissue ash  
Anal. Chem. 28(6):1035-1037
- 280 Groh, Jindrich, Petr Zdarsky, Jaroslav Sistek, and Eva Kvasnickova. 1967  
Magnesium distribution after infusions of normal solutions of magnesium chloride hydrochloric acid, and magnesium chloride sodium bicarbonate (in anuric rats)  
Sb. Ved. Lek. Fak. Karlovy. Univ. Hradci. Kralove. Suppl. 10(5):649-655
- 281 Guggenheim, R., J. Ilan, M. Postick, and E. Tal. 1963  
Prevention of meat anemia in mice by copper and calcium  
J. Nutr. 79(2):245-250
- 282 Guilbault, P., and E. Caraboeuf. 1962  
Comparison of the sensitivity of the myocardium of the rat and guinea pig to ions  
Compt. Rend. Congr. Natl. Soc. Savantes, Sect. Sci. 87:1285-1298
- 283 Gulyi, M.P., D.Ya. Vasilenko, D.A. Mel'nichuk, M.M. Shabel'nik, G.Ya. Korniyaka, M.M. Os'makovi, and L.M. Kolesnichenko. 1968  
Increasing the productivity of agricultural animals and fowl  
U.S.S.R. Pat. 210,646 issued Feb. 6, 1968
- 284 Gusev, G.P. 1970  
Renal cation excretion in dogs following injection of CaCl<sub>2</sub> and MgCl<sub>2</sub> following initial water diuresis or antidiuresis  
Fiziol. Zh. SSSR Sechenov. 56:782-790
- 285 Guyot, Rene. 1929  
Darnsteine medikamentosen ursprungs  
J. Pharm. Chim. 10(8):500-505
- 286 Gwathmey, J.I. 1928  
Synergism of magnesium sulphate and morphine  
J. Am. Med. Assoc. 91:1774-1776
- 287 Haarmann, Walter, and Kurt Bartscher. 1936  
Über den einfluss der chloride der alkalien und erdalkalien auf die stärkeverzuckerung durch diastatische fermente  
Biochem. Z. 283:301-311
- 288 Haase, E. 1931  
Value of magnesium sulphate  
Med. Welt. 5:1494-1496
- 289 Haden, Russell L., and Thomas G. Orr. No date  
Die wirkung anorganischer salze auf die chemischen veränderungen im blut des hundes nach verschluss des zwölffingerdarms  
J. Exp. Med. 39:321-330



- 290 Hadnagy, C., and I. Klepsz. 1957  
Mitotische Zellteilung und Antikörperbildung.  
(Mitotic cell division and antibody formation)  
Z. Immunitätsforsch. 114(5/6):493-501
- 291 Haffner, F., and W. Schaid. 1943  
As purgative and laxative  
Munch. Med. Wchnschr. 90:311
- \* 292 Haggard, Harold W., and Leon A. Greenberg. 1932  
Antidotes for strychnine poisoning  
J. Am. Med. Assoc. 98(14):1113-1136
- 293 Hahn, J., D.J. Tuma, and M.A. Quaife. 1967  
Solvent extraction and determination of magnesium  
in biological materials  
Anal. Chem. 39(10):1169-1171
- 294 Haldi, J., J. Larkin, and P. Wright. 1929  
Effects produced on weight relations in rabbit's  
brain by anesthetics and intravenous injections  
Am. J. Physiol. 88:112-116
- 295 Hansen, Olga S. 1921  
Magnesiumsulfat bei Arsenikvergiftung  
J. Pharmacol. Exp. Ther. 17:105-113
- 296 Hanzlik, P.J., and Elisabeth Prescho. No date  
Therapeutische Wirkung verschiedener Stoffe bei  
chronischer Vergiftung von Tauben  
durch metallisches Blei  
J. Pharmacol. Exp. Ther. 21:131-143
- 297 Harris, A.S., A. Estandia, H.T. Smith, R.W.  
Olsen, T.J. Ford, Jr., and R.F. Tillotson. 1953  
Magnesium sulfate and chloride in suppression of  
ectopic ventricular tachycardia accompanying  
acute myocardial infarction  
Am. J. Physiol. 172:251-258
- 298 Harrison, W.W., M.G. Netsky, and M.D. Brown. 1968  
Trace elements in human brain: copper, zinc, iron  
and magnesium  
Clin. Chim. Acta 21:55-60
- 299 Hartwich, Adolf. 1926  
Einfluss pharmakologisch wirksamer Substanzen auf  
die isolierte Froschniere. I. Mitt. Methodik,  
Einfluss des mechanischen und osmotischen  
Druckes, der Wasserstoffionenkonzentration, des  
Zuckers und des Magnesium- und Natriumsulfates.  
[Effect of pharmacologically active substances  
on the isolated frog-kidney. I. Method, effect of  
mechanical and osmotic pressure, of pH, of sugar  
Arch. Exp. Path. Pharmacol. 111(1/2):81-98
- 300 Hassib, Moquib Ali. 1964  
Biochemical changes in animals by the action of  
glucocorticoids and electrolytes  
Egypt. Pharm. Bull. 44(4):209-215
- 301 Haubrich, R. 1939  
Über die Retraktion bei der Blutgerinnung unter  
Druck  
Pflügers Arch. Ges. Physiol. 243(1):39-42
- 302 Haugaard, Niels, Ella S. Haugaard, and Nam Hea Lee  
1969  
Role of magnesium, phosphate and ATP in the  
regulation of calcium uptake by rat-  
liver mitochondria  
Proc. Kon. Ned. Akad. Wet. Ser. C Biol. Med. Sci.  
72(1):1-15
- 303 Haury, V.G. 1939  
Effect of intravenous injections on volume of  
extremities  
J. Lab. Clin. Med. 24:951-952
- \* 304 Haury, V.G., and A. Cantarow. 1940  
Effect of magnesium sulfate on serum and  
peritoneal fluid calcium, magnesium and  
inorganic phosphorus  
Proc. Soc. Exp. Biol. Med. 43:335-336
- 305 Haury, Victor G. 1939  
The effect of intravenous injections of magnesium  
sulfate on the vascular system  
J. Pharmacol. 65:453-460
- 306 Hayashi, Kazuji. 1927  
Vergleichende Untersuchung über die  
pharmakologische Reaktion des Vorhofes und  
der Kammer (Jap.)  
Folia Pharmacol. Jap. 5(1/2):73-85
- 307 Hazard, R., and E. Savini. 1952  
Nicotinolytic of magnesium ion  
Arch. Int. Pharmacodyn. Ther. 88(4):393-400
- \* 308 Hazard, R., and L. Wurmsier. 1931  
Action on excitability of sympathetic nervous  
system  
Compt. Rend. Soc. Biol. 108:944-946
- 309 Hazard, R., and L. Wurmsier. 1932  
Chloride, action on renal vasoconstrictors  
Compt. Rend. Soc. Biol. 110:525-528
- 310 Hazard, Rene, and Lise Wurmsier. 1938  
Demonstration, on the intestine in situ, of the  
adrenaline discharge-provoking action of the  
potassium ion  
Compt. Rend. Soc. Biol. 128:491-493
- 311 Heagy, F.C., and A.C. Burton. 1948  
Chloride, effect of intravenous injection on body  
temperature of unanesthetized dog, with  
observations on magnesium levels and body  
temperature in man  
Am. J. Physiol. 152:407-416
- 312 Hedon, L. 1931  
Sur une cause d'erreur dans l'inscription de la  
pression arterielle par la technique classique  
Compt. Rend. Soc. Biol. 106(11):893-895
- 313 Heffer, A. 1927  
CaCl<sub>2</sub> and MgCl<sub>2</sub> action on duodenal secretion  
Russ. Klin. 7:838-852
- 314 Heianzan, N. 1925  
Über die Ionenwirkung auf die Leberfunktion und  
den Zuckerstoffwechsel. I.  
Mitt. Elektrolytenwirkung auf die sekretorische  
Leberfunktion. II. Mitt. Elektrolytenwirkung auf  
den Zuckerstoffwechsel. III. Mitt. Die Wirkung  
des Ca und Na auf die Leberfunktion in Bezug auf  
den Zuckerstoffwechsel  
J. Biochem. 5:171-184
- 315 Heller, V.G., and C.H. Larwood. 1930  
Salziges Trinkwasser  
Science 71:223-224
- 316 Hemmeter, J.C. 1922  
Effect of magnesium sulphate on liver and biliary  
apparatus  
N.Y. Med. J. 116:645-648
- 317 Henry J. Brown, Cleveland, Ohio. 1923  
Arzneimittel gegen Nicotinvergiftungen  
A. Pat. 1,474,140 issued Nov. 13, 1923
- 318 Herrmann, Julian B. 1941  
Effects of certain drugs on temperature  
regulation and changes in their toxicity, in  
rats exposed to cold  
J. Pharmacol. 72:130-137
- 319 Hessler, Anita Y. 1968  
Magnesium reversal of proflavine inhibition of  
bacteriophage T2H  
Virology 35(4):619-622
- 320 Hillis, W.G., Clarence B. Ammerman, Phillip E.  
Loggins, and Rudolph Hendrickson. 1969  
Citrus pulp dehydrated with magnesium oxide and  
its subsequent use as a feed  
Proc. Fla. State Hort. Soc. 1968(81):297-301
- 321 Hirschberg, Louis. 1928  
Sekretorische Funktion der Leber, des Pankreas  
und des Magens nach Einführung von MgSO<sub>4</sub> ins  
Duodenum. [Effect of MgSO<sub>4</sub> in duodenum on  
secretion by liver, pancreas and stomach]  
Arch. Verdau.-Kr. 44(5/6):393-405

- 322 Hirschfelder, A.D. 1933  
Enhanced action of morphine in experimental nephrosis after oral ingestion of magnesium sulphate  
Proc. Soc. Exp. Biol. Med. 30:1057-1058
- 323 Hirschfelder, A.D. 1934  
Clinical manifestations of high and low plasma magnesium; dangers of epsom salt (magnesium sulphate) purgation in nephritis  
J. Am. Med. Assoc. 102:1138-1141
- 324 Hirschtelder, Arthur D. 1934  
Effect of renal insufficiency upon plasma magnesium and magnesium excretion after ingestion of magnesium sulfate  
J. Biol. Chem. 104(3):647-653
- 325 Hodgkins, J.R., M. Ragheb, H.I. Abdou, and M. Nashed. 1943  
Euthanasia of equines. (The action of magnesium)  
Vet. Rec. 55(28):269-270
- 326 Hoesch, K. 1924  
Zur magnesiumwirkung  
Munch. Med. Wchschr. 71:1679-1680
- 327 Holland, William C. 1954  
The action of anesthetic agents on the loss of potassium from isolated guinea pig auricles  
J. Pharmacol. Exp. Ther. 3(1):1-8
- 328 Hollander, Vincent P., David E. Smith, and Thelma E. Adamson. 1959  
Studies on estrogen-sensitive transhydrogenase. The effect of estradiol-17-beta on alpha-ketoglutarate production in noncancerous and cancerous human breast tissue  
Cancer 12(1):135-138
- 329 Holobut, Wieslaw. No date  
Wirkung des strychnins auf die motorischen nerven und der einfluss der alkaliionen  
Compt. Rend. Soc. Biol. 100:1232-1234
- 330 Honoki, Janos, Istvan Lukacs, and Constantin E. Sekeris. 1967  
Effect of ammonium sulfate and other salts on the ribonuclease activity of isolated rat liver nuclei  
Hoppe-Seyler's Z. Physiol. Chem. 348(11):1392-1394
- 331 Hoper, M. 1963  
Microdetermination of magnesium and calcium in animal tissue  
Experientia 19:367-368
- 332 Hori, Ichiro, and Nobuaki Inoue. 1957  
The clouding of sake. II. The component sugars of the clouding matter  
Hakko Kagaku Zasshi 35:419-422
- 333 Horimoto, Akira. 1962  
Variations of metabolic products in the blood of animals exposed to high and low temperatures and effect of various drugs thereon. I. Seasonal variations of metabolic products in the blood of rabbits exposed to high temperature  
Nara Igaku Zasshi 13:20-24
- 334 Horn, D.B., and A.L. Latner. 1963  
The estimation of magnesium by atomic absorption spectrophotometry  
Clin. Chim. Acta 8:974-976
- 335 Houghtaling, Cora I., Francis S. Houghtaling, Nestor E. Houghtaling, and Robert W. Kilburn. 1966  
Citrus fruit juice concentrates  
U.S. Pat. 3,227,562 issued Jan. 4, 1966
- 336 Huang, T'ing-Fei. 1964  
Effects of various ions, glucose, and insulin on the electrocardiogram in experimental animals  
Kuo Li Taiwan Tah Hsueh Yi Hsueh Yuan Yen Chiu Pao Kao 10(1):54-65
- 337 Hubbard, J.I., R.F. Schmidt, and T. Yokota. 1965  
Effect of acetylcholine on mammalian motor nerve terminals  
J. Physiol. 181(4):810-829
- 338 Huber, John T., Roy S. Emery, John William Thomas, and I.M.K. Youser. 1969  
Milk fat synthesis on restricted-roughage rations containing whey, sodium bicarbonate, and magnesium oxide  
J. Dairy Sci. 52(1):54-59
- 339 Huidobro, F., and J.V. Loco. 1939  
The effect of MgSO<sub>4</sub> on the superior cervical ganglion  
Anal. Acad. Biol., Univ. Catol. Chile 3:133-145
- 340 Hulyi, M.F., D.O. Mel'nychuk, and M.D. Klymenko. 1968  
Vplyv bikarbonatu natriyu, Mn++, Mg++ i Zn++ na intensyvniat'onovlennyya bilkiv, hlikohenu ta lipidiv u pechintsita m'yazakh kroliu. (Effect of sodium bicarbonate, Mn++, Mg++, and Zn++ on the intensity of renewal of proteins, glycogen and lipids in the rabbit liver and muscles)  
Ukr. Biokhim. Zh. 40(2):167-172
- 341 Hunt, B.J. 1969  
The estimation of magnesium in plasma, muscle and bone, by atomic absorption spectrophotometry  
Clin. Chem. 15(10):979-996
- 342 Husdan, H., and A. Rapoport. 1969  
Estimation of calcium, magnesium and phosphorous in diet and stool  
Clin. Chem. 15(8):669-679
- 343 Huston, John, Jr., D.P. Wallach, and Geo. J. Cunningham. 1952  
Pulmonary reaction to barium sulfate in rats  
Arch. Pathol. 54:430-438
- 344 Hutchinson, Harry T., Myron M. Nichols, Charles R. Kuhn, and Alois Vasicka. 1964  
Effects of magnesium sulfate on uterine contractility, intrauterine fetus, and infant  
Am. J. Obstet. Gynecol. 88(6):747-758
- 345 Iberti, U. 1939  
Modifications of ratio of porphyrin in blood, urine and feces under influence of calcium chloride and magnesium sulfate in normal and pathologic conditions, with special reference to saturnism  
Clin. Med. Ital. 70:237-248
- 346 Iida, C., K. Fuwa, and W.E.C. Wacker. 1967  
A general method for magnesium analysis in biological materials by atomic absorption spectroscopy  
Anal. Biochem. 18:18-26
- 347 Inamura, Shozo. 1951  
Effect of various sorts of ions on the sensoria of the labyrinth of eel (*Anguilla japonica*)  
Tohoku J. Exp. Med. 54(2):145-150
- 348 Inoue, T. 1937  
Permeability of Bowman's capsule in the kidney. I. The effect of diuretics on its permeability to water and dye  
Sei-i-kai Med. J. 56(10):1866-1887
- 349 Irving, E.A., and P.S. Watts. 1961  
Estimation of calcium and magnesium in blood serum by the cathode ray polarograph  
Biochem. J. 79:429-432
- 350 Ivy, A.C., and G.B. McIlvain. 1923  
Die erregung der magensekretion durch applikation von substanzen auf die duodenal- und jejunal-schleimhaut  
Am. J. Physiol. 67:124-140
- 351 Iwao, Taijiro. 1936  
The toxicity of metals  
Jikken Yakubuts. Z. 10:357-410

- 352 Iwata, Osamu. 1963  
The action potential of the optic nerve. II.  
Effects of minerals (Jap.)  
Nippon Ganka Gakkei Zasshi 67(10):1499-1514
- 353 Jackson, S.H., and F. Brown. 1964  
Simultaneous determination of calcium and  
magnesium of serum by a single chelometric  
titration  
Clin. Chem. 10(2):159-169
- 354 Jacobasch, G., W. Binus, H. Vierus, and C. Boese.  
1967  
Disturbances of the dental metabolism in dogs  
caused by magnesium deficiency (Ger.)  
Acta Biol. Med. Ger. 18(3):409-422
- 355 Jaswal, A.S. 1970  
Effects of various chemical blanchings on the  
texture of French fries  
Am. Potato J. 47(1):13-18
- 356 Jaulmes, P., and G. Pons. 1947-1948  
Rapid test for adulterants in grape musts  
Trav. Soc. Pharm. Montp. 7:66-68
- 357 Jensen, Soren Lovstrup. 1945  
Influence of ionic strength on the acid liver  
phosphatase  
Acta Physiol. Scand. 9:357-361
- 358 Jirgensons, Br. 1928  
Beitrage zur kolloidchemie der hamoglobine. I  
Biochem. Z. 194:140-150
- 359 Joglekar, G.V., and J.H. Balwani. 1966  
Effect of CNS (central nervous system)  
depressants on hexobarbitone sleeping time in  
rats  
Arch. Inst. Pharmacodyn. Ther. 159(1):140-143
- 360 Johannsson, O., G. Perrault, L. Savoie, B.  
Tuchweber. 1968  
Action of various metallic chlorides on calcemia  
and phosphatemia  
Brit. J. Pharmacol. Chemother. 33(1):91-97
- 361 Johnson, R.G., and B.L. Stierbold. 1959  
The effect of urea magnesium chloride, and a low  
protein diet, on the production of calcium  
oxalate vesicle calculi in the rat  
J. Urol. 81(5):691-694
- 362 Jonas, E.E.E. 1969  
Effect of magnesium sulphate and cryoprotective  
agents on free fatty acid formation in cold-  
stored fish muscle  
J. Fish. Res. Board Can. 26(8):2237-2240
- 363 Jonderko, Gerard, and Zdzislaw Dabrowski. 1964  
The effect of heavy metal salts and calcium  
disodium Versenate on the activity of  
serum phosphomonoesterase. I. In vitro studies  
Arch. Immunol. Ther. Exp. 12(5):592-601
- \* 364 Jones, J.E., P.C. Desper, S.R. Shane, and E.B.  
Flink. 1966  
Magnesium metabolism in hyperthyroidism and  
hypothyroidism  
J. Clin. Invest. 43(6):891-900
- 365 Jones, J.E., R. Manalo, and E.B. Flink. 1967  
Magnesium requirements in adults  
Am. J. Clin. Nutr. 20(6):632-635
- 366 Jullien, A., J. Ripplinger, J. Cardot, M. Joly,  
and J. Moulin. 1959  
The action of individual or associated alkaline  
and alkaline-earth ions on the pulmonary  
circulation in the frog  
J. Physiol. 51:495-496
- 367 Jung, Frederic T., and Franklin Cook. 1927  
Die wirkung intraperitoneal gereichten  
magnesiumsulfats an parathyreodextonierten ratten  
Proc. Soc. Exp. Biol. Med. 24:586-587
- 368 Junge, Douglas. 1967  
Multi-ionic action potentials in molluscan giant  
neurons  
Nature 215(5100):546-548
- 369 Juul-Christensen, E.K., H.P.M. Kerckhoffs, and T.  
Huizinga. 1967  
The influence of gastric antacids on the release  
in vitro of tetracycline hydrochloride  
Pharm. Weekbl. 102(21):463-473
- 370 Kaidin, D.A. 1968  
Effect of malic acid and alkali-metal and  
alkaline earth-metal carbonates on  
the development of alimentary  
hypercholesterolemia and lipidosis of the aorta  
in rabbits  
Byull. Eksp. Biol. Med. 66(12):29-31
- 371 Kakkar, Bimla, and D.W. Mullick. 1963  
Effect of intramuscular injection of magnesium  
sulfate solution on the growth rate and serum  
composition in rats  
Ann. Biochem. Exp. Med. (Calcutta) 23(6):229-230
- 372 Kalant, H., and R.A. Hickie. 1968  
Effects of divalent cations of K<sup>+</sup> transport in  
rat liver and morris hepatoma 5123tc  
Cancer Res. 28:2086-2091
- 373 Karo, Armand, M., Jr., Metin O. Unal, Oliver  
Carrier, Jr., Clifton R. Daniel, Watts R. Webb,  
and William C. Holland. 1967  
Preliminary observations on drug responses in  
viably frozen mammalian hearts  
Cryobiology 3(4):335-340
- 374 Kato, Hidekazu. 1939  
The influence of some pharmaceutical agents on  
the blood sugar and the contents of  
glycogen, creatine and phosphate of the muscle  
of rabbits  
Mitt. Med. Akad. Kioto 25:406-407
- 375 Kawabata, Juro. No date  
Uber die zusammenwirkungen von cocain und  
ammonium, calcium, strontium, barium sowie  
magnesium  
Folia Pharmacol. Jap. 4:35-57
- 376 Kawamura, Yojiro, Junichi Okamoto, and Masaya  
Funakoshi. 1968  
A role of oral afferents in aversion to taste  
solutions  
Physiol. Behav. 3(4):537-542
- 377 Kayaalp, S. Oguz, and Robert J. McIsaac. 1970  
Differential blockade and potentiation of  
transmission in a sympathetic ganglion  
J. Pharmacol. Exp. Ther. 173(1):193-204
- 378 Keeser, E. 1931  
Effect on action of enzymes  
Arch. Exp. Pathol. Pharmacol. 160:663-686
- 379 Keeser, E. 1939  
The effect of pharmaceuticals on the activity of  
histaminase  
Dtsch. Med. Wochschr. 65:94-95
- 380 Keeser, Ed. 1944  
Etiology and therapy of atherosclerosis  
Dtsch. Z. Verdau. Stoffwechselkr. 8:44-48
- 381 Keeser, Ed. 1947  
On the character of cholesterol solution  
Arch. Exp. Path. Pharmacol. 204:36-40
- 382 Kegley, Larry M., Bruce W. Brown, and Allen R.  
Berutzen. 1969  
Mesocostoides corti: inorganic components in  
calcareous corpuscles  
Exp. Parasitol. 1969(25):85-92
- 383 Keil, W. 1934  
Magnesium, effect of combining tribrom-ethanol  
with magnesium chloride  
Arch. Exp. Pathol. Pharmacol. 174:490-492

- 384 Kelco Co. 1969  
Canned created meat-containing food products containing a water-soluble alginate and a soluble magnesium salt  
U.S. Pat. 3,480,450 issued Nov. 25, 1969
- 385 Kesik, Maria. 1967  
Effect of some sodium and magnesium salts on the absorption of glucose in the small intestine of the rat in vivo (Pol.)  
Stul. Soc. Sci. Torun., Sect. G 3(1):42 pp
- 386 Khan, Wali Mohammad. 1943  
Pitto or gillar in sheep and goats  
Indian Farming 4(2):85
- 387 Kim, Young Hi. 1967  
Effect of salts on the inactivation of antiphage serum by drying  
New Med. J. (Seoul, Korea) 10(7):98-100
- 388 Kirsch, Ulrich. 1970  
Beginning of rigor mortis in the ischemic guinea pig heart in normothermia. Effects of procaine, potassium, and magnesium (Ger.)  
Arzneimittel-Forschung 20(8):1071-1074
- 389 Kirsner, Joseph E. 1941  
A further study of the effect of various antacids on the hydrogen-ion concentration of the gastric contents  
Am. J. Dig. Dis. 8:53-56
- 390 Kisch, Bruno. No date  
Differenzierende Wirkungsanalysen von Herzgiften. V. Mitt. Die Wirkung von  $MgCl_2$ ,  $SeCl_2$  und  $BaCl_2$  auf die Herzreizbildung beim Kaltblutler  
Arch. Exp. Pathol. Pharmacol. 124:210-230
- 391 Kisin, I.E. 1967  
Effect of vasodilating agents on basal tonicity and vascular response reactions following electric stimulation of sympathetic nerves  
Farmakol. Toksikol. 30(6):674-678
- \* 392 Klein, B., and M. Oklander. 1967  
The automated fluorometric determination of serum magnesium. II. Procedure using 8-hydroxyquinoline-5-sulfonic acid  
Clin. Chem. 13(1):26-35
- \* 393 Klein, B., J.H. Kaufman, and E. Oklander. 1967  
Automated atomic absorption spectrophotometry. II. The determination of serum magnesium  
Clin. Chem. 13(9):788-796
- 394 Kleinau, H., and Ernest Schubert. 1969  
Variations in length of heart muscle fibers from sheep after local application of calcium chloride compared with the effect of magnesium, strontium, sodium, and potassium chlorides  
Pfluegers Arch. 306:135-145
- 395 Kobayashi, Choe. 1936  
Extraction of phosphatase and carboxylase of yeast  
J. Biochem. (Japan) 24:369-387
- 396 Kochetkova, G.V. 1960  
The influence of reserpine and magnesium sulfate on the coagulability of the blood in experimental dicoumarin hypoprothrombinemia  
Nauch. Tr., Kuban. Gos. Med. Inst. 18:87-90
- 397 Kochetkova, G.V. 1961  
The mechanism of action of some recent antihypertensive drugs on the rate of blood clotting  
Mater. 3-ei Tret'eil Nauchn. Konf. Uch. Goroda Samark., Sekts. Med. Nauk, Samark. Sb. 1961:72-73
- 398 Kochetkova, G.V. 1962  
Effect of magnesium sulfate and dieldrol on blood clotting rate and arterial blood pressure in experimental hypoprothrombinemia  
Nauchn. Tr. Samark. Med. Inst. 21:213-219
- 399 Kofler, L., and R. Fischer. No date  
Magnesiummuckose durch orale verabreichung von Magnesiumsulfat und saponin  
Arch. Exp. Pathol. Pharmacol. 130:319-322
- 400 Kofler, L., and R. Fischer. 1930  
Eine biologische prufung der darmresorption von kalkpreparaten  
Arch. Exp. Path. Pharmacol. 149:326-335
- 401 Kohler, F. Peter, and Charles A.W. Uhle. 1966  
Oral magnesium intake and its effect on various urinary constituents in the prophylaxis of urolithiasis  
J. Urol. 96(5):812-815
- 402 Kolda, Jaroslav. 1928  
Vliv koloidniho hydroxydu horecnateho na srazeni krve. (The influence of colloidal magnesium hydrate on the clotting of blood)  
Biol. Spizy Vys. Skoly Zverolek. Brno 7:93-96
- 403 Kono, Tomosaburo. 1959  
On the absorption of calcium chloride and magnesium sulfate from rabbit intestine. (Jap.)  
Folia Pharmacol. Jap. 55(5):1089-1114
- 404 Kooistra, John A., and John A. Troller. 1968  
Food preservative compositions and method for inhibiting microbial growth in food  
U.S. Pat. 3,404,987 issued Oct. 8, 1968
- \* 405 Koshy, K.T., R.W. Duvall, A.E. Troup, and J.W. Pyles. 1965  
Factors involved in the browning of spray-dried lactose  
J. Pharm. Sci. 54(4):549-554
- 406 Kotzareff, A., J. De Morsier, and A. Morin. 1929  
Action du chlorure de magnesium sur l'evolution du cancer greffe sur la souris blanche. (Action of  $MgCl_2$  on evolution of cancer graft in white mice)  
Bull. Acad. Med. (Paris) 102(25):11-12
- 407 Kovacs, Edit F., and Gyorgy Bot. 1965  
Inhibition of phosphoglucosutase by anions, phosphate esters and Mg ion  
Acta Physiol. Acad. Sci. Hung. 27(4):327-340
- 408 Kovtun, S.D. 1960  
Labil'nist' nerva, al'terovanobo khloridyda bariyu ta khloridom mahniyu. (Lability of a nerve altered with barium and magnesium chloride) (Ukr.)  
Fiziol. Zh. 6(4):481-489
- 409 Kravchenko, A.T., and G.L. Ryazanova. 1967  
The effect of magnesium salts on thermal inactivation of myxoviruses (Russ.)  
Vopr. Virusol. 12(3):342-348
- 410 Krylova, N.I. 1965  
Use of aminazine in a complex of hypotension agents in hypertensive crises (Russ.)  
Klin. Med. 43(9):124-125
- 411 Krylova, N.N., and L.D. Zueva. 1953  
Stabilizing blood used for food and industrial purposes  
Tr. Vses. Nauch.-Issled. Inst. Myasn. Prom. 5:139-145
- 412 Kumar, D. 1963  
In vitro and in vivo effects of magnesium sulfate on human uterine contractility  
Am. J. Obstet. Gynecol. 86:1036-1040
- 413 Kunicki, T. 1967  
A case of acute intoxication with magnesium sulphate administered into the duodenum  
Wiad. Lek. 20:77-79
- 414 Kunitsa, L.D. 1968  
Gipotenzivnoe i kaliureticheskoe deistvie gipotiazida. (Hypotensive and kaliuretic effect of hydrochlorothiazide)  
Kardiologiya 8(9):110-116

- 415 Kuppaswamy, S., and R.G. Rao. 1970  
Dehydration of green peas  
J. Food Sci. Technol. (Mysore) 7(Suppl.):18-21
- 416 Kusano, Chuji. 1965  
Stimulating effectiveness of cations and anions on the tarsal chemoreceptors of the cabbage butterfly (*Pieris rapae crucivora*)  
Tottori Nogakkaiho 18:1-22
- 417 Kusiak, Marian, and Jan Markiewicz. 1961  
Śmiertelne zatrucia po doustnym stosowaniu roztworu siarczanu magnezu. (Fatal intoxication caused by intraduodenal administration of magnesium sulphate. Report of 2 cases)  
Pol. Iyg. Lek. 16(11):674-679
- 418 Kusnetzow, A.J. 1926  
Action of salts of alkalis and alkali earths on adrenal secretion  
Z. Ges. Exp. Med. 48(6):671-678
- \* 419 La Barre, J., and G. Kettenmeyer. 1941  
Effects of hypnotics with paralyzing action on thalamic centers; effect of barbital sodium and magnesium sulfate on epinephrine secretion  
Arch. Int. Pharmacodyn. Ther. 66:315-324
- 420 La Barre, J., and G. Kettenmeyer. 1941  
Effects of hypnotics with paralyzing action on thalamic centers; effects of magnesium sulfate on gastric secretion and motility  
Arch. Int. Pharmacodyn. Ther. 66:305-314
- \* 421 La Barre, J., and O. Vesselovsky. 1942  
Study of hypnotics with paralyzing action on thalamic centers; effect of magnesium sulfate on external secretion of pancreas  
Arch. Int. Pharmacodyn. Ther. 67:267-275
- 422 La Bossa, F. Buonomo. 1931  
Effects of magnesium chloride on hemolytic amoebocytes  
Boll. Ist. Sieroter. Milan. 10:665-669
- 423 La Bossa, F. Buonomo. 1931  
Effects on micro-organisms in vitro  
Gior. Batt. Immunol. 6:248-258
- 424 Labkovskiy, B.M. 1967  
Effect of magnesium salts on induction by urethan of pulmonary adenomas in mice  
Vopr. Onkol. 13(8):121
- 425 Laforet, C. Colmeiro. 1932  
Magnesium sulphate and tribrom-ethanol  
Med. Latina 5:487-495
- 426 Lamberth, J.L. 1969  
Effect of magnesium sulfate administration on the rumen contents of normal cattle on high-grain rations  
Austral. Vet. J. 45(5):223-224
- 427 Lanari, Alfredo. 1939  
Pharmacological study of myotonia  
Arch. Int. Pharmacodyn. 61:360-369
- 428 Lanyi, Janos K., and Joann Stevenson. 1969  
Effect of salts and organic solvents on the activity of *Halobacterium cutirubrum* catalase  
J. Bact. 98(2):611-616
- 429 Larski, Zdzislaw. 1965  
The influence of magnesium chloride on the thermal inactivation rate of Newcastle disease virus  
Bull. Vet. Inst. Pulawy 9(1/2):1-7
- 430 Lassner, A. 1943  
Osmotherapy with electrolyte solutions (magnesium sulfate) in secondary diphtheria  
Wien. Klin. Wchnschr. 56:690
- 431 Laver, W.G., A. Neuburger, and S. Udenfriend. 1958  
Initial stages in the biosynthesis of porphyrins. 1. The formation of delta-aminolaevulinic acid by particles obtained from chicken erythrocytes  
Compt. Biochem. J. 70(1):4-14
- 432 Lawhoin, C.C. 1944  
Treatment of Meniere's syndrome (with intravenous injections of magnesium sulfate)  
Med. Rec. 157:220
- 433 Lazzara, R.K., T.K. Yun, and G.E. Burch. 1965  
Effects of magnesium sulfate, calcium gluconate, potassium chloride, and ammonium chloride on the urinary excretion of magnesium in patients with idiopathic cardiomyopathy and congestive heart failure  
Am. J. Med. Sci. 250(2):193-198
- \* 434 Lebduska, J., and F. Cervinka. 1930  
Effect of colloidal magnesium hydroxide and sodium thiosulfate on white blood corpuscles  
Compt. Rend. Soc. Biol. 103:366-368
- 435 Lebedinskaja, S.I., and E.N. Speranskaja-Stepanova. 1930  
Action of epinephrine on contraction of striated muscle of warm-blooded animals in magnesium sulphate poisoning  
Z. Ges. Exp. Med. 71:323-328
- 436 Lehman, Arnold J. 1951  
Chemicals in foods: report to the Association of Food and Drug Officials on current developments  
Assoc. Food Drug Offic. U.S., Q. Bull. 15:82-89
- 437 Leibowitz, J., and A. Schweitzer. 1930  
Vergleich der wirkungen von  $MgCl_2$  auf atmungsgrosse und kontraktilität des trosschventrikels  
Biochem. Z. 229:291-295
- 438 Lendle, L. No date  
Untersuchungen über den verschiedenen angrißspunkt einiger narkotika im zentralnervensystem  
Arch. Exp. Path. Pharmacol. 143:108-116
- 439 Leo, Cosimo. 1926  
Anestesia generale col solfato di magnesio per via sottocutanea. Ricerche sperimenta  
19. (General anesthesia with  $MgSO_4$  administered subcutaneously)  
Clin. Chir. 29(11):1096-1111
- 440 Lepskaya, M.V., and A.F. Gol'dberg. 1936  
Influence of x-ray radiation and nonspecific stimuli on the carbohydrate metabolism and erythropoiesis in experimental sarcoma in rats. III  
Z. Krebsforsch. 44:432-439
- 441 Lesnik, Helena, J.W. Guzek, and W.Z. Traczyk. 1969  
The release of antidiuretic hormone from the pituitary stalk after intravenous calcium chloride or magnesium chloride infusion in rats  
J. Endocrinol. 45(1):83-90
- 442 Leusen, I. 1948  
Influence de la concentration en magnesiu du liquide Cephalorachidien sur le systeme vasomoteur. (The effect of magnesium concentration of the cephalorachidian fluid on the vasomotor system)  
Arch. Int. Pharmacodyn. Ther. 77(1):50-51
- 443 Levy, J. Cl. 1968  
Modifications of muscle excitability by depolarizing and repolarizing substances  
Prod. Probl. Pharm. 23(5):303-309
- 444 Lieberman, T.Yu. 1963  
Effects on cerebrovascular tonus caused by some headache remedies in hypertensive patients  
Farmakol. Toksikol. 26(5):573-578
- 445 Ligon, E. Wm., Jr. 1943  
Cathartic salts on motility of Thirty-Vella jejunal loops  
Proc. Soc. Exp. Biol. Med. 52:282-284

- 446 Linetskii, G.S. 1967  
Ekskurgatornyi metod v sochetanii s  
vitaminoterapie i pri nekotorykh dermatozakh.  
(The purgative method in conjunction with  
vitamin therapy in some dermatoses)  
Sb. Nauch Tr. L'vov Nauch Obshchest Dermatovenerol 5:385-389
- \* 447 Lipsitz, P.J. 1971  
The clinical and biochemical effects of excess  
magnesium in the newborn  
Pediatrics 47:501-509
- \* 448 Lipsitz, P.J., and I.C. English. 1967  
Hypermagnesemia in the new born infant  
Pediatrics 40:856-862
- 449 Lister, D., and P.W. Ratcliff. 1971  
The effect of the pre-slaughter injection of  
magnesium sulphate on glycolysis and meat  
quality in the pig  
Proc. 2nd Int. Symp. Cond. Meat Qual. Pigs  
pp. 139-144
- 450 Lium, R., and H.W. Florey. 1939  
The action of magnesium sulphate on the intestine  
of the cat  
Q. J. Exp. Physiol. 29(3):303-319
- 451 Ljunggren, E. 1965  
Prophylactic measures against oxalate kidney  
stone formation  
Z. Urol. Nephrol. 58(7):507-512
- 452 Loiseleur, Jean, and Renee Colliard. 1937  
The adsorption of polypeptides by the proteins of  
blood plasma  
Compt. Rend. 205:261-263
- 453 Lomborg, Florence. 1950  
The action of sodium and magnesium sulfate on  
Salmonellae, Shigellae and Vibrios.  
Am. J. Dig. Dis. 17(2):38-39
- 454 Lott, J.A., and T.S. Herman. 1971  
Increased autocatalyzer dialysis of calcium and  
magnesium in presence of protein  
Clin. Chem. 17(7):614-621
- 455 Luco, J.V., R. Pichard, and F. Huidobro. 1940  
Action of  $MgSO_4$  on skeletal muscles  
Anal. Acad. Biol., Univ. Chile 3:69-75
- 456 Luhrig, B. 1926  
Beitrage zur beurteilung von kakao  
Pharm. Zentralhalle 67:129-135
- 457 Lumiere, Auguste, and Malespine. 1928  
Schutz gegen die anaphylaktoiden shocks mit hilfe  
von magnesiumhyposulfit  
Compt. Rend. Acad. Sci. 187:736-737
- 458 Lur'e, G.S., and B.A. Temper. 1930  
Influence of cations on gastric secretion  
Odess. Med. Z. 5(3/5):261-264
- 459 Lynch, H.T., H.M. Lemon, H.J. Henn, R.J.  
Ellingson, and R.L. Grissom. 1964  
Vitamin D-intoxicated patient with  
hypoparathyroidism. Hypercalcemia,  
acute cerebellar ataxia, and EEG  
(electroencephalographic) changes: magnesium  
sulfate therapy  
Arch. Intern. Med. 114(3):375-380
- 460 Lynch, William P. 1956  
Experimental modification of the rate of  
metamorphosis of Bugula larvae  
J. Exp. Zool. 133(3):589-612
- 461 Maaske, C.A., and B. Gibson. 1939  
Effects on denervated mammalian muscle  
Am. J. Physiol. 127:486-491
- 462 Mac Andrews and Forbes Co. 1934  
Abfuhrmittel  
U.S. Pat. 1,976,668 issued Oct. 9, 1934
- \* 463 MacIntyre, I. 1963  
An outline of magnesium metabolism in health and  
disease--a review  
J. Chronic Dis. 16:201-215
- \* 464 MacIntyre, I., S. Boss, and V.A. Troughton. 1963  
Parathyroid hormone and magnesium homeostasis  
Nature 198:1058-1060
- 465 Magnus-Levy, A. 1928  
Stationary metabolism of magnesium chloride  
injected subcutaneously; experimental study  
Z. Klin. Med. 107:671-673
- 466 Malanud, T., and B. Iussea. 1944  
Therapeutic dose of magnesium sulfate  
Prensa Med. Argent. 31:1461-1462
- 467 Malinconico, E. 1932  
Foscal (calcium phosphate-magnesium carbonate  
preparation)  
Rass. Clin. Ter. Sci. Aff. 31:34-36
- 468 Malkus, Zdenek. 1956  
Determination of magnesium oxide in cocoa powder  
Prun. Potravin. 7:524-526
- 469 Malmstadt, H.V., and T.P. Hadjiioannou. 1959  
Automatic titration of calcium or magnesium in  
blood serum  
Clin. Chem. 5(1):50-56
- 470 Marchi, C. 1930  
Possible significance of absence of alimentary  
magnesium chloride and greater frequency of  
carcinomas  
Riforma Med. 46:55-56
- 471 Marchi, C. 1930  
Deficiency of magnesium chloride and greater  
frequency of cancer; preliminary report  
Scr. Biol. 5:295-302
- 472 Mariano, Calderone. 1936  
Magnesiumsulfat in endovenoser gabe bei der  
behandlung von uraemie und hypertension  
Arch. Farmacol. Sper. Sci. Affini 61(35):60-79
- 473 Markovitz, Alvin, and Susan Sylvan. 1962  
Effect of sodium sulfate and magnesium sulfate on  
heteropolysaccharide synthesis in gram-negative  
soil bacteria  
J. Bact. 83(3):483-489
- 474 Markuse, K.P., and D.A. Losinsky. 1936  
Influence of nonspecific stimuli upon the course  
of experimental rat sarcoma. II. The combined  
effect of x-ray irradiation and mineralization  
Z. Krebsforsch. 44:422-431
- 475 Maruo, E. 1970  
Metoclopramide and electrocardiographic changes  
caused by KCl, BaCl<sub>2</sub>, CaCl<sub>2</sub>, MgCl<sub>2</sub>, aconitine, k-  
strophanthoside, vasopressin and by asphyxia  
Arzneimittelforschung 20:12-18
- 476 Martin, Lay. 1927  
Biliary, pancreatic and duodenal studies. I. The  
hydrogen ion concentration of successive  
portions of duodenal contents following  
stimulations with magnesium sulphate  
Arch. Intern. Med. 39(2):275-280
- \* 477 Martinek, R.G., and R.E. Berry. 1964  
Micromethod for the estimation of magnesium in  
biological fluids  
Clin. Chim. Acta 10:365-374
- 478 Marullaz. 1929  
Effect of magnesium salts on development of ter-  
tumors  
Bull. Acad. Med., Paris 102:48-50
- 479 Marullaz. 1930  
Action des sels de magnesium sur la genese des  
tumeurs du goudron  
Rev. Int. Med. Chir. 41(2):20-24

- 480 Maruyama, K., and H. Kobayashi. 1957  
Effect of desoxycorticosterone acetate upon the  
apyrase activity of rat liver mitochondria  
*Enzymologia* 18(2):135-141
- 481 Marxer, A. 1933  
Konservierung von pflanzlichen oelen  
Belg. Pat. 392,235 issued May 19, 1933
- 482 Masasune, H. No date  
Salze und blutzucker  
*Fukuoka-Ikwadaigaku-Zasshi* 20:3
- 483 Mashima, Shizuyoshi, and Yonosuke Ikeda. 1958  
Selection of mutagenic agents by the *Streptomyces*  
reverse mutation test  
*Appl. Microbiol.* 6:45-49
- 484 Maslov, N.M., and S.M. Sverdlov. 1950  
Relation between muscle impedance and its  
polarization potential in the presence of  
some chemical agents  
*Dokl. Akad. Nauk S.S.S.R.* 71:49-52
- 485 Massione, R., and F. Zanotti. 1940  
Lead in blood, urine and feces after injections  
of calcium (chloride) and magnesium sulfate  
*Med. Lavoro* 31:25-32
- \* 486 Massry, S.G., J.W. Coburn, and C.R. Kleeman. 1969  
Renal handling of magnesium in the dog  
*Am. J. Physiol.* 219(6):1460-1467
- 487 Massry, Shaul G., Jack W. Coburn, and Charles R.  
Kleeman. 1970  
Evidence for suppression of parathyroid gland  
activity by hypermagnesemia  
*J. Clin. Invest.* 49(9):1619-1629
- 488 Massry, Shaul G., Juan J. Ahumada, Jack W.  
Coburn, and Charles R. Kleeman. 1970  
Effect of magnesium chloride infusion on urinary  
calcium and sodium during reduction in their  
filtered loads  
*Am. J. Physiol.* 219(4):881-885
- 489 Matignon, M.C. 1930  
Sur la chaleur de dissolution limite de  
l'hyposulfite de soude et du sulfate de magnesie  
hydrates  
*Compt. Rend. Acad. Sci.* 190:429-431
- \* 490 Maxwell, George M., Robert B. Elliott, and  
Richard H. Burnell. 1965  
Effects of hypermagnesemia on general and  
coronary hemodynamics of the dog  
*Am. J. Physiol.* 208(1):158-161
- 491 McAleese, D.M., and R.M. Forbes. 1961  
The requirement and tissue distribution of  
magnesium in the rat as influenced  
by environmental temperature and dietary calcium  
*J. Nutr.* 73(1):94-106
- 492 McCann, H.G. 1959  
Determination of microgram quantities of  
magnesium in mineralized tissues  
*Anal. Chem.* 31(12):2091-2092
- 493 McCaughan, J.M., B.L. Sinder, and C.J. Sullivan.  
1938  
The external secretory function of the human  
pancreas. Physiologic observations  
*Arch. Intern. Med.* 61(5):739-754
- 494 McCutcheon, Morton, and Balduin Lucke. No date  
Der einfluss gewisser elektrolyte und  
nichtelektrolyte auf die durchlassigkeit  
lebender zellen fur wasser  
*J. Gen. Physiol.* 12:129-138
- 495 McShan, W.H., and Roland K. Meyer. 1945  
Augmentation of sheep pituitary gonadotropin by  
insoluble metallic hydroxides  
*Proc. Soc. Exp. Biol. Med.* 59:239-242
- 496 Medway, William, Morley B. Kare, and Eugene Kravis.  
1957  
The physiological effect of massive doses of  
mineral salts in pigs and chickens  
*Rep. N.Y. State Vet. Coll. Cornell Univ.*  
1955-56:22
- 497 Megnin, J. 1932  
Therapy of posttraumatic edema with intravenous  
injections of hypertonic solution of magnesium  
sulphate  
*J. Chir.* 40:193-213
- 498 Melton, Charles G., Jr. 1965  
Blastular arrest and partial cleavage of frog  
embryos injected with actinomycin D, mitomycin  
C, and homologous liver macromolecules  
*Dev. Biol.* 12(2):287-308
- 499 Mendelsohn, Simon. 1938  
Dispergierende mittel in backpulvern zwecks  
stabilisierung. II  
*Food Manuf.* 13:333-336
- 500 Mendenhall, Walter L., C.W. McClure, and Mildred  
Cate. 1926  
Die gallentreibenden eigenschaften des  
magnesiumsulfats  
*Boston Med. Surg. J.* 195:76-79
- 501 Merck and Co., Inc. 1966  
Antacid compositions  
*Neth. Pat.* 6,605,734 issued Nov. 21, 1966
- 502 Merritt, John A., and Jane F. Desforges. 1967  
Effect of magnesium on reticulocyte maturation  
*J. Lab. Clin. Med.* 70(3):452-457
- 503 Meyer, H., and H. Grund. 1963  
Absorption of magnesium by cattle  
*Tieraerztl. Umsch.* 18(4):181-182
- 504 Micelli, O., E. Avolio, and J. Leondeff. 1960  
Electrolytes in blood coagulation. I.  
Experimental methods employed  
*Boil. Soc. Ital. Biol. Sper.* 36:70-73
- 505 Mikhailov, V.A., and K.N. Klyachina. 1961  
Effect of manganese chloride and aminazine on  
thyroid function  
*Vopr. Gig., Fiziol. Tr., Prof. Patol. Prom.*  
*Toksikol., Sverd., Sb.* 6:361-367
- 506 Mikhailova, Lilyana. 1958  
Influence of magnesium chloride on the embryonal  
and post-embryonal development of some fish  
*Izv. Zool. Inst., Bulg. Akad. Nauk* 7:315-341
- 507 Miller, Aaron, and James K. Rocks. 1969  
Algin gel compositions  
*U.S. Pat.* 3,455,701 issued Jul. 15, 1969
- \* 508 Miller, E.R., D.E. Ullrey, C.I. Zutout, I.A.  
Hofer, and R.W. Luecke. 1965  
Mineral balance studies with the baby pig:  
Effects of dietary magnesium level upon calcium,  
phosphorus and magnesium balance  
*J. Nutr.* 86:209-212
- 509 Miller, J. Roscoe, and T.R. Van Dellen. 1938  
Electrocardiographic changes following  
intravenous administration of magnesium sulfate.  
An experimental study in dogs  
*J. Lab. Clin. Med.* 23:914-918
- 510 Miller, J. Roscoe, and T.R. Van Dellen. 1941  
Electrocardiographic changes following the  
intravenous administration of magnesium sulfate.  
III. Combined effect with digitalis  
*J. Lab. Clin. Med.* 26:1116-1120
- 511 Misumi, Kosei. 1925  
Studien uber den einfluss der CaCl<sub>2</sub>- und MgCl<sub>2</sub>-  
injektion auf die harnsekretion  
*J. Biochem.* 5:441-448

- 512 Mitolo, M. 1929  
Paralyzing effect  
Rell. Soc. Ital. Biol. Sper. 4:378-380
- 513 Miyahara, James T., Don W. Esplin, and Barbara Zablocka. 1966  
Differential effects of depressant drugs on presynaptic inhibition  
J. Pharmacol. Exp. Ther. 154(1):119-127
- 514 Miyao, Noboru, and Mitsuaki Hayashi. 1961  
Strontium metabolism following strontium-calcium discrimination in domestic animals. V. Effects of various salts and whole-body cobalt-60 gamma irradiation on the strontium-calcium discrimination in the goat and owl  
Nippon Juigaku Zasshi 23:181-191
- \* 515 Moineddin, J.F., and H.W. Lee. 1960  
Alimentary, blood and other changes due to feeding manganous sulfate, magnesium sulfate and sodium sulfate  
Am. J. Physiol. 199:77-83
- 516 Molumphy, Paul E., and Raphael Garcia. 1959  
Treatment of severe toxemia of pregnancy with intravenous magnesium sulfate and apresoline  
Obstet. Gynecol. 14(2):193-197
- \* 517 Monacelli, B., H. Tanaka, and J.H. Yoe. 1956  
Spectrochemical determination of magnesium, chromium, nickel, copper, and zinc in human plasma  
Clin. Chim. Acta 1:577-582
- 518 Monif, G.R. 1972  
Iatrogenic maternal hypocalcemia following magnesium sulfate therapy  
J. Am. Med. Assoc. 219:1469-1470
- 519 Monsanto Co. 1970  
Granular food acids  
Jap. Pat. 933/70 issued in 1970
- 520 Mori, Y. 1969  
Starch noodles  
Jap. Pat. 17,976/69 issued in 1969
- 521 Morita, Genpei. No date  
Über den einfluss der pharmaca auf die reizleitung zwischen vorhof und kammer  
Folia Pharmacol. Jap. 4:246-257
- 522 Morita, Sachikado. 1922  
Die hyperglykämie und glykosurie durch ammoniumchlorid und magnesiumsulfat bei der katze  
Tohoku J. Exp. Med. 3:363-413
- 523 Morris, Eugene Ray, and Boyd L. O'Dell. 1969  
Effect of magnesium deficiency in guinea pigs on kidney function and plasma ultrafiltrable ions  
Proc. Soc. Exp. Biol. Med. 132(1):105-110
- 524 Mortissey, Edmund J. 1925  
Der einfluss von magnesiumsulfat auf den druck der cerebrospinalflüssigkeit und das hirnvolumen  
Arch. Surg. 11:778-789
- 525 Motomura, Isao. 1938  
Effect of some salt solutions on the parthenogenetic membrane formation of sea-urchin eggs  
Sci. Rep. Tohoku Imp. Univ., 4th Ser. 13(1):85-88
- 526 Mozes, Magda, and Gy. Porika. 1961  
Action of electrolytes on thermoregulation of rats  
Rev. Med. (Targu Mures) 7:422-424
- 527 Mukoyama, Y. No date  
Über die wirkung der ionen Ca, K und Mg auf den sekretionsmechanismus der magendrüsen  
Biochem. Z. 157:303-332
- 528 Munck, O. 1954  
On the pathogenesis of the dumping-syndrome  
Acta Med. Scand. 148(5):329-342
- 529 Myhrman, G. 1930  
Effects, modification of tissue oxidation with chlorides of sodium, potassium, magnesium and calcium  
Acta Paediatr. 10:140-147
- 530 Nadasdi, M. 1961  
Influence of electrolytes on the cardiovascular effects of unilateral renal ischemia in rats  
Brit. J. Exp. Pathol. 42:324-330
- 531 Nagase, Ichiro, Mitsu Eguchi, and Shizuko Tanabe. 1962  
Fundamental studies on a stomach model. II. Digestion of starch and casein in the new stomach model  
Yakuzaijaku 22:80-82
- 532 Nagnibeda, N.I. 1957  
The treatment of advanced malignant tumours with magnesium sulphate  
Vopr. Onkol. 3(5):563-567
- 533 Nakazawa, Masumi. 1937  
The influence of diuretics on basal metabolism of rabbits  
Keio Igaku 17:1593-1646
- 534 Nanaeva, M.T., and I.A. Rakhinova. 1968  
Effect of papaverine, dibazol, and magnesium sulfate at high altitudes  
Mater. Sim. "Vysokogor'e Krasnaya Krov" Konf. "Vysokogor'e Lek." 1968:155-156
- 535 Nash, C.W., T. Tu, and H.J. Martin. 1972  
The influence of inorganic ions on the uptake and retention of titrated noradrenaline by isolated perfused rat hearts  
Can. J. Phys. Pharm. 50:645-654
- 536 Natale, P. 1932  
Magnesium sulphate: 43 cases  
Ann. Ostet. Gynec. 54:513-538
- \* 537 Neal, John B., and Marybelle Neal. 1962  
Effect of hard water and MgSO4 on rabbit atherosclerosis  
Arch. Pathol. 73:400-403
- 538 Neauport, C. 1971  
Estimation of Mg++ levels in the cellular fractions of tadpoles of Discoglossus pictus (Oth): influence of MgCl2 concentration in the breeding media. Study at several stages of development (Fr.)  
Arch. Sci. Physiol. (Paris) 25:361-375
- 539 Nichita, G., and I. Pahomeanu-Mesaros. 1945  
Effect of intravenous injections in dogs; clinical study  
Bull. Acad. Med. Roum. 10:87-93
- 540 Nichita, G., and M. Stanesco. 1945  
Effect on salivary, biliary and renal secretions of dog  
Bull. Acad. Med. Roum. 10:116-125
- 541 Nichita, G., I. Popesco, and I. Pahomeanu-Mesaros. 1945  
Effect of intravenous injections in dog; graphic study  
Bull. Acad. Med. Roum. 10:126-132
- 542 Nichita, H., and G. Nichita. 1945  
Effect of basal metabolism of dogs under normal conditions and during chloralose anesthesia  
Bull. Acad. Med. Roum. 10:236-245
- 543 Nichita, H., and G. Nichita. 1945  
Effect on basal metabolism of dog  
Bull. Acad. Med. Roum. 10:228-235
- 544 Nikiforov, M.I. 1967  
O farmakologicheskoy aspekte magnezial'noy narkoza. (The pharmacological aspect of magnesium anesthesia)  
Ref. Zh. Otd. Vyp. Farmakol. Khimioter. Sredstva Toksikol. 7(54):113



- 545 Nikiforov, M.I., and A.G. Gorodnik. 1969  
Influence of magnesium ions on the bioelectric activity of the brain against the background of armine (physostigmine)-induced activation (Russ.)  
Farmakol. Toksikol. (Moscow) 32(5):519-521
- 546 Nikolajev, O. 1929  
Die rolle der ionen und elektrolyte im prozesse der speichelsekretion  
Pflüger's Arch. Ges. Physiol. 223(1/2):95-103
- 547 Nishina, T. 1925  
Über die intravenöse injektion von magn. sulf. und die gallensekretion  
Nippon Naika Gakkai Zasshi 13
- 548 Noguer-More, J. 1932  
Effects, cytophyllactic effects of halogen salts; role of magnesium deficiency in human pathology; 3 cases  
Ars. Med., Barc. 8:419-421
- 549 Nolting, David F., and Omar M. Althabe. 1943  
El tratamiento de la hipertension arterial, en el embarazo, por el sulfato de magnesia  
Rev. Asoc. Med. Argent. 57(509/510):245-247
- 550 Northup, David W., and Edward J. Van Liere. 1939  
The effect of anoxemia on the absorption of magnesium sulfate from the small intestine  
Arch. Int. Pharmacodyn. 62:175-178
- 551 Oda, Yoshinobu. 1927  
Über den einfluss verschiedener substanzen auf den insulinblutzuckerspiegel. Leichtmetallsalze, alkalioide und alt-tuberkulin. (Effect of salts of light metals, alkaloids, and old tuberculin on the insulin-blood-sugar level) (Jap.)  
Folia Endocrinol. Jap. 3:630-657
- 552 Okamura, Kazuhiro. 1960  
Action of phosphates in Kawaboko and fish-sausage products. VII. Influence of added salts on the water intake of fish meat  
Nippon Suisan Gakkaishi 26:60-65
- 553 Okamura, Kazuhiro, Mituko Kodama, Hiroko Sonoda, and Jyunko Yamada. 1966  
Kawaboko and fish sausage products. XIX. Influence of added salt on the water intake and the protein solubility of water-washed raw fish flesh  
Nippon Suisan Gakkaishi 32(1):80-88
- 554 Okamura, Teiji. 1929  
Über den einfluss der verschiedenen vegetativen nervengifte, alkalioide und leichtmetallsalze auf die gallensaureausscheidung  
Arb. Med. Univ. Okayama 1:400-412
- \* 555 Oklander, Morris, and Bernard Klein. 1966  
Automated fluorometric determination of serum magnesium  
Clin. Chem. 12(4):243-247
- 556 Olin, T.E. 1936  
Über die desinfizierende bzw. antiseptische wirkung einiger chemischer verbindungen in trockenem oder leicht feuchten zustand  
Zentralbl. Bakt. Parasitenkd. Infektionskr. Abt. I. 137:283-287
- 557 Omel'chenko, O.T. 1957  
Effect of black radish and beet juice and of magnesium sulfate on bile secretion in patients with common bile duct drainage  
Fiziol. Zh., Akad. Nauk Ukr. R.S.S.R. 3(4):45-51
- 558 Onodera, N., Y. Nishina, K. Yukawa, S. Kata, and S. Faura. 1932  
Über eine die peristaltik anregende substanz in der magenschleimhaut  
Arch. Verdauungs-Drankh. 51:197-207
- 559 Opitz, K., G. Bornemann, and A. Loesser. 1961  
Possibility of influencing the absorption of magnesium  
Arzneimittel-Forschung 11:1066-1067
- 560 Oppenheimer, Morton J., and Frank C. Mann. 1941  
Influence of cathartics on the activity of small intestine  
Am. J. Dig. Dis. 8:90-93
- 561 Organon, N.V. 1971  
Method for producing a sweetener composition  
Meth. Pat. 6,918,568 issued in 1971
- 562 Orru, Antonietta. 1940  
Sul comportamento del tuorlo di uovo di gallina immerso in soluzioni variamente concentrate di diversi elettroliti. (The behavior of the yolk of hen's egg immersed in different concentrations of various electrolytes)  
Arch. Sci. Biol. (Bologna) 26:593-610
- 563 Overbeck, Henry W., Robert M. Daugherty, Jr., and Francis J. Haddy. 1969  
Continuous infusion indicator dilution measurement of limb blood flow and vascular response to magnesium sulfate in normotensive and hypertensive men  
J. Clin. Invest. 48(10):1944-1956
- 564 Owles, W.H. 1937  
Investigations of the functions of the small intestine in men by intestinal intubation. I. Technic of intestinal intubation in man  
Clin. Sci. 3:1-10
- \* 565 Özsoylu, S. 1964  
Effect of magnesium sulfate on blood sugar levels of diabetic children  
Türk. J. Pediatr. 6:211-217
- 566 Pacaud, A. 1933  
Action de la lecithine et du chlorure de magnesium sur la vie et la reproduction des Cladoceres  
Compt. Rend. Acad. Sci. (Paris) 197(2):194-196
- 567 Packett, L.V., and J.P. Hauschild. 1964  
Phosphorus, calcium, and magnesium relations in ovine urolithiasis  
J. Nutr. 84(2):185-190
- \* 568 Paixao, L.M., and J.H. Yoe. 1959  
Spectrochemical determination of magnesium, chromium, nickel, copper, and zinc in human plasma and red cells  
Clin. Chim. Acta 4:507-514
- 569 Palmer, L.S., C.H. Eckles, and D.J. Schutte. 1928  
As factor in retention of calcium and phosphorus in cattle  
Proc. Soc. Exp. Biol. Med. 26:58-62
- 570 Pandazi, Andrew A. 1964  
The mode of action of magnesium sulfate in the treatment of toxemia of pregnancy  
Marquette Med. Rec. 30(1):24-27
- 571 Parfentjev, I.A., A. Waldschmidt, and A.J. Weil. 1942  
Absorption of bacterial antigen from diphtheric toxin and toxoid by magnesium hydroxide  
J. Immunol. 44:325-328
- 572 Parker, C.B. 1969  
Insure keeping quality, texture of tomato pack with 3 way salt tablet  
Food Proc. 30(5):30-31
- 573 Patton, Alva R. 1948  
Processes for controlling potato chip color  
U.S. Pat. 2,448,152 issued Aug. 31, 1948
- 574 Paul H. Raymer, East Orange, N.J. 1927  
Kaugummi  
A. Pat. 1,630,763 issued May 31, 1927
- 575 Paul, F. 1971  
Effect of oral Ca and Mg on gastric secretion of HCl Munch. Med. Wochenschr. 113:422-425

- 576 Paulus, W. 1962  
Der magnesiumgehalt im leichenurin nach aufnahme grosserer mengen von magnesiumsulfat. (Magnesium content of cadaver urine after taking large amounts of magnesium sulfate)  
Arch. Toxikol. 19(5):325-326
- 577 Pelikan, Zdenek. 1969  
Activity of pancreatic lipase influenced by antituberculous in the presence of ions,  $Hg^{++}$  and  $Mn^{++}$ , in vitro  
Scr. Med. 42(6-7):351-354
- 578 Pentschew, A., and H. Kassowitz. 1932  
Vergleichende untersuchungen über die wirkung verschiedener metallsalze auf das zentralnervensystem von kaninchen  
Arch. Exp. Pathol. Pharmacol. 164:667-684
- 579 Penzoldt, Franz. 1923  
Über alte und neue abfuhrmittel  
Munch. Med. Wchschr. 71:33-35
- 580 Perrin, Maurice, and Alain Cuenot. 1930  
Versuch mit einigen anagotischen salzen  
Compt. Rend. Soc. Biol. 102:1038-1039
- 581 Petersen, Carl. 1931  
Über den calcium- und phosphorstoffwechsel der schweine. III. Der einfluss verschiedener mineralzulagen auf den saure-basen-haushalt bei schweinen und insbesondere auf die acidität und den ammoniakgehalt des harns  
Wiss. Arch. Landwirtsch. Abt. B. Arch. Tierernähr. Tierzucht. 5(4):567-597
- 582 Petrovskaja, B. 1939  
The action of drugs on the isolated perfused lungs of the pig  
Q. J. Exp. Physiol. 29:277-282
- 583 Petrovskii, Yu.A. 1936  
The mechanism of the action of a series of medicinal agents on the biliary tract  
J. Physiol. (USSR) 21:293-310
- 584 Pfeiffer, E. 1952  
Magnesium malcosis in carp  
Maugnyn-Schmiedeberg's Arch. Exp. Path. Pharmacol. 216:375-380
- 585 Pfeiffer, Carl, Howard G. Glass, and Robert H. Dreisbach. 1941  
Effect of varying intake of dietary salts on formation of galactose cataracts  
J. Pharmacol. 72:31-32
- 586 Pines, N. 1933  
Magnesium sulphate in treatment of angiospasm  
Lancet 1:577-579
- 587 Pinilla, Casto. 1942  
Intravenous or intraspinal injections of magnesium sulfate in therapy, especially of Sydenham's chorea  
Prensa Med., La Paz 2:254-257
- 588 Pjatnitsky, N., and P. Olefirenko. 1927  
Einfluss von intravenösen  $MgCl_2$ - und  $CaCl_2$ -injektionen auf die durch morphin erzeugte magensaltsekretion (Russ.)  
J. Exp. Biol. Med. 1927:382-397
- 589 Plante, Norwand. 1961  
Le sulfate de magnesie dans de sevrage alcoolique et ses complications. (Magnesium sulfate in alcohol withdrawal and its complications)  
Laval Med. 31(2):147-155
- 590 Pleshchitsky, A.Ya. 1938  
The velocity of blood circulation in the lesser circulatory system of rabbits after parenteral injection of magnesium sulfate (Ger.)  
J. Physiol. (USSR) 25:150-153
- 591 Plotnikoff, N., and A.V. Evans. 1967  
Enhancement of conditioned photic evoked responses in the rabbit by penoline and magnesium hydroxide  
Int. J. Neuropsychiatr. 3(3):263-267
- 592 Plotnikoff, Nicholas. 1970  
Comparison of PMH (penoline and magnesium hydroxide) and penoline activity after electroconvulsive shock  
Arch. Int. Pharmacodyn. Ther. 184(1):175-185
- 593 Plotnikoff, Nicholas, F. Will, and W. Ditzler. 1969  
Stimulant activity of penoline and magnesium hydroxide (magnesium penoline, PMH)  
Arch. Int. Pharmacodyn. Ther. 181(2):441-458
- 594 Polukhin, P.S., and A.G. Savoiskii. 1962  
Changes in carbohydrate and fat metabolism in productive cows fed on trace element rations (Russ.)  
Tr. Mosk. Vet. Akad. 44:116-123
- 595 Popoff, M. 1942  
The effect of chemical stimulation on the growth of mouse tumors  
Z. Krebsforsch. 52:32-36
- 596 Popoff, Methodi, Minko Dobreff, and Georgi Paspaletti. 1926  
Beeinflussung der avitaminose bei tauben durch zellstimulation  
Zellstimulationsforsch 2:47-57
- 597 Popova, E.A. 1969  
Assessment of the cholagogic effect of sunflower seed oil and magnesium sulfate on the basis of the urochecystokinin level  
Sov. Med. 32:57-62
- 598 Pora, Eugen A., and Florica Stoicovici. 1964  
Dynamics of blood cations as a function of the salt balance in rabbits  
Stud. Cercet. Biol., Ser. Zool. 16(4):307-318
- 599 Prandi, F., and J. Jimenez-Vargas. 1954  
Effect of different salts on the contractility of bronchial muscle  
Rev. Esp. Fisiol. 10(3):159-177
- 600 Presl, J. 1970  
Magnesium sulfate in the therapy of pre-eclampsia and eclampsia  
Cesk. Gynecol. 35:193-194
- 601 Pribyl, E. 1927  
Farmakoterapeuticky ucinek koloidniho hydroxydu horecnateho na zanety jater a ledvin toxikogenneho puvodu. (The pharmacotherapeutic action of colloidal magnesium hydroxide in experimental hepatitis and nephritis of toxic origin)  
Biol. Spisy. Vys. Skoly Zverolek. Brno 6:291-318
- 602 Pribyl, E. 1929  
Effects of magnesium ion on metabolism of calcium in organism  
Compt. Rend. Soc. Biol. 102:258-260
- 603 Pribyl, E. 1931  
Therapeutic use of colloidal magnesium hydroxide in inflammations produced experimentally by toxic drugs  
Arch. Exp. Pathol. Pharmacol. 160:255-268
- 604 Prodanski, P. 1968  
The effects of some salts on the rennetability of various milks  
Milchwissenschaft 23(8):491-497
- 605 Puetzer, Bruno, Leonard Mackles, and Martin M. Perl (to Tintex Corp.). 1967  
Compositions for relief of gas-caused gastrointestinal distress in warm-blooded animals  
U.S. Pat. 3,355,354 issued Nov. 28, 1967
- 606 Quaranta, L. 1928  
Effect of parenteral administration on vegetative nervous system  
Studia 18:439-442

- 607 Raab, Ernst. No date  
Über die Bedeutung anorganischer Salze für das  
Kontraktionsvermögen vegetativer Muskeln  
Pflügers Arch. Physiol. 215:651-666
- 608 Raciszewski, Z., and E. Pijanowski. 1961  
Improvement of the quality of baking powders by  
means of magnesium stearate  
Zesz. Nauk. Szk. Gł. Gospod. Wiejsk. Warsz., Roln.  
4:239-245
- 609 Rademaekers, A., and Torald Sollmann. 1924  
Untersuchungen über salinische Abführmittel. I.  
Magnesiumsulfat am herausgenommenen  
Darm: minimalkonzentrationen in lockerscher  
Lösung; ausserlich beigebracht; magnus' methode  
Arch. Int. Pharmacodyn. Ther. 29:481-487
- \* 610 Radway, Jerrold R. 1966  
Magnesium oxide in the raw sugar house  
Jaw. Assoc. Sugar Technol. J. 1966(27):48-58
- 611 Kaida, Hanns. 1923  
Kalkbewegung durch Neutralsalze  
Z. Ges. Exp. Med. 37:266-273
- 612 Rakhimova, I.A. 1969  
Vliyaniye dibazola, papaverina i sulfata magniya  
na krovoobrashchenie i dykhanie sobak v  
usloviyakh vysokogor'ya. (Effect of dibazol,  
papaverine and magnesium sulfate on  
blood circulation and respiration in dogs under  
high-altitude conditions)  
Farmakol. Toksikol. 32(2):287-289
- 613 Rakusin, M. No date  
Magnesiumhydroxyd als Gegengift gegen sublimat  
J. Exp. Biol. Med. (Russia) 4:52-54
- 614 Rakusin, M. 1927  
Magnesiumhydroxyd als Antidotum arsenici  
Munch. Med. Wchschr. 74:1667-1668
- 615 Rakusin, M.A. No date  
Magnesiumhydroxyd als Gegengift gegen sublimat  
Munch. Med. Wchschr. 74:325-326
- \* 616 Randall, R.L., M.D. Cohen, C.S. Spray, Jr., and  
E.C. Rossmelisl. 1964  
Hypermagnesemia in renal failure: Etiology and  
toxic manifestations  
Ann. Intern. Med. 61:73-88
- 617 Rangan, C.H., and J.C. Gupta. 1962  
Effect of parenteral magnesium on serum  
cholesterol in rabbits  
Indian J. Med. Res. 50:18-22
- 618 Ranzi, S., B. Arosio, P. Citerio, P. Menotti, and  
F. Semenza. 1946  
The mechanism of action of substances which  
modify embryonal development (Ital.)  
Experientia 2:315-316
- 619 Reach, Felix. No date  
Die Beeinflussung des Choleodochus-sphincter durch  
Magnesiumsulfat  
Arch. Exp. Path. Pharmacol. 100:379-384
- \* 620 Rees, P.O. 1970  
A case of poisoning by Epsom salt  
J. Forensic Sci. Soc. 10:91-94
- 621 Reznikoff, Paul. 1928  
Mikrographische Studien über Zellphysiologie. V. Der  
Antagonismus der Kationen bei ihrer Wirkung auf  
das Protoplasma von Amoeba dubia  
J. Gen. Physiol. 11:221-232
- 622 Rhodes, Goodrich B. 1925  
Mit Magnesiumhydroxyd behandelter Dorschlebertran  
bei der Behandlung chirurgischer tuberkulose  
J. Lab. Clin. Med. 11:227-230
- 623 Rice, E.W., and C.Z. Lapara. 1964  
Rapid ultramicrospectrophotometric determination  
of magnesium  
Clin. Chim. Acta 10:360-364
- 624 Richardson, Alan E. 1965  
Clinical (and pharmacotherapeutic) aspects of  
cerebral edema  
Proc. Royal Soc. Med. 58(8):604-606
- 625 Kriessner, Otto, and Keniti Yamada. 1933  
Über die Beeinflussung des Muskelchemismus durch  
intravenöse Infusion von vegetativen Giften und  
Narkotika  
Naunyn-Schmiedeberg's Arch. Exp. Path. Pharmacol.  
170:208-225
- 626 Rigo, Janos, and Istvan Szelenyi. 1967  
Effect of magnesium on the fat-induced increased  
coagulability (Hung.)  
Kiserl. Orvostud. 19(1):17-20
- 627 Rising, J.D., and M.C. Carroll. 1946  
Comparative value of pentothal sodium (barbital  
derivative), curare, and magnesium sulfate for  
modification of metrazol convulsions  
J. Kans. Med. Soc. 47:297-299
- 628 Ritter, U. 1955  
Bewegungsmechanismen der Papilla Vateri.  
(Mechanisms of motility of the papilla of Vater)  
Z. Ges. Exp. Med. 126(5):444-449
- 629 Robin, P., and P. Bourdin. 1966  
Action of sodium sulfate, magnesium sulfate, and  
magnesium chloride on rinderpest virus adapted  
to cell culture (Fr.)  
Rev. Elev. Med. Vet. Pays Trop. 19(4):451-456
- 630 Roche, Jean, Nguyen-Van-Thoai, and Abderrhaman  
Dziri. 1945  
Action on amylases of amino acids associated with  
metallic ions  
Compt. Rend. Soc. Biol. 139:519-521
- 631 Roffo, A.H. 1925  
Die Wirkung gewisser metallischer Ionen auf die  
Entwicklung normaler und neoplastischer Gewebe  
in vitro  
Bol. Inst. Med. Exp. 1:307-323
- 632 Rogers, K.T. 1956  
Re-examination of the production of cyclopia in  
Fundulus heteroclitus with magnesium chloride  
and ethyl alcohol  
Biol. Bull. 110(3):344-351
- 633 Rogers, T.A., F.L. Haven, and P.E. Mahan. 1960  
Exchange of radioactive magnesium in Walker  
carcinoma 256: A note 1,2  
J. Natl. Cancer Inst. 25:887-888
- 634 Roine, Paavo, and C.A. Elvehjem. 1950  
Significance of the intestinal flora in nutrition  
of the guinea pig  
Proc. Soc. Exp. Biol. Med. 73(2):308-310
- 635 Rokhlina, M.L. 1929  
The role of ions and hormones in the excitation  
process of effectors  
J. Biol. Med. Exp. 13(35/36):108-120
- \* 636 Roller, Dietrich. 1936  
A case of magnesium sulfate poisoning  
Wien. Klin. Wchschr. 49:241-242
- 637 Rona, P., and H. Kleinmann. 1927  
Nephelometrische Untersuchungen über fermentative  
Eiweisspaltung. VII. Mitt. Der Einfluss von  
Ionen auf die Stabilität des Trypsins  
Biochem. Z. 196:177-196
- 638 Rondeau, Edward. 1937  
Action of salts on Amoeba proteus  
Mendel Bull. 9:119-121
- 639 Rosamund, Eugene. 1937  
Prontosil  
J. Tenn. State Med. Assoc. 30:200-202
- 640 Rosen, H., J. McCallum, A. Blumenthal, and H.P.K.  
Agersborg. 1964  
Effects of K and Mg aspartates on the isolated  
guinea pig heart  
Arch. Int. Pharmacodyn. 147(3-4):476-481

- \* 641 Rosier, O.A. 1952  
Old and new cures of tapeworm infection; acute poisoning due to oral administration  
Wien. Klin. Wchnschr. 64:942-945
- 642 Rossi, Emil R., and Aldo A. Rainero, Jr. 1967  
Sugar block with handle  
U.S. 3,312,555 issued Apr. 4, 1967
- 643 Roth, George B., and Phoebe J. Crittenden. 1934  
Sulphates of sodium and magnesium on gastrointestinal activity  
Proc. Soc. Exp. Biol. Med. 32(1):91-94
- 644 Rothberger, C.J., and L. Zwillinger. 1936  
Über die Wirkung von Magnesium auf die Strophanthin- und die Bariumtachykardie  
Münch.-Schwabebergs Arch. Exp. Path. Pharmacol. 181:301-316
- 645 Roufogalis, B.D., and J. Thomas. 1968  
The acceleration of acetylcholinesterase activity at low ionic strength by organic and inorganic cations  
Mol. Pharmacol. 4(2):181-186
- 646 Rowland, Robert C., and Jack A. Pritchard. 1964  
The effect of parenteral magnesium sulfate on blood loss at delivery  
Am. J. Obstet. Gynecol. 89(2):261-263
- 647 Royer, M., and P.M. Manfredi. 1944  
Pressure within the common bile duct of the normal dog  
Rev. Soc. Argent. Biol. 20:232-245
- 648 Rozen, J.S., and G.L. Perusse, Jr. 1929  
Effect of calcium and magnesium ions on gastric hunger contractions  
Am. J. Physiol. 91:298-304
- 649 Rozental, D.L. 1948  
Contraction of transversely striped muscle caused by calcium, barium, and magnesium chlorides  
Dokl. Akad. Nauk. S.S.S.R. 63:593-596
- 650 Rubentschik, L. 1924  
Über die einwirkung von salzen auf die lebensfähigkeit der urobakterien  
Zentralbl. Bakt. Parasitenkd. Abt. II 67:167-194
- 651 Rubin, Mitchell I., and Milton Rapoport. 1941  
The mode of action of magnesium sulfate in reducing the hypertension of acute glomerulonephritis  
Am. J. Med. Sci. 201:734-745
- 652 Runnstrom, John. 1928  
Struktur und atmung bei der entwicklungs-erregung des seeigels  
Acta Zool. (Stockholm) 9(3):445-492
- 653 Russetzky, I.I. 1928  
Effect of intravenous injections on muscular tonus in parkinsonism  
Med. Biol. J. 4(3-4):84-88
- 654 Rutishauser, E., and M. Bianchi. 1938  
Experimentelle studie über die veränderungen der serumphosphatase bei metallvergiftungen  
Arch. Int. Pharmacodyn. Ther. 58:240-247
- 655 Ryan, William J., and Lawrence Kogan. 1968  
Dusting starch, especially for bakery dough  
U.S. Pat. 3,377,171 issued Apr. 9, 1968
- 656 Sabalitschka, Th., and E. Pilger. 1938  
Einfluss des wassers auf den kaffeegeschmack  
Umsch. Wiss. Technol. 42:920-922
- 657 Sabrodski, A. 1935  
Der einfluss der wassersalze auf das malz beim prozess der malzextraktgewinnung  
Gar.-Ind. (Russ.-Brodilnaja Prom-st.) 12(1):22-26
- 658 Saegesser, Max. 1932  
Das magnesiumpulver in der behandlung des wundstarrkrampfes  
Schweizerische Med. Wochenschr. 62(13):306-309
- 659 Sahu, Gadhadhar, and Achintya K. Mukherjee. 1966  
Effect of magnesium on sugar, calcium, and magnesium levels in the blood of hypothyroid and thyroid-fed rabbits  
Indian J. Exp. Biol. 2(2):111-113
- 660 Saito, Chyutaro. 1933  
Über die veränderung der elektrischen erregbarkeit des herzvagus durch verschiedene mittel. II. Mitt. Über die wirkung der salze.  
Acta Sch. Med. Univ. Kioto 16:152-158
- 661 Sakhanova, L.V. 1966  
Ob izbiratel'nom deistvii sernokislologo magniya i nembutala na zritel'nyi, slukhovoi i kozhnyi analizatory. (Selective activity of magnesium sulfate and nembutal on the visual, auditory and skin analysors)  
Ref. Zh. Biol. 1966(1p):385
- 662 Sambuc, E., G. Raymond, and M. Maudet. 1971  
Oxidation of salted margarine  
Rev. Fr. Crops Gras 18(1):17-20
- 663 Sarudi, Iare. 1938  
Über die bestimmung der mineralbestandteile in pflanzlichen stoffen und nahrungsmitteln  
Osterr. Chem.-Ztg. 41:436-438
- 664 Saunders, J. 1969  
Verfahren zur herstellung von kochsalz. (Process for manufacturing cooking salt)  
W. Ger. Pat. 1,492,998 issued in 1969
- 665 Saunier, S., and A. Lemaitre. 1951  
Volumetric determination of calcium and magnesium in sugar-factory juices  
Sucr. Fr. 92:231-233
- 666 Sborgi, U., and A. Golanti. 1937  
Ricerche chimiche e chimicofisiche sulle acque minerali della perla (pisa)  
Ann. Chim. Appl. 27(5):213-232
- 667 Schiff, E. 1920  
Action of magnesium sulphate on calcium and magnesium metabolism in infants  
Jahrb. Kinderheilkd. 91:43
- 668 Schijvarger, L. 1943  
Glomerulotubular nephropathies (mixed nephritis) treated with magnesium sulfate; 4 cases  
Dia Med. 15:114-115
- 669 Schmalbruch, H. 1964  
Effects of alkali and alkaline earth ions (Na+, K+, Li+, Mg++, Ca++) on certain structures of the mouse diaphragm (Ger.)  
Z. Zellforsch. Mikroskop. Anat. 62(2):246-278
- 670 Schmidt, Carl F., and W. Benson Harer. 1923  
Die wirkung von arzneimitteln auf die atmung. I. Die morphinreihe; II. Ather, chloroform, chloral, urethan, luminal, magnesium, coffein, strychnin und atropin  
J. Exp. Med. 37:47-67; 69-81
- 671 Schmidt, G.R., R.G. Cassens, and E.J. Briskey. 1970  
Changes in tension and certain metabolites during the development of rigor mortis in selected red and white skeletal muscles  
J. Food Sci. 35(5):571-573
- 672 Schmidt, Hans. No date  
Die steigerung der antikörperbildung durch intravenöse manganchlorur-einspritzungen  
Zentralbl. Bakt. Parasitenkd. 95:74-81
- 673 Schmidt, S. 1925  
Metallsalze und antikörper  
Z. Immunitätsforsch. Exp. Ther. I. 45:305-322
- 674 Schnitzer, R.J., and F.L. Pundsack. 1970  
Asbestos hemolysis  
Environ. Res. 3(1):1-13

- 675 Schoonover, Janetta Wright. 1936  
Enzyme bei Krebs. II. Die glycerophosphatasen der menschlichen erythrocyten.  
Biochem. J. 30:1097
- 676 Schroeder, H.A. 1960  
Relations between hardness of water and death rates from certain chronic and degenerative diseases in the United States  
J. Chronic Dis. 12:586-591
- \* 677 Schroeder, H.A., A.P. Mason, and I.H. Tipton. 1969  
Essential metals in man: Magnesium  
J. Chronic Dis. 21:815-841
- 678 Schroeder, Hermann, and Bernard B. Raginsky. 1932  
Über die harnsaureausscheidung durch den darm und ihre pharmakologische beeinflussung  
Naunyn-Schmiedeberg's Arch. Exp. Path. Pharmacol. 168:413-423
- 679 Schuphan, W. 1953  
The experimental results of the bundesanstalt für qualitätsforschung pflanzlicher erzeugnisse on the question of the potassium diarrhea after spinach consumption  
Landwirtsch. Forsch. 4:154-161
- 680 Seiden, Rudolf (to Haver-Lockhart Laboratories, Inc.). 1966  
Minimizing weight loss in slaughter animals  
U.S. Pat. 3,241,974 issued Mar. 22, 1966
- \* 681 Selisko, O., and H. Ackermann. 1959  
Magnesium tetany and toxicity of various magnesium salts on intraperitoneal injection  
Naturwissenschaften 46:385-386
- \* 682 Selisko, O., and H. Ackermann. 1963  
On the mean lethal dose (L.D.50) of several magnesium salts in aqueous solution  
Naturwissenschaften 50(13):479-480
- 683 Sell, Jerry L. 1969  
Magnesium nutriture of the hen: influence on retention of magnesium, calcium, and nitrogen, and on ration metabolizable energy value  
Poult. Sci. 48(4):1437-1442
- 684 Selye, H., and G. Gabriani. 1965  
The role of electrolytes in the pathogenesis of experimental cardiopathies without vascular involvement  
Electrolytes Cardiovasc. Dis. 1:135-160
- 685 Selye, Hans, and Eors Bajusz. 1960  
The prevention of experimental myopathies by various chlorides  
J. Nutr. 72:37-45
- \* 686 Serbescu, P. 1934  
Influence of magnesium on experimental cancer  
Bull. Acad. Med. (Paris) 111(12):435-438
- 687 Shah, Yu Ting. 1927  
Changes in the oxygen consumption of kidney during saline diuresis  
Proc. Imp. Acad. 3(9):627-629
- 688 Shapiro, Leonard. 1967  
Rapid analysis of rocks and minerals by a single-solution method  
U.S. Geol. Surv. Prof. Pap. 575-B:187-191
- 689 Shay, Harry, S.A. Komarov, M. Siple, and Margot Gruenstein. 1947  
An evaluation of some antacid and anti-peptic agents in the prevention of gastric ulceration in the rat  
Am. J. Dig. Dis. 14(3):99-104
- 690 Shcherbakov, S.A., I.B. Bakhtemeev, I.B., and P.N. Andreev. 1936  
The oligodynamic action of salts on the enzymic activity of pepsin (Ger.)  
Arch. Fisiol. 36:315-322
- 691 Shchipitsina, L.P. 1967  
Uncoupling of oxidative phosphorylation in the mitochondria of rat heart and kidneys under conditions of vitamin B2 poisoning (Russ.)  
Vopr. Pitan. 26(2):38-42
- 692 Shimizu, Yutaka, and Wataru Simidu. 1953  
Studies on muscle of aquatic animals. XIV. Effect of neutral salts on protein denaturation (Jap.)  
Bull. Jap. Soc. Sci. Fish. 19(6):757-760
- 693 Shinagawa, Minoru. 1925  
Vergleichende untersuchung über die spontane bewegung und die wirkung der pharmaca an verschiedenen teilen des ausgeschnittenen kaninchenuterus  
Folia Jap. Pharmacol. 1:421-439
- 694 Shionoya, Takuji. 1927  
Studies in experimental extracorporeal thrombosis. V. Influence of certain chemical substances on extracorporeal thrombosis with special reference to the efficacy of a combination of heparin and magnesium sulfate  
J. Exp. Med. 46(6):963-977
- 695 Shitov, E.E. 1965  
Effect of magnesium sulfate associated with tranquilizing agents on the bio-electrical and cholinesterase activity of the brain  
Farmakol. Toksik. 28:13-17
- 696 Shoji, Kiyoshi. 1937  
Adrenaline content of the suprarenal glands of rabbits after administration of ammonium chloride and magnesium sulfate  
Tohoku J. Exp. Med. 30:259-265
- 697 Shrivastava, Dasharath Lal, Baghunath Sahai Gupta, Mata Prasad, and Shanti Swarupa Bhatnagar. No date  
Ein weiterer beweis für die chemische theorie der schutzwirkung der zucker  
J. Phys. Chem. 29:166-177
- 698 Shvedenko, L.A. 1965  
Some biochemical indexes in patients with hypertension  
Vrach. Delo 1965(12):105-106
- 699 Shvedsky, B. 1930  
Coagulation of pure solution of fibrinogen and plasma combined with magnesium sulphate; methods of preparation  
J. Eksp. Biol. Med. (Ser. A., No. 39) 13:53-57
- 700 Shytov, Ye. Ye. 1963  
Vplyv sirchanokysloyi mahneziyi, pokhidnykh dytenilotstovoyi kysloty ta aminazynu na dykhannya tkany mozku bilykh shuriv. (The effect of magnesium sulfate, diphenylacetic acid derivatives, and chlorpromazine on the respiration of brain tissue in white rats)  
Sb. Nauk Prats' L'vivskiy Med. Inst. 24:94-97
- \* 701 Silver, L., J.S. Robertson, and L.K. Dahl. 1960  
Magnesium turnover in the human studied with Mg28  
J. Clin. Invest. 39:420-425
- 702 Simici, D., M. Popescu, and C. Covaceanu. 1935  
Recherches faites en Roumanie sur la fréquence de l'infection du duodenum et des voies biliaires par les lamblies intestinalis  
Bull. Mem. Soc. Med. Hôp. Bucarest 17(9):154-160
- 703 Simon, Alexander. No date  
Über die wirkung der verschiedenen konzentrierten kationenlösungen auf die osmotische resistenz der roten blutkörperchen  
Biochem. Z. 170:244-253
- 704 Simon, Hans. No date  
Über die wirkung von karlsbader-, glauher- und bittersalz auf die aussere sekretion von leber und bauchspeicheldrüse  
Dtsch. Med. Wchschr. 50:1012-1014

- 705 Simpson, J.I., I.C.L. Chang, E.C. Appel, and M.C. Bollman. 1955  
Effect of water hardness in reconstituting dehydrated fruits and vegetables  
Food Technol. 9(12):613-615
- 706 Singh, Chanan, and C.R. Krishna Murti. 1967  
Influence of polyanions and polycationic complexes of metals on cells and cellular organelles  
Indian J. Biochem. 4(3):171-174
- 707 Sinha, Bindesri Prosad. 1950  
Effect of magnesium chloride on gastric secretion  
Antiseptic (Madras) 47:890-891
- 708 Skala-Rosenbaum, J. 1928  
Acidosis and magnesium hydroxide  
Cas. Lek. Cesk. 67:1437-1438
- 709 Sky-Peck, H.H. 1964  
A method for determination of magnesium in serum and urine  
Clin. Chem. 10(5):391-398
- 710 Sloan, J.H. 1926  
Über den Mechanismus der Vorbeugung einer tödlichen parathyreoideen tetanie durch perorale Zuführung von Magnesiumchlorid  
Am. J. Physiol. 79:100-111
- 711 Smith, E.A., S. Alfend, and L.C. Mitchell. 1926  
Detection of added pepper shells in pepper  
J. Assoc. Offic. Agric. Chem. 9(3):333-342
- \* 712 Smith, P.K., A.W. Winkler, and B.M. Schwartz. 1939  
Distribution of magnesium following parenteral administration  
J. Biol. Chem. 129:51-56
- 713 Smith, R.B., and F.M. Giesy. 1923  
The alkalinity of magma magnesiae as determined by the hydrogen electrode  
J. Am. Pharm. Assoc. 12:955-957
- 714 Smith, R.B., and F.M. Giesy. 1924  
Die Alkalität von Magnesiaöl, bestimmt durch die Wasserstoffelektrode  
J. Am. Pharm. Assoc. 13:1118-1120
- 715 Sobel, Albert Edward, and Albert Hanok. 1952  
Calcification. VII. Reversible inactivation of calcification in vitro and related studies  
J. Biol. Chem. 197(2):669-685
- 716 Societe Industrielle pour la Fabrication des Antibiotiques. 1966  
Dietetic compositions  
Fr. Pat. M3789 issued Jan. 31, 1966
- 717 Soffer, Louis J., Clarence Cohn, Gerson Lesnick, Harry Sobotka, and Mildred Jacobs. 1944  
Effect of sodium iodide, magnesium sulfate, thyroxin, and thyrotropic hormone on the blood magnesium partition  
J. Clin. Invest. 23(2):263-266
- 718 Sollmann, Torald, and A. Rademaekers. 1926  
Investigations on saline cathartics. II. Magnesium sulphate on the peristalsis and the propulsion in small intestines  
Arch. Int. Pharmacodyn. Ther. 31:39-69
- 719 Sonjen, G.G., and E.W. Baskerville. 1968  
Effect of excess magnesium on vagal inhibition and acetylcholine sensitivity of the mammalian heart in situ and in vitro  
Nature 217(5129):679-680
- 720 Sotskii, O.P. 1969  
Effect of gangliosides on the process of swelling and contraction of rat brain mitochondria (Russ.)  
Vopr. Biokhim. Mozga 1969(15):139-146
- 721 Soudan, P. 1955  
Chemical aspects of the salting of codfish  
Rev. Trav. Inst. Sci. Tech. Peches Marit. 19:129-136
- \* 722 Spare, P.D. 1962  
A study of the titan yellow dye lake methods for the estimation of serum magnesium  
Am. J. Clin. Path. 37(2):232-236
- 723 Spek, Josef. 1924  
Studien an zerschnittenen Zellen  
Arch. Entw.-Mech. 101:451
- 724 Spencer, H., I. Levin, and J. Samachson. 1967  
Effect of magnesium on radiostrontium excretion in man  
Int. J. Appl. Radiat. Isot. 18(6):407-415
- \* 725 Speranskaja-Stepanova, E.M. 1933  
Influence on excretion of urine by dogs  
Z. Ges. Exp. Med. 88:639-641
- 726 Speranskaja-Stepanova, E.M. 1934  
Über die Wirkung von Morphin und schwefelsaurem Magnesium auf die Diurese bei hunden  
Tr. Vsessojuznogo Inst. Exp. Med. 1(1):163-171
- 727 Spira, Gaston. 1964  
Food preservation  
Fr. Pat. 1,380,648 issued Dec. 4, 1964
- 728 Stander, H.J. No date  
Wirkung von intravenöser Zufuhr von Magnesiumsulfat. Experimentelle Untersuchung  
J. Am. Med. Assoc. 92:631-636
- 729 Stander, H.J. 1928  
Effect of intravenous administration; experimental studies  
Tr. Sect. Obstet., Gynecol. Abd. Surg., Am. Med. Assoc. pp. 210-224
- 730 Steele, T.H., S.-P. Wen, M.A. Evenson, and E.E. Fieselbach. 1968  
The contribution of the chronically diseased kidney to magnesium homeostasis in man  
J. Lab. Clin. Med. 71:455-463
- 731 Steigmann, F., and Arthur R. Marks. 1944  
Inhibition of peptic activity in the treatment of peptic ulcer  
Am. J. Dig. Dis. 11:173-178
- 732 Stein, Herman H., and Tobias O. Yellin. 1967  
Penicillin and magnesium hydroxide: Lack of effect on RNA and protein synthesis  
Science 157(3784):96-97
- 733 Steinkraus, Keith H., and Roger A. Morse. 1966  
Factors influencing the fermentation of honey in mead production  
J. Apic. Res. 5(1):17-26
- 734 Stern, Philip A., and Raymond L. Cahen. 1956  
Evaluation of the laxative potency of saline cathartics and aperients  
Arch. Int. Pharmacodyn. 107:178-193  
Arch. Intern. Pharmacodyn. 107:178-193
- \* 735 Stevens, Alexander R., and Harold G. Wolff. 1950  
Magnesium intoxication. Absorption from the intact gastrointestinal tract  
Arch. Neurol. Psychiatr. 63:749-759
- \* 736 Stone, Stuart R., and Jack A. Pritchard. 1970  
Effect of maternally administered magnesium sulfate on the neonate  
Obstet. Gynecol. 35(4):574-577
- 737 Storry, J.E., and J.A.F. Rook. 1963  
Magnesium metabolism in the dairy cow. V. Experimental observations with a purified diet low in magnesium  
J. Agric. Sci. 61:167-171
- 738 Stuber, Bernhard, and Minoru Sano. 1923  
Untersuchungen zur Lehre von der Blutgerinnung. VIII. Mitt. Über die Gerinnungshemmung durch Neutralsalze  
Biochem. Z. 140:42-62

- 739 Studnický, J. 1969  
Use of flocculants to improve filtering of sugar  
cane juice using MgO  
-Bol. Sobre Deriv. 3(3):3-20
- 740 Suckawa, Takeshi. 1929  
Über den einfluss einiger pharmaka auf die  
magnesiumhyperglykämie und -glykosurie  
Mitt. Med. Akad. Kyoto 3:225-288
- 741 Sugiura, Kanematsu, and Stanley R. Benedict. 1923  
Der einfluss anorganischer salze auf das wachstum  
der geschwulste bei weissen ratten  
J. Cancer Res. 7:324-369
- 742 Sumi, Kenshi. 1931  
Über die beziehung zwischen der wirkung  
verschiedener pharmaka auf den darme  
vom hühnerembryo und den entwicklungsstufen des  
narkose. IV, V und VI. Mitt. (Jap.)  
Folia Pharmacol. Jap. 13(1):132-148
- \* 743 Sunderman, F.W., and J.E. Carroll. 1965  
Measurements of serum calcium and magnesium by  
atomic absorption spectrophotometry  
Am. J. Clin. Path. 43(4):302-310
- 744 Svendsen, A. 1943  
Magnesium sulfate injections in therapy of  
encephalopathy due to hypertension in children  
Ugeskr. Laeg. 105:769
- 745 Svendsen, A. 1945  
Magnesium sulfate in treatment of acute  
glomerulonephritis  
Nord. Med. (Hosp.) 26:809-814
- 746 Swingle, W.W., and A.J. Swingle. 1967  
Activation, inhibition, and reversal of  
inhibition of plasma volume changes  
in adrenalectomized and intact dogs  
Proc. Soc. Exp. Biol. Med. 125(3):815-818
- 747 Syozi, K. 1937  
Epinephrine load in suprarenal glands of rabbits  
after administration of ammonium chloride and  
magnesium sulphate  
Tohoku J. Exp. Med. 30:259-265
- 748 Szekacs, Bela, Istvan Szathmari, Tamas Toth,  
Istvan Szelenyi, and Janos Rigo. 1969  
Effect of magnesium on the renin sensitivity of  
rats (Hung.)  
Kiserl. Orvostud. 21(3):225-228
- 749 Szelenyi, I., E. Nemesanszky, I. Perenczi, and J.  
Rigo. 1967  
Effect of aspartic acid salts on the vascular  
wall in vitro and in vivo (Ger.)  
Arzneim.-Forsch. 17(4):502-505
- 750 Szelenyi, Istvan, Elemer Nemesanszky, Istvan  
Perenczy, and Janos Rigo. 1967  
Effect of Mg salts on vascular tonus (Hung.)  
Kiserl. Orvostud. 19(4):377-380
- 751 Szelenyi, Istvan, Janos Rigo, Elemer Nemesanszky,  
Jozsef Sos, and Margit Jona. 1969  
Effect of magnesium salts on cardiac function in  
vivo (Hung.)  
Kiserl. Orvostud. 21(6):622-626
- 752 Tabor, Herbert, and Sanford M. Rosenthal. 1945  
Experimental chemotherapy of burns and shock.  
VIII. 1. Effects of K administration, of Na  
loss, and fluid loss in tourniquet shock  
U.S. Pub. Health Repts. 60:373-381
- 753 Taketa, H. 1929  
Diuretic action of ions  
Tohoku J. Exp. Med. 12:366-388
- 754 Tadini, R. 1942-1943  
Gli agenti chimici che determinano la ciclopia  
negli anfibi. (Chemical agents inducing cyclopia  
in amphibians)  
R. Ist. Lomb. Sci. Lett. Cl. Sci. 76:115-120
- 755 Tanura, Toyoyuki, Eiichiro Miyazawa, and Shinji  
Miyasawa. 1968  
Toxicity of drugs. V. Subchronic toxic tests on  
mice with the use of amber acid (succinic acid)  
and magnesium phosphate drugs  
J. Nippon Univ. Sch. Dent. 10(3):83-90
- 756 Tanaka, J. 1931  
Effect of magnesium sulphate on phosphorus  
metabolism in blood and muscle of rabbits  
Kyoto-Ikadaigaku-Zasshi 5:77-163
- 757 Tanaka, J. 1932  
Effect of magnesium sulphate and calcium chloride  
on phosphate metabolism in toad muscle  
Mitt. Med. Akad. Kyoto 6:2481-2513
- 758 Tanaka, Shutaro, Ichiro Kato, Tsuneo Iwasura, and  
Torii Sorimachi. 1953  
The buffer ability of sake. I  
J. Soc. Brew. (Japan) 48:24-26
- 759 Tannenbaum, A., and H. Silverstone. 1953  
Effects of varying the level of mineral in the  
diet  
Cancer Res. 13(6):460-463
- 760 Tarbell, H.A. 1937  
Sulfanilamide must be used with care  
J. Med. Soc. N. J. 34:506-507
- 761 Tauber, Henry. 1938  
Carboxylase enzyme system  
J. Biol. Chem. 125:191-199
- \* 762 Teloh, H.A. 1958  
Estimation of magnesium in serum by means of  
flame spectrophotometry  
Am. J. Clin. Path. 30:129-132
- 763 Terauchi, Kojiro. 1929  
Über die sogenannte curarewirkung verschiedener  
substanzen  
Tohoku J. Exp. Med. 12:505-513
- 764 Terhorst, B., and H. Melchior. 1969  
Periodic prophylaxis of urolithiasis  
Z. Urol. Nephrol. 62(10):761-773
- 765 Tezner, Otto. 1929  
Der einfluss verschiedener salze auf die  
retraktion des blutgerinnsels  
Z. Ges. Exp. Med. 64(3/4):462-469
- \* 766 Thatcher, Harvey S. 1928  
A case of magnesium sulphate poisoning  
J. Am. Med. Assoc. 91(16):1185-1186
- 767 Thomas, J.W., and B.S. Emery. 1969  
Effects of sodium bicarbonate, magnesium oxide,  
and calcium hydroxide on milk fat secretion  
J. Dairy Sci. 52(1):60-63
- 768 Thomas, John William, and Roy S. Emery. 1969  
Additive nature of sodium bicarbonate and  
magnesium oxide on milk fat concentrations of  
milking cows fed restricted-roughage rations  
J. Dairy Sci. 52(11):1762-1769
- 769 Thomas, L., and L. Kreitmman. 1931  
Chemotherapy of epithelioma in salmon with  
halogen salts of magnesium  
Bull. Acad. Med., Paris 106:124-126
- 770 Thompson, J. 1968  
Lard rearrangement  
U.S. Pat. 3,392,031 issued in 1968
- 771 Thurnher, H., and E. Kresbach. 1961  
Beitrag zur klinik und behandlung der  
magnesiumsulfatvergiftung. (Contribution to the  
clinical aspects and treatment of magnesium  
sulfate poisoning)  
Wien. Klin. Wochschr. 73(20):356-357
- 772 Timofeeva, T.A. 1947  
Influence of magnesium sulfate on the higher  
nervous activity of dogs  
Akad. Med. Nauk SSSR 1:159

- 773 Tisdall, Frederick P., T.G.H. Drake, and Alan Brown. 1926  
The production of a lowered carbohydrate tolerance in dogs  
Am. J. Dis. Child. 32(6):854-861
- 774 Toi, Bunichi, Masahiro Takahashi, Takeshi Ogasawara, Tetsuo Hino, and Noriko Matsuda. 1960  
Production of dense soy sauce  
Nippon Nogei Kagaku Kaishi 34:1017-1022
- 775 Trabucchi, E. 1932  
Effect of magnesium sulphate; experimental study  
Arch. Int. Pharmacodyn. Ther. 42:65-86
- 776 Trabucchi, Emilio. 1931  
Variazioni di tossiata ottenute per alcuni farmaci del respiro dopo il taglio dei vaghi. (Modifications of toxicity of drugs acting on respiration, following vagotomy)  
Boll. Soc. Ital. Biol. Sper. 6(1):49-52
- 777 Tudoranu, G., C.C. Dimitriu, and M. Vasiliu. 1939  
Influence of rectal absorption on the motor reactions of the human stomach (viscerography) (French)  
Bull. Acad. Med. Roum. 4:315-319
- 778 Tudoranu, G., C.C. Dimitriu, L. Wasserman, and R. Negrutzi. 1939  
Study of the influence of rectally absorbed materials on the chemistry of human gastric contents (French)  
Bull. Acad. Med. Roum. 4:112-123
- 779 Uchida, Yasumi, Kazuaki, Kamisaka, and Hideo Ueda. 1969  
Responses of the cardiac sympathetic receptors to various substances  
Jap. Heart J. 10(3):225-242
- 780 Umeda, Susura, and Seit Kitasima. 1939-1940  
Studies on the permeability of the peripheral vessels  
Jippon J. Clin. Angio-Cardiol. 3:373-378, 432-436, 497-501, 552-556
- 781 Umezawa, Shun-ichi. 1958  
Role of intracellular phosphate ions on contraction and coagulum formation of an isolated frog muscle fibre  
Annot. Zool. Jap. 31(3):127-135
- 782 Underhill, Frank P., and Edward C. Wood. No date  
Die einwirkung von magnesiumpulver auf den strychninkrampf  
J. Pharmacol. Exp. Ther. 36:129-153
- 783 Utescher, Ernst. 1920  
Verfahren zum konservieren von eiern  
D.R. Pat. 328,423 issued Oct. 23, 1920
- 784 V. Parkas, G. 1930  
Nierenstudien. II. Über die wirkung harntfahiger substanzen und deren konzentration auf die quellung der nierezelle  
Z. Ges. Exp. Med. 70:708-719
- 785 Vacek, Tomas. 1928  
O znenach v cinnosti obvodoveho nervstva vlivem koloidniho hydroxydu horecnateho. (On influence of colloidal magnesium hydrate on the function of the nerve-fibers)  
Biol. Spisy. Vys. Skoly Zverolek. Brno 7:87-92
- 786 Vadrot, P.C. 1970  
Production of aromatic substances from animal and vegetable substances  
Indian Pat. 121,290 issued 1970
- 787 Vainshtein, S.G., D.Kh. Maksudova, and M.I. Pivikova. 1970  
Klinicheskoe izuchenie ksilita kak zhelchegonnogo sredstva. (Clinical study of Xylite as a choleric drug)  
Klin. Med. 48(1):126-128
- 788 Van der Pol, C. 1954  
Magnesium oxide as a clarifying agent for raw sugar solutions  
Commun. Sugar Milling Res. Inst. 1954(21)
- 789 Van Dellen, T.R., and J. Roscoe Miller. 1939  
Electrocardiographic changes following the intravenous administration of magnesium sulfate. II. An experimental study on dogs  
J. Lab. Clin. Med. 24:840-843
- \* 790 Van Fossen, D.D., E.E. Baird, and G.S. Tekell. 1959  
A simplified flame spectrophotometric method for estimation of magnesium in serum  
Am. J. Clin. Path. 31(4):368-370
- 791 Van Leeuwen, Gerald. 1965  
Magnesium sulfate for respiratory distress  
Pediatrics 35(4):719
- 792 Van Liere, E.J., D.W. Northup, J.C. Stickney, and R.E. Richard. 1947  
Effect of anemic anoxia on absorption of isotonic magnesium sulfate from the small intestine  
Proc. Soc. Exp. Biol. Med. 64(1):62-64
- 793 Van Sonsbeek, J.J.M. 1969  
In vitro studies of antacids  
Pharm. Weekbl. 104(27):825-831
- \* 794 Van't Riet, Bart, and James E. Wynn. 1969  
Potentiometric determinations of calcium, magnesium, and complexing agents in water and biological fluids  
Anal. Chem. 41:158-162
- 795 Vasil'eva, V.F., G.P. Gusev, Yu. V. Matochin, and E.I. Shakhmatova. 1969  
Vydelenie elektrolitov pochkoi stavridy i skorpeny posle inektsii khloridov Na, K, Ca ili Mg. (Elimination of electrolytes by the kidney in horse-mackerel and rock perch after injection of Na, K, Ca or Mg chlorides)  
Vopr. Ikhtiol. 9(3):546-555
- 796 Vasilenko, D.Ya., A.I. Vertiichuk, V.G. Kebko, and I.V. Povidzen. 1970  
Effect of carboxylation stimulation on carbohydrate and fat metabolism in lactating cows  
Visn. Sil's'kogospod. Nauki 13(1):90-93
- 797 Vavrincz, Gabor. 1969  
Formation and composition of beet molasses. I. Changes in the optical rotatory power of sucrose caused by divalent halides  
Z. Zuckerind. 19(11):607-610
- 798 Vazifdar, N.F. 1930  
Epsom salts poisoning  
Indian Med. Rec. 50:301-304
- 799 Vedove, Angel Delle, and Moises Sloer. 1943  
El tratamiento de la hipertension arterial por el sulfato de magnesio y su accion sobre el electrocardiograma  
Rev. Med. Rosario 33(7):664-670
- 800 Vignes, E.C. 1968  
Effect of temperature on rate of reaction during liming (of sugar cane juice)  
Mauritius Sugar Ind. Res. Inst., Annu. Rep. 1968:117-120
- 801 Vinogradov, V.S. 1939  
A method for the prevention of coagulation of animal blood  
Lab. Prakt. (USSR) No. 5:18-19
- 802 Voegtlin, W.L., and A.C. Ivy. 1934  
An investigation concerning certain substances reported to affect the motility of the gall bladder  
Am. J. Dig. Dis. Nutr. 1(3):174-177
- 803 Vogel, J.H. No date  
Chlormagnesium und schwefelsaure magnesia in den taglichen nahrungsmitteln  
Kali 20:181-184



- 804 Volf, V. 1965  
Effect of phosphates, carbonates, and magnesium oxide upon the intestinal absorption of Sr-85 in rats  
*Experientia* 21(10):571-572
- 805 Von Messe, J.H., and D.O. Stephens. 1969  
Kaltwasserlösliche saure Zubereitungen mit guter fließfähigkeit. (Free-flowing, cold-water-soluble drink mixes)  
W. Ger. Pat. 1,517,076 issued in 1969
- 806 Vorontsov, D.S. 1966  
Effects of alkaline and alkaline-earth metal cations on physical electrotone of denuded frog nerve (Russ.)  
*Fiziol. Zh. S.S.S.R.* 52(1):22-28
- 807 Voza, P., and P. Natale. 1930  
Effects of parenteral injections of magnesium sulphate on calcium and potassium content of blood  
*Ann. Obstet. Gynecol.* 52:1111-1119
- 808 Wacker, Leonhard. No date  
Die rolle des magnesiumphosphats bei der ermüdung und totenstarre des muskels  
*Munch. Med. Wchschr.* 74:1222-1223
- \* 809 Wacker, W.E.C., and A.P. Parisi. 1968  
Magnesium metabolism  
*New Engl. J. Med.* 278(12):658-663
- 810 Wacker, W.E.C., and R.J.F. Williams. 1968  
Magnesium/calcium balances and steady states of biological systems  
*J. Theor. Biol.* 20(1):65-78
- \* 811 Wacker, W.E.C., C. Iida, and K. Fuwa. 1964  
Accuracy of determinations of serum magnesium by flame emission and atomic absorption spectrometry  
*Nature* 202:659-661
- 812 Wallis, Craig, and Joseph L. Melnick. 1962  
Magnesium chloride enhancement of cell susceptibility to poliovirus  
*Virology* 16(2):122-132
- 813 Wallis, Craig, Fred Morales, Joycelyn Powell, and Joseph L. Melnick. 1966  
Plaque enhancement of enteroviruses by magnesium chloride, cysteine, and pancreatin  
*J. Bact.* 91(5):1932-1935
- 814 Wallis, Craig, Joseph L. Melnick, and Fred Rapp. 1965  
Different effects of MgCl<sub>2</sub> and MgSO<sub>4</sub> on the thermostability of viruses  
*Virology* 26(4):694-699
- 815 Watanabe, Tamotsu. 1924  
Studien zur physiologie und experimentellen therapie der gallenabsonderung. II. Mitteilung.  
*Z. Ges. Exp. Med.* 40:201-235
- 816 Watson-Williams, E. 1932  
Magnesium sulphate  
*Practitioner* 128:556-557
- 817 Watson, C.B. 1931  
Effect of intravenous magnesium sulphate on blood pressure  
*Brit. Med. J.* 2:1086-1087
- 818 Webb, Watts R., Ross P. Dodds, Metin O. Unal, Armand M. Karow, William A. Cook, and. 1966  
Suspended animation of the heart with metabolic inhibitors: effect of magnesium sulfate or fluoride and adrenochrome in rats  
*Ann. Surg.* 164(3):343-351
- 819 Weiner, Philip D., and Albert M. Pearson. 1969  
Calcium chelators influence some physical and chemical properties of rabbit and pig muscle  
*J. Food Sci.* 34(6):592-596
- 820 Welsh, R.S. 1964  
Polymerizing effect, on the isolated DNA, of the addition of ATP-MgCl<sub>2</sub> to calf thymus nuclei before extraction  
*Int. Congr. Biochem.* 6(1):94
- 821 Wen, S.-F., N.L.H. Wong, and J.H. Dirks. 1971  
Evidence for renal magnesium secretion during magnesium infusions in the dog  
*Am. J. Physiol.* 220(1):33-37
- 822 Wen, Sung-Feng, Raphael L. Evanson, and John H. Dirks. 1970  
Micropuncture study of renal magnesium transport in proximal and distal tubule of the dog  
*Am. J. Physiol.* 219(3):570-576
- 823 Westenberg, J. 1951  
Experiment in fish salting  
Indo-Pac. Fish. Council., Cronulla, N.S. Wales, Proc. 2(Sect. II/III):123-125
- 824 Whelan, Mary. No date  
Die wirkung intravenöser injektion von anorganischen chloriden auf die zusammensetzung von blut und harn  
*J. Biol. Chem.* 63:585-620
- 825 Wierbicki, Eugen, V.R. Cahill, and P.E. Deatherage 1957  
Effects of added sodium chloride, potassium chloride, calcium chloride, magnesium chloride, and citric acid on meat shrinkage at 70° and of added sodium chloride on drip losses after freezing and thawing  
*Food Technol.* 11:74-76
- \* 826 Wies, F.A. 1933  
Effect of magnesium sulphate on the brain of the fetal rat  
*Yale J. Biol. Med.* 5:545-553
- \* 827 Willner, G. 1971  
Toxicologic-chemical findings in a fatal magnesium poisoning following duodenal administration of magnesium sulfate  
*Arch. Toxikol.* 27:129-135
- 828 Winkelstein, A., and P.W. Aschner. 1926  
The mechanism of the flow of bile from the liver into the intestines  
*Am. J. Med. Sci.* 171(1):104-111
- 829 Winkler, Alexander W., Paul K. Smith, and Hebbel E. Hoff. 1942  
Intravenous magnesium sulfate in the treatment of nephritic convulsions in adults  
*J. Clin. Invest.* 21(2):207-216
- 830 Winogradow, A.P. 1927  
Die wirkung von arzneisubstanzen auf die absonderung der galle. (The action of drugs on the secretion of bile)  
*Arch. Exp. Path. Pharmacol.* 126(1/2):17-30
- \* 831 Winter, J.E., and H.G. Barbour. 1928  
Influence of magnesium salts upon the toxicity and antipyretic action of the salicylates  
*Proc. Soc. Exp. Biol. Med.* 25:587-590
- \* 832 Winter, J.E., C.H. Richey, and H.G. Barbour. 1931  
Magnesium oxide as an aid to the antipyretic action of phenacetin in dogs  
*J. Pharm. Exp. Ther.* 38:343-347
- 833 Witter, Robert F., and Ethel Newcomb. 1952  
Roles of magnesium chloride and pH in preservation of fatty acid oxidase activity of (rat liver) mitochondria heated at 30°  
*Fed. Proc.* 11(1):313
- 834 Witter, Robert F., Ethel H. Newcomb, Mary A. Cottone, and Elmer Stotz. 1954  
The effect of magnesium chloride and pH on heat inactivation of the fatty acid oxidase of mitochondria  
*Proc. Soc. Exp. Biol. Med.* 87(2):465-469

- 835 Wittmann, G. 1967  
The influence of 1 mol  $\text{MgCl}_2$  and 2 mol  $\text{NaCl}$  on  
the 50% C-inactivation of virus causing foot and  
mouth-disease  
Zentralbl. Bakt. Parasitenkd. Infektionskr. Hyg.  
Abt. I. Orig. 202(1):133-134
- 836 Woitaschewski, J.B. 1926  
Action of magnesium sulfate  
Arch. Klin. Chir. 141:135-150
- 837 Wojnar, Stefan, Edward Fuzylo, Eugeniusz Butruk,  
and Wiktor B. Szostak. 1967  
Effect of intravenous infusions of magnesium  
sulfate upon triglyceride lipolysis at the peak  
of alimentary lipemia in atherosclerotic patients  
Pol. Tygod. Lek. 22(35):1325-1327
- 838 Wolff, H., and S. Manjean. 1932  
Action on reduced glutathione content of organs  
of dog  
Compt. Rend. Soc. Biol. 111:779-781
- 839 Woodbury, Dixon M., and Virginia D. Davenport.  
1949  
Brain and plasma cations and experimental  
seizures in normal and desoxycorticosterone-  
treated rats  
Am. J. Physiol. 157(2):234-239
- 840 Motherspoon, J.R. 1965  
The evaluation of magnesium oxide in (sugar)  
juice clarification  
Jaw. Assoc. Sugar Technol. J. 26:65-69
- 841 Wright, E. Percy. 1934  
Die anwendung von magnesiumsulfat bei krebs  
Can. Med. Assoc. J. 31:404-406
- 842 Wunsch, L., and M. Kratka. 1965.  
Chelatometrische magnesium bestimmung in plasma  
Clin. Chim. Acta 12:37-45
- 843 Yamamoto, T. 1967  
The effect of ion concentration and pH on the  
thermal stability of a canine adenovirus  
Can. J. Microbiol. 13(9):1139-1149
- 844 Yanagimachi, Ryuzo. 1957  
Some properties of the sperm-activating factor in  
the micropyle area of the herring egg  
Annot. Zool. Jap. 10(3):114-119
- 845 Yanagishima, Naohiko. 1967  
Induction of heritable respiration deficiency in  
yeast by salt solutions  
Plant Cell Physiol. (Tokyo) 8(1):211-215
- 846 Yasukawa, Yutaka. 1933  
Experiments on sea water and *Vibrio cholerae*  
Jap. J. Exp. Med. 11(2):119-127
- 847 Yokoyama, M., and J.S. Finlayson. 1961  
Effects of salt concentration, pH, and certain  
ions on the agglutination of human erythrocytes  
by isohemagglutinins  
Transfusion 1:175-178
- 848 Yurisson, E. 1956  
The mechanism of the action of sodium and  
magnesium sulfates on the motility of the  
intestine of hibernating animals  
Sb. Nauchn. Tr. Est. S-kh. Akad. 1956(2):238-244
- 849 Zajack, Peter. 1968  
Use of cation-exchange resins in the hydrogen  
cycle for the neutralization of filtered cane  
sugar juice treated with magnesium oxide  
Sobre Deriv. Cane Azucar 2(1):30-40
- 850 Zak, B.R. 1941  
The effect of electrolytes on the stimulating  
action of hormones  
J. Physiol. U.S.S.R. 30:656-658
- 851 Zakrivdoroga, S.P. 1945  
Comparative influence and therapeutic range of  
action of some drugs on isolated hearts of cold-  
blooded animals (frogs)  
Parakol. Toksikol. 8(5):10-14
- 852 Zakusov, V.V. 1940  
Changes in the impulse-summation capacity of the  
central nervous system under the influence of  
some narcotics and analgesics  
Parakol. Toksikol. 3(6):4-10
- 853 Zel'tser, Ya.V. 1959  
Solubility of salts, which create hardness in  
vodka  
Spir. Prom. 25(3):28-32
- 854 Zenon, Bubien, Tadeusz Garbulinski, and J.  
Wasilewski. 1969  
Effects of magnesium sulfate and atropine on  
gastrointestinal motility in sheep poisoned with  
the organophosphorus compound Z-50 (Pol.)  
Pol. Arch. Wet. 11(4):571-580
- 855 Zucker, Marjorie B., and Jennie Borrelli. 1958  
Some effects of divalent cations on the clotting  
mechanism and the platelets of EDTA blood  
J. Appl. Physiol. 12(3):453-460
- 856 Zwillinger, L. 1936  
Magnesium sulfuricum bei einer  
strophanthinvergiftung. II. Mitt. Zur  
magnesiumwirkung auf das herz  
Wien. Klin. Wschr. 49:594-595